

# The Geometry of Ellipses and Planetary Orbits

**Moti Ben-Ari**

<http://www.weizmann.ac.il/sci-tea/benari/>

Version 0.12

October 6, 2023

© Moti Ben-Ari 2023

This work is licensed under Attribution-ShareAlike 4.0 International. To view a copy of this license, visit <http://creativecommons.org/licenses/by-sa/4.0/>.

# Contents

<b>1</b>	<b>Introduction</b>	<b>1</b>
<b>2</b>	<b>The sizes of the Earth, Moon and Sun</b>	<b>4</b>
2.1	Eratosthenes's measurement of the radius of the earth . . . . .	4
2.2	Aristarchus's measurements . . . . .	4
<b>3</b>	<b>The Sun-centered solar system</b>	<b>8</b>
3.1	The length of the year and the length of the seasons . . . . .	8
3.2	The location of the center of the Earth's orbit . . . . .	10
<b>4</b>	<b>Elliptical orbits</b>	<b>13</b>
4.1	Determining the radius of the Earth's orbit . . . . .	13
4.2	Measuring the angles in the triangle Sun-Earth-Mars . . . . .	14
4.3	A new location for the center of the Earth's orbit . . . . .	15
4.4	Orbits are ellipses . . . . .	15
<b>5</b>	<b>Gravitation</b>	<b>18</b>
5.1	Newton's laws of motion . . . . .	18
5.2	The inverse square law for gravitation . . . . .	21
5.3	Universal gravitation . . . . .	22
5.4	Kepler's third law . . . . .	23
<b>6</b>	<b>A proof Proposition XI, Problem VI</b>	<b>24</b>
6.1	A formula for $QR$ . . . . .	24
6.2	A formula for $QT$ . . . . .	25
6.3	A formula for $QR/QT^2$ . . . . .	26
6.4	Approaching the limit . . . . .	27
<b>7</b>	<b>Ellipses</b>	<b>28</b>
7.1	Fundamental properties . . . . .	28
7.2	A circle circumscribing an ellipse . . . . .	30
7.3	The latus rectum of an ellipse . . . . .	30
7.4	The area of an ellipse . . . . .	31
7.5	The angles between a tangent and the lines to the foci . . . . .	32
7.6	Conjugate diameters . . . . .	32

7.7	The parametric representation of an ellipse . . . . .	34
7.8	Areas of parallelograms . . . . .	34
<b>8</b>	<b>Ellipses in Euclidean geometry</b>	<b>37</b>
8.1	The definition of an ellipse using the focus and the directrix . . . . .	37
8.2	A right angle at the focus of an ellipse . . . . .	38
8.3	Ratios of perpendiculars to the axes . . . . .	40
8.4	A circle circumscribing an ellipse . . . . .	41
8.5	The latus rectum of an ellipse . . . . .	42
8.6	Areas of parallelograms . . . . .	42
<b>A</b>	<b>Theorems of Euclidean Geometry</b>	<b>48</b>
A.1	Constructing a circle from three points . . . . .	48
A.2	The product of two subsegments . . . . .	48
A.3	Adjacent pairs of similar triangles . . . . .	49
A.4	The angle bisector theorems . . . . .	50
	<b>Sources and further reading</b>	<b>52</b>

# 1 Introduction

Everyone “knows” that Kepler discovered that the orbits of the planets are ellipses and that Newton showed that an elliptical orbit implies that the force of gravity must be inversely proportional to the square of the distance from the Sun. Although I knew these facts, I had never seen them demonstrated.

*Calculus in Context* [5] by Alexander J. Hahn is a comprehensive textbook on introductory calculus that augments theory with applications in physics and astronomy, such as the work of Kepler, Newton and Galileo, as well as applications in engineering such as building bridges and domed structures. These are not just historical anecdotes but detailed computations.

This document contains a detailed explanation of one topic from Hahn’s book: the determination of orbits by Aristarchus, Copernicus, Kepler and Newton. The presentation is mathematical, since the historical and astronomical aspects are thoroughly described in [5], as well as in other works. The document is intended to enrich the learning of mathematics by secondary-school students and students in introductory university courses. The prerequisites are a very good knowledge of Euclidean geometry along with some trigonometry, a bit calculus and Newton’s laws of motions.

Section 2 presents the measurements of the radii and distances of the Earth, Moon and Sun by Eratosthenes and Aristarchus. Section 3 describes the construction of a model of a Sun-centered system by Nicolaus Copernicus. Section 4 shows how Johannes Kepler developed his three laws of planetary motion and Section 5 presents Isaac Newton’s derivation of the inverse-square law of gravitation from of Kepler’s laws. One step of Newton’s derivation requires a theorem whose proof is very long, so it is split off into Section 6. Section 7 contains more than you ever wanted to know about the mathematics of ellipses, but these theorems are necessary. I suggest that you look up each theorem (and its proof) as needed, rather than trying to study them all at once. Section 8 is discussed in the next paragraph. Theorems of Euclidean geometry that are used but do not concern ellipses are collected in Appendix A.

## Euclidean geometry

It is easy to measure angles. We are familiar with the use of a protractor in school and these can be scaled-up to obtain more accurate measurements. Measuring long distances was impossible until the recent inventions of radar and lasers. At most one could pace-off distances with low accuracy. The only *measured* distance used here is the estimate of 800 km by Eratosthenes for the distance between two places in Egypt (Section 2.1). For this reason, the mathematics used is primarily Euclidean geometry, in particular, similar triangles and ratios of their sides.

The final steps in Newton’s derivation requires the use of limits, which had been used already by Archimedes to compute the circumference and area of a circle by approximating the circle. Newton (along with his contemporary Gottfried Wilhelm Leibniz) developed

the calculus from the concept of limits. However, Newton's *Principia* uses Euclidean geometry almost exclusively, although analytic geometry had been developed by René Descartes and Pierre de Fermat even before Newton was born.

Newton expected his readers to have an extensive knowledge of geometry. This expectation continued until relatively recently:

In book 1, prop[osition] 10 (and notably in prop[osition] 11), Newton made use of a property of conics which he presents without proof, merely saying that the result in question comes from “the *Conics*.” Here, as elsewhere in the *Principia*, Newton assumes the reader to be familiar with the principles of conics and of Euclid. In the eighteenth and nineteenth centuries, when Newton's treatise was still being read in British universities, authors of books on “conic sections”—for example, W. H. Besant, W. H. Drew, Isaac Milnes—supplied the proof of this theorem in order to help readers of the *Principia* who might be baffled by the problem of finding a proof. They even chose letters to designate points on the diagrams so that the final result would appear in exactly the same form as in the *Principia* [2, p. 330].

To keep this spirit alive, Section 8 brings the necessary proofs of ellipses using Euclidean geometry. The reader is forewarned, however, that as Euclid said, “There is no royal road to geometry!”


## Style and notation

The computations in Hahn's book are faithful to the historical record, for example, measuring distances in units such as the stadia of the Greeks. Here, the computations are fully modernized and use modern units such as kilometers.

Diagrams are used to facilitate understanding each step in the geometrical proofs, more than appear in other sources. For example, Newton proved his difficult theorem (Section 6) using only a single diagram. The diagrams are not to scale and are often distorted. Otherwise, it would be impossible to draw planetary orbits on a piece of paper.

The following shortcuts facilitate a less verbose presentation:


- $AB$  denotes both a line segment and its length.
- $\triangle ABC$  denotes both a triangle and its area.
- The phrase “ $AB$  intersects  $CD$ ” is used even if  $AB$  needs to be *extended*<sup>1</sup> until it intersects  $CD$ :




---

<sup>1</sup>The term used by Newton is *produced*.

- The notation in the following diagram will be used consistently to refer to elements of an ellipse. The diagram is for reference since the terms will only be defined when needed.
  - $O$  is the center of the ellipse and  $a, b$  are the semi-major and semi-minor axes of an ellipse.  $A, A'$  are the vertices on the major axis and  $B, B'$  are the vertices on the minor axes.
  - $S$  and  $H$  are the foci and  $c$  is the distance from a focus to the center.
  - $X$  is the intersection of the major axis with the directrix  $d$ . (There is a second directrix on the right side of the ellipse.)



## 2 The sizes of the Earth, Moon and Sun

### 2.1 Eratosthenes's measurement of the radius of the earth

The ancient Greeks knew that the Earth is round and Eratosthenes was able to measure the radius of the Earth (Figure 1). Choose two points  $A, B$  on the same longitude and measure the distance  $d$  between them. Plant a vertical stick (red) in the ground at  $A$  and another (blue) at  $B$ . On a day in the year when the stick at  $A$  produces no shadow at noon, at the same time the stick at  $B$  produces a shadow whose angle is  $\alpha$ . The sun is so far away from the Earth that over the relatively short distance  $d$ , the rays of the Sun are essentially parallel. By alternate interior angles, the angle between the two sticks as measured from the center of the Earth is also  $\alpha$ .

The angle that Eratosthenes measured at the blue stick was

$$\alpha = 7.5^\circ \cdot \frac{2\pi}{360} \approx 0.131 \text{ radians},$$

and the distance  $d$  between  $A$  and  $B$  was known to be approximately 800 km. The arc  $\widehat{AB}$  subtends the angle  $\alpha = d/r_e$  where  $r_e$  is the radius of the Earth, so

$$r_e = \frac{d}{\alpha} = \frac{800}{0.131} \approx 6107 \text{ km}. \quad (1)$$

This value is quite close to the modern measurement of 6370 km.

### 2.2 Aristarchus's measurements

Using  $r_e$ , Eratosthenes's measurement of the radius of the Earth, Aristarchus was able to measure and compute the following values:

- $r_m$ : the radius of the Moon,
- $r_s$ : the radius of the Sun,
- $d_m$ : the distance from the Earth to the Moon,
- $d_s$ : the distance from the Earth to the Sun.




Figure 1: Eratosthenes's measurement of the radius of the earth




Figure 2: Observing a first quarter moon

### Computing $d_s/d_m$

An observer on Earth can follow the phases of the Moon as it revolves around the Earth. At one point in the month the phase will be first quarter, meaning that the one half of the moon is illuminated while the other half is not (Figure 2). The angle between the Sun and the Moon will be  $87^\circ$ . Since exactly half of the moon is illuminated, we know that the angle  $\angle$  Earth-Moon-Sun is a right-angle so

$$\begin{aligned}\cos 87^\circ &= \frac{d_m}{d_s} \\ \frac{d_s}{d_m} &= \frac{1}{\cos 87^\circ} \approx 19.\end{aligned}\tag{2}$$

### Computing $r_s/r_m$ and $d_m/r_m$

The Moon is much, much smaller than the Sun, but it is also much, much closer to the Earth. When the Moon is precisely positioned between the Earth and the Sun, its “disk” exactly covers the “disk” of the Sun, causing a total solar eclipse (Figure 3).




Figure 3: A solar eclipse



The angle subtended by the Moon is  $2^\circ$  degrees. Bisecting the angle creates two right triangles with an acute angle of  $1^\circ$ , where the right angles are the tangents to Moon and the Sun. By similar triangles, Equation 2 and Figure 3,

$$\frac{r_s}{r_m} = \frac{d_s}{d_m} = 19 \quad (3)$$

$$\frac{d_m}{r_m} = \frac{1}{r_m/d_m} = \frac{1}{\sin 1^\circ} \approx 57. \quad (4)$$

### Computing the radii and distances

Figure 4 shows a lunar eclipse. Unlike a solar eclipse where the Moon exactly covers the Sun, the Earth more than covers the Moon and its shadow is four times the Moon's radius.

Figure 5 show a lunar eclipse annotated with the distances  $d_m, d_s$  and the radii  $r_m, r_e, r_s$ . The ray from the top of the Sun is tangent to both the Sun and the Earth, so it forms right angles with their radii, as well as with the extension of the Moon's radius. The thick horizontal lines are constructed parallel to the line connecting the centers, forming two similar right triangles, so using Equation 3,

$$\begin{aligned} \frac{r_s - r_e}{r_e - 2r_m} &= \frac{d_s}{d_m} = \frac{r_s}{r_m} \\ r_s r_e + r_m r_e &= 3r_s r_m. \end{aligned}$$

Again from Equation 3,  $r_s = 19r_m$ , so

$$r_m = \frac{20}{57} r_e.$$

By Equation 1,  $r_e \approx 6107$  km, by Equation 4,  $d_m = 57r_m$ , and by Equation 2,  $d_s = 19d_m$ , so we can compute the radii and distances:

$$r_m = \frac{20}{57} r_e \approx 2143 \text{ km}$$

$$r_s = 19r_m \approx 40,713 \text{ km}$$

$$d_m = 57r_m \approx 122,140 \text{ km}$$

$$d_s = 19d_m \approx 2,320,660 \text{ km}.$$




Figure 4: A lunar eclipse




Figure 5: Detail of a lunar eclipse

The following table summarizes these data together with the modern values [5, Table 1.3]. While the computed values for the radii of the Earth and the Moon are not far off from the modern values, the other computed values are not anywhere near the modern values. Nevertheless, they do show that the Greeks understood the immense size of the solar system.

		Computed (km)	Modern (km)
$r_e$	radius of Earth	6107	6370
$r_m$	radius of Moon	2143	1740
$r_s$	radius of Sun	40,713	695,500
$d_m$	distance Earth-Moon	122,140	384,570
$d_s$	distance Earth-Sun	2,320,660	150,000,000

### 3 The Sun-centered solar system

As everyone living far from equator knows, the time between sunrise and sunset varies with the seasons. The reason is that the axis of the rotation of the Earth is offset by  $23.5^\circ$  relative to the orbit of the Earth. The plane of the orbit of the Earth around the Sun is called the *ecliptic*. Measuring the length of the day as the time from sunrise to sunset, there is a day in June, called the *summer solstice*, when the length of the day is longest. Similarly, there is a day in December, called the *winter solstice*, when the length of the day is shortest.<sup>2</sup> There are also two days when the length of the day equals the length of the night: the *autumn equinox* in September and the *spring equinox* in March.

Today we know that the universe is immensely large and that the stars are moving at extremely high speeds, but an observer on Earth sees them as if their positions are fixed on a sphere around the earth, called the *celestial sphere*. This solstices and equinoxes can be associated with the projection of the Sun on the celestial sphere as seen from the Earth. The details of the Earth's orbit can be found in books on astronomy, as well as in the Wikipedia articles on *Equinox* and *Solstice*.

#### 3.1 The length of the year and the length of the seasons

Let us assume that the Earth orbits the Sun in a circle, such that the center of the orbit  $O$  is the center of the Sun  $S$ . In Figure 6 the inner circle is orbit of the Earth and the outer circle is the celestial sphere. The orbit can be divided into four quadrants called *seasons*: spring, summer, autumn, winter.

---

<sup>2</sup>This holds for the northern hemisphere; in the southern hemisphere the opposite holds.




Figure 6: The orbit of the Earth and the seasons





Figure 7: The lengths of the seasons are not equal

The length of a year is approximately  $365\frac{1}{4}$  days. The extra  $\frac{1}{4}$  day is accounted for by adding a day in leap years.<sup>3</sup> The length of each season as determined by the equinoxes and the solstices is  $365.25/4 = 91.3125$  days. However, measurements by the Greek astronomer Hipparchus showed that the actual lengths of the seasons differed from this number and a model of the solar system must be able to explain these differences:

Season	Days	%
Spring	$94\frac{1}{4}$	25.8
Summer	$92\frac{1}{2}$	25.3
Autumn	$88\frac{1}{8}$	24.1
Winter	$90\frac{1}{8}$	24.7

In his Earth-centered solar system, Hipparchus proposed that the center of the Earth's orbit is offset from the center of the Sun. Copernicus used the same idea in his Sun-centered solar system (Figure 7). If the center of the Earth's orbit is in the upper-left quadrant of the coordinate system defined by the equinoxes and the solstices, the angles for spring and summer are obtuse, so the seasons are longer than one-fourth of a year, whereas the angles for the autumn and winter are acute, so they are shorter than one-fourth of a year. Figure 8 shows a magnified and distorted view of Figure 7. It has been annotated with additional lines and labels that will facilitate the demonstration of Copernicus's computation. The axes  $A'C'$  and  $B'D'$  have their origin  $O$  at the center of the Earth's orbit and are parallel to the axes in the ecliptic. The dashed lines from  $O$  are all radii of the Earth's orbit that will be denoted  $r$ . The dotted right triangles will be used in the computation.

<sup>3</sup>The length of a year is actually 365.2425. In the sixteenth century, the *Gregorian calendar* accounted for the difference by removing three leap years in every four hundred years.



Copernicus's task was to locate the position of the center of the Earth's orbit  $O$  relative to the center of the Sun  $S$ . This will be given in polar coordinates  $OS = c$  and  $\angle FSB = \lambda$  (the label is on the large circle of the ecliptic). The strategy of the computation is as follows:

- ### Computing the angles of the arcs $\widehat{AB}, \widehat{AC}$

$$\widehat{AB} = 2\pi \cdot \frac{92\frac{51}{60}}{365.25} = 2\pi \cdot \frac{92.85}{365.25} = 1.5972 \text{ radians}$$

$$\widehat{AC} = 2\pi \cdot \frac{92\frac{51}{60} + 93\frac{14.5}{60}}{365.25} = 2\pi \cdot \frac{186.09}{365.25} = 3.2012 \text{ radians}.$$

### Computing the angles of the arcs $\widehat{AA'}$ , $\widehat{BB'}$

Let us express the arcs  $\widehat{AC}$  and  $\widehat{AB}$  in terms of the arcs that comprise them. Since  $AC$  is parallel to  $A'C'$ ,  $\widehat{AA'} = \widehat{C'C}$ . Compute  $\widehat{AA'}$ :

$$\widehat{AC} = \widehat{AA'} + \widehat{A'C'} + \widehat{C'C} = 2\widehat{AA'} + \pi$$

$$\widehat{AA'} = \frac{1}{2}(3.2012 - \pi) = 0.0298 \text{ radians}.$$

Now that we have computed  $\widehat{AB}$  and  $\widehat{AA'}$  we can compute  $\widehat{BB'}$ :

$$\widehat{AB} = \widehat{AA'} + \widehat{A'B'} - \widehat{BB'}$$

$$\widehat{BB'} = 0.0298 + \frac{\pi}{2} - 1.5927 = 0.0034 \text{ radians}.$$

### Computing the lengths of the arcs $\widehat{AA'}$ , $\widehat{BB'}$

$OA$  and  $OB$  are radii of the Earth's orbit so

$$\sin \alpha = \frac{a}{r} \approx \alpha$$

$$a \approx r\alpha = r\widehat{AA'} = 0.0298r$$

$$\sin \beta = \frac{b}{r} \approx \beta$$

$$b \approx r\beta = r\widehat{BB'} = 0.0034r,$$

where we have used the assumption that  $O$ , the center of the Earth's orbit, is very close to the Sun  $S$  so that  $\lim_{\alpha \rightarrow 0} \frac{\sin \alpha}{\alpha} = \lim_{\beta \rightarrow 0} \frac{\sin \beta}{\beta} = 1$ .

### Computing the position of $O$ relative to $S$

Figure 9 shows a magnified diagram of a portion of Figure 8. In the dotted triangles, we have already computed the lengths  $a$  and  $b$ . Since  $OT$  is parallel to  $A'A$  and  $TS$  is parallel to  $BB'$ , we can label  $OT$  by  $a$  and  $TS$  by  $b$ . The emphasized triangle is a right triangle and  $c$ , the distance of  $O$  from  $S$ , can be obtained from Pythagoras's theorem:

$$c = \sqrt{a^2 + b^2} = r\sqrt{(0.0298)^2 + (0.0034)^2} = 0.03r.$$

$\lambda$  can be obtained from trigonometry:

$$\lambda = \tan^{-1} \frac{b}{a} = \tan^{-1} \frac{0.0034}{0.0298} = 0.1129 \text{ radians} \approx 6.47^\circ.$$

The distance  $0.03r$  is shown in the following table using the values of  $r$  from the table on page 7.

12

## 4 Elliptical orbits

Towards the end of the sixteenth century, the astronomer Tycho Brahe carried out extremely precise observations. In 1600 he hired Johannes Kepler as his assistant and when Tycho died soon afterwards, Kepler was appointed to his position. Here we explain how Kepler was able to establish that planetary orbits are ellipses.

### 4.1 Determining the radius of the Earth's orbit

A Martian year is 687 days, that is, it equals  $\frac{687}{365.25} = 1.88$  Earth years. We know when Mars reaches a “new year” by observing its projection on the celestial sphere, but each time the position of the Earth in its orbit will be different. Figure 10 shows the orbit of the Earth—its center  $O$  offset from the Sun  $S$  as Copernicus showed—at four occasions when the position of Mars  $M$  at its new year was observed. Four triangles are created  $\triangle OE_iM$ . Figure 11 shows one of the triangles with the angles labeled. Using the law of sines,

$$\frac{OE_i}{\sin \beta} = \frac{OM}{\sin \alpha}$$

$$OE_i = OM \frac{\sin \beta}{\sin \alpha}.$$

Tycho Brahe was able to measure all three angles:<sup>4</sup>

	$\alpha$	$\beta$	$\gamma$	$OE_i$
$E_1$	127.1	20.8	32.1	$0.6682 \cdot OM$
$E_2$	84.2	35.8	60.5	$0.6721 \cdot OM$
$E_3$	41.3	42.4	96.4	$0.6785 \cdot OM$
$E_4$	1.6	3.4	175.0	$0.6805 \cdot OM$

<sup>4</sup>The values are rounded to one decimal point.




Figure 10: Observations of the orbit of Mars from the Earth






Figure 11: One triangle Earth-Sun-Moon

The values of  $OE_i$  given in the fourth column of the table are not equal. Assuming (as everyone did at that time) that the Earth's orbit is circular, the only solution was to move the center of the orbit so that  $\{E_1, E_2, E_3, E_4\}$  were all on the circle.

## 4.2 Measuring the angles in the triangle Sun-Earth-Mars

How can the angles  $\alpha, \beta, \gamma$  be measured? Since  $\triangle E_iOM$  is a triangle, it is sufficient to measure two of the angles.  $\gamma$  is easily measured by observing Mars and the Sun at the same time. However, neither  $\alpha$  nor  $\beta$  can be measured directly since they are not accessible to an observer on Earth.

Tycho's measurement used the known periods of the orbits to compute the angles. The Earth moves counterclockwise around Sun. Given any point  $E$ , for some  $t$ ,  $t$  days later the Earth will have moved to  $E'$  and Mars will have moved to  $M'$ , so that they are in *opposition*, that is, Mars will be on the continuation of the Earth-Sun line (Figure 12). Since the Earth completes an orbit in about half the time that Mars takes to complete an orbit,  $t$  will be such that neither the Earth nor Mars has completed a full orbit. The angles  $\theta_E$  and  $\theta_M$  are fractions of a circular orbit of  $360^\circ$ , so

$$\begin{aligned}\frac{t}{365.25} &= \frac{\theta_E}{360} \\ \theta_E &= \frac{360}{365.25} t \\ \frac{t}{687} &= \frac{\theta_M}{360} \\ \theta_M &= \frac{360}{687} t.\end{aligned}$$

This gives values for  $\theta_E$  and  $\theta_M$ . Since  $E'M'$  is a straight line, we have that  $\alpha - \theta_M = 180^\circ - \theta_E$ , so that  $\alpha = 180 - \theta_E + \theta_M$ , and the values of  $OE_i$  can be computed.




Figure 12: The Earth and Mars in opposition

### 4.3 A new location for the center of the Earth's orbit

Kepler's next task was to obtain a new value  $O'$  for the center of the Earth's orbit such that the  $E_i$ 's are on the orbit. Given the new locations of the Earth  $\{E_1, E_2, E_3, E_4\}$ , by Theorem A.1 a circle centered at  $O'$  can be constructed that goes through  $\{E_1, E_2, E_3\}$  (Figure 13). To verify that this is the correct orbit, check that  $E_4$  is on the circle.

### 4.4 Orbits are ellipses

While Kepler was able to modify the center of the orbit of the Earth to be consistent with the observations, he was not able to adequately describe the orbit of Mars. After years of work, he came to the conclusion that the orbit must be oval like an egg. Oval, perhaps, but certainly not an ellipse, because he was certain that it would have been discovered by




Figure 13: Observations of the orbit of Mars from the new Earth's orbit




Figure 14: The orbit of Mars as an oval “egg”

Archimedes! Figure 14 shows  $C$ , a position on a circular orbit, and an oval orbit (dashed), where  $M$  is the position of Mars on the oval corresponding to  $C$ . The radius of the circular orbit is labeled  $a$  and the unknown distances to  $M$  are labeled  $s = SM$  and  $t = OM$ .

Kepler's computed that  $\frac{a-t}{t} = 0.00429$  and  $\frac{s}{t} = 1.00429$ , so that

$$\begin{aligned}\frac{a-t}{t} &= \frac{s}{t} - 1 \\ a-t &= s-t,\end{aligned}$$

and therefore  $SM = s = a = AO = CO$ . The dashed oval is likely an ellipse, because in an ellipse  $SM = AO$  (by Theorem 7.2). Kepler then computed the projections of the observations of Mars on the  $x$ -axis (Figure 15) and obtained for all of them that

$$\frac{M_i O_i}{C_i O_i} = \frac{1}{1.00429} = 0.99573.$$

By Theorem 7.4, since the ratio  $MO/CO = b/a$  is constant for in an ellipse, Kepler was able to conclude that the orbit of Mars is an ellipse.




Figure 15: The orbit of Mars as an ellipse

## 5 Gravitation

In the *Principia* Isaac Newton proved the following theorem led to the theory of universal gravitation.

**Theorem 5.1** *If a planet subject to a centripetal force follows an elliptical orbit around the Sun, then the force decreases as the inverse square of the distance from the Sun.*

After a review of Newton's Laws of force and motion, we show that Kepler's second law must hold in *any* system subject to a centripetal force. The next step is to show the inverse square law and then it is a small step to universal gravitation and Kepler's third law.

### 5.1 Newton's laws of motion

1. A body in uniform motion (including a body at rest) continues with the same motion unless a force is applied.
2. A force  $F$  applied to a body causes an acceleration  $a$  in the direction of the force whose magnitude  $a = F/m$ , where  $m$ , the constant of proportionality, is called the *mass* of the body.
3. If one body exerts a force on a second body, the second body exerts a force on the first of equal magnitude but in the opposite direction.

Forces are denoted by vectors, where the direction of the vector represents the direction of the force and the length of the vector represents the magnitude of the force. Forces can be decomposed into perpendicular components (Figure 16), or into components in any directions (Figure 17). The components form a parallelogram whose diagonal is the resultant force.

Newton was interested in *centripetal force* which is a force exerted by a single body on another, in particular, the gravitational force exerted by the Sun on a planet (Figure 18). Since the only force is that directed towards the Sun, the planet does not move "up" or "down" so its orbit is in a plane.




Figure 16: Perpendicular components of a force




Figure 17: Arbitrary components of a force




Figure 18: Centripetal force




Figure 19: Equal areas in equal times

Kepler's second law states that a planet in orbit sweeps out equal area in intervals of equal duration, that is, if it takes the planet time  $t$  to move from  $P_1$  to  $P_2$  and also  $t$  to move from  $P_3$  to  $P_4$ , then the area of the sector  $P_1SP_2$  is equal to the area of  $P_3SP_4$  (Figure 19). (Obviously, this means that the speed of the planet must vary as it traverses its orbit  $v_{P_1P_2} \gg v_{P_3P_4}$ .) Newton proved that this must be true in any system where a body is subject to a centripetal force from another body.

The proof is based on dividing an area into very small sectors and then taking the limit. Consider three points  $A, B, C$  on the orbit (Figure 20) that represent the positions of the planet at intervals of  $\Delta t$ . For clarity we have drawn them spaced out, but the intention is that they are very close together. Newton assumed that the planet does not smoothly traverse the arcs, but rather that it every  $\Delta t$  it jumps in discrete steps from one point on the orbit to the next.

Figure 21 shows how the force is exerted in discrete steps. The planet moves from  $A$  to  $B$  and we expect that the centripetal force at  $B$  will cause an acceleration that moves the planet to  $C$ , the next point on the orbit. Instead, we "pretend" that the force is not applied at  $B$ , but, in the absence of an applied force, planet continues to move in the same direction and at the same speed. After another period of  $\Delta t$  as passed and the planet has reached point  $C'$ , the force is now applied *in the same direction* as it would have been applied at  $B$ , moving the planet to  $C$ .

**Theorem 5.2** *The area of  $\triangle ASB$  is equal to the area of  $\triangle BSC$ .*

**Proof** The proof will be done in two steps by showing that  $\triangle ASB = \triangle BSC'$  and then that  $\triangle BSC' = \triangle BSC$ .

- In Figure 22,  $\triangle ASB$  is shown in blue and  $\triangle BSC'$  is shown in red. It is assumed that  $AB = BC'$  (the planet moves from  $B$  to  $C'$  during the same interval  $\Delta t$ ), so since  $SH$  is the height of both triangles, their areas are equal.
- In Figure 23,  $\triangle BSC$  is shown in blue and  $\triangle BSC'$  is shown in red. It is assumed  $CC'$  is parallel to  $SB$  (the planet is subject to the centripetal force at  $C'$  in the *same*




Figure 20: “Small” sectors of an orbit




Figure 21: Exerting force at discrete times

direction as the force at B), so the heights of both triangles to the common side  $SB$  are equal and their areas are equal. It follows that  $\triangle ASB = \triangle BSC' = \triangle BSC$ , which we denote by  $\Delta A$ . ■

We assume that the sectors of the orbit are each divided up into small sectors of uniform duration  $\Delta t$ . By the theorem, each sector has the same area  $\Delta A$ . Therefore (see Figure 19),

$$\frac{A_{P_1SP_2}}{\Delta A} = \frac{t}{\Delta t} = \frac{A_{P_3SP_4}}{\Delta A},$$

from which Kepler’s second law follows:  $A_{P_1SP_2} = A_{P_3SP_4}$ .

The proof used two approximations:

- $\Delta A$  is an approximation of the area of each sector.
- The force at  $C'$  is an approximation to the force at  $B$ .




Figure 22:  $\triangle ASB = \triangle BSC'$




Figure 23:  $\triangle BSC' = \triangle BSC$




Figure 24: The derivation of the inverse square law

In the limit as the size of the sectors decreases, the errors become negligible.

**Definition 5.3** For a given elliptical orbit,  $\kappa = \frac{A}{t}$ , where  $A$  is the area of the ellipse and  $t$  is the period of the orbit, is called Kepler's constant.

## 5.2 The inverse square law for gravitation

Newton's next step was to show that if the orbit of a planet is elliptical, the centripetal force must be proportional to the mass of the planet and inversely proportional to the square of its distance from the Sun. In Figure 24,  $S$  is the Sun, and  $P$  and  $Q$  are points on the orbit that are very close to each other.  $PR$  is the tangent to the ellipse at  $P$ , and  $R$  is chosen so that  $QR$  is parallel to  $SP$ .  $QT$  is constructed perpendicular to  $SP$ . Denote the lengths  $r_p = SP$ ,  $h = QT$ ,  $q = QR$  and the time interval from  $P$  to  $Q$  by  $\Delta t$ .

When a point is subject to an acceleration  $a$  for a period of  $\Delta t$ , its displacement is  $\frac{1}{2}a(\Delta t)^2$ . From Newton's second law we know that at point  $R$ , the planet is subjected to an acceleration of  $F_P/m$ , so

$$q = \frac{1}{2} \frac{F_P}{m} (\Delta t)^2$$

$$F_P = \frac{2mq}{(\Delta t)^2}.$$

Now we compute the area of  $\Delta A_{PSQ} \approx \Delta SPQ = (1/2)hr_p$  and use Kepler's constant:

$$\Delta t = \frac{\Delta A_{PSQ}}{\kappa} = \frac{hr_p}{2\kappa}$$

$$F_P = 2mq \cdot \frac{4\kappa^2}{(hr_p)^2} = 8\kappa^2 m \cdot \frac{q}{h^2} \cdot \frac{1}{r_p^2}.$$

To obtain an inverse-square law for the force, the first two factors have to be independent of the distance. For a given planet  $m$  is constant and for a given elliptical orbit  $\kappa$  is constant, so the first factor does not depend on the distance. What about the second factor  $q/h^2$ , in particular, what value does it have as  $\Delta t$  approaches zero?



**Theorem 5.4** *In an elliptical orbit*

$$\lim_{\Delta t \rightarrow 0} \frac{q}{h^2} = \frac{1}{L},$$

where  $L$  is the length of the latus rectum of the ellipse (Definition 7.5).

Newton's proof is very complex and is presented separately in Section 6.

Since  $L$  is constant for any given ellipse, the inverse square law can be written

$$F_P = \frac{8\kappa^2 m}{L} \cdot \frac{1}{r_P^2}. \quad (5)$$

The formula can be re-written so that the constant values appearing are more familiar:  $a$ , the semi-major axis and  $T$ , the period of the orbit. By Theorem 7.6,  $L=2b^2/a$  and by Theorem 7.7,  $\kappa = A_{\text{ellipse}}/T=\pi ab/T$ , so

$$F_P = \frac{8\kappa^2 m}{L} \cdot \frac{1}{r_P^2} = \frac{8(\pi ab)^2 m}{T^2} \cdot \frac{a}{2b^2} \cdot \frac{1}{r_P^2} = \frac{4\pi^2 a^3 m}{T^2} \cdot \frac{1}{r_P^2}. \quad (6)$$

Newton was able to show that:

- The inverse square law applies to all conic sections including a parabola and a hyperbola, not just to an ellipse and, of course, to a circle. Some comets have hyperbolic orbits and orbit the Sun only once.
- The converse holds: if a planet is subject to an inverse-square centripetal force then the orbit must be an ellipse (or another conic section).
- The proof assumes that a planet is a very small point, but the result holds even for large planets as long as the density of the planet is radially symmetric, that is, for a given distance from the center the density is constant.

### 5.3 Universal gravitation

By Newton's third law, we can equate the force  $F_{S \leftarrow E}$  that the Sun  $S$  exerts on the Earth  $E$  with the force  $F_{E \leftarrow S}$  that the Earth exerts on the Sun. Let  $m$  be the mass of the Earth and  $M$  be the mass of the Sun, then by Equation 5,

$$\begin{aligned} F_{S \leftarrow E} &= \frac{8\kappa_E^2 m}{L_E} \cdot \frac{1}{r^2} = \frac{C_E m}{r^2} \\ F_{E \leftarrow S} &= \frac{8\kappa_S^2 M}{L_S} \cdot \frac{1}{r^2} = \frac{C_S M}{r^2} \\ \frac{C_E}{M} \cdot \frac{1}{r^2} &= \frac{C_S}{m} \cdot \frac{1}{r^2}, \end{aligned}$$

from some constants  $C_E, C_S$ .

Why are the constants different? The Earth and the Sun both rotate around their center of mass called the *barycenter*, which is very close to the center of the Sun since the Sun is so much more massive than the Earth. The ellipse of the Sun's orbit is very small relative to the Earth so  $A$  and  $L$  are smaller, and the Sun's period is large so  $T$  is larger. The different values for  $8\kappa^2/L$  are encapsulated into the constants  $C_E, C_S$ . Let  $G = \frac{C_E}{M} = \frac{C_S}{m}$  so that

$$F_{S \leftarrow E} = F_{E \leftarrow S} = G \frac{mM}{r^2}. \quad (7)$$

This is Newton's law of universal gravitation. It is not specific to planetary orbits but holds between any two bodies with masses  $m, M$ .

## 5.4 Kepler's third law

**Theorem 5.5 (Kepler's third law)** *Let  $P_1, P_2$  be two planets whose elliptical orbits have semi-major axes  $a_1, a_2$  and whose orbital periods around the Sun are  $T_1$  and  $T_2$ . Then*

$$\frac{a_1^3}{T_1^2} = \frac{a_2^3}{T_2^2}.$$

**Proof** By Equations 6 and 7,

$$F = \frac{4\pi^2 a_i^3 m}{T_i^2} \frac{1}{r_i^2} = \frac{GmM}{r_i^2}.$$

After canceling  $m$  and  $r_i$  we get

$$\frac{a_i^3}{T_i^2} = \frac{GM}{4\pi^2}.$$

$GM/4\pi^2$  is a constant that depends only on the mass of the Sun and the gravitational constant, so  $a_i^3/T_i^2$  is constant for all planets rotating around the Sun. ■

## 6 A proof Proposition XI, Problem VI

Theorem 5.4 is Book I, Section III, Proposition XI, Problem VI of the *Principia*.

Study Figure 25:<sup>5</sup>

- Let  $P, Q$  be two points on the ellipse that represent the movement of a body in an elliptical orbit separated by a time interval  $\Delta t$ . Construct lines from  $P$  to the center  $C$  and the foci  $S, H$ .
- Construct the tangent at  $P$  and choose  $R$  on the tangent such that the body would move from  $P$  to  $R$  if it continued for time  $\Delta t$  not subject to any force. Construct the parallelogram  $PRQX$  and extend  $QX$  until it intersects  $PC$  at  $V$ .
- Construct a line parallel to  $RP$  through  $H$  and let  $I$  be its intersection with  $PS$ .
- Construct  $DC$  the conjugate diameter to  $PC$  (Definition 7.9), and let  $E$  be its intersection with  $PS$ .

### 6.1 A formula for $QR$

**Theorem 6.1**  $QR = PV \cdot \frac{CA}{CP}$ .

**Proof** By Theorem 7.8,  $\angle RPX = \angle ZPH = \alpha$  and by alternate interior angles,

$$\angle PHI = \angle ZPH = \alpha = \angle RPX = \angle PIH,$$

so  $\triangle IPH$  (red) is isosceles and  $PI = PH = d$ .

<sup>5</sup>The bottom half of the ellipse is not shown, but we still refer to lines  $DC, PC$  as diameters.




Figure 25: Geometry of an elliptical orbit (1)




Figure 26: Geometry of an elliptical orbit (2)

$SC = CH = c$  are equal because they are the distances of the foci from the center of the ellipse. Let  $SE = e$ . By construction  $EC \parallel IH$  so  $\triangle ESC \sim \triangle ISH$  (blue) and

$$\frac{SC}{SE} = \frac{SH}{SI}$$

$$SI = \frac{SH \cdot SE}{SC} = \frac{2c \cdot e}{c} = 2e.$$

By definition of an ellipse  $SP + PH = SI + IP + PH = 2CA$  so  $2e + d + d = 2CA$  and  $EP = e + d = CA$ .

$QV \parallel EC$  so  $\triangle EPC \sim \triangle XPV$  and

$$\frac{PX}{PV} = \frac{EP}{PC} = \frac{CA}{PC}$$

$$PX = PV \cdot \frac{CA}{PC}.$$

Since  $PRQX$  is a parallelogram  $QR = PX$ ,  $QR = PV \cdot \frac{AC}{PC}$ . ■

## 6.2 A formula for $QT$

Construct a perpendicular from  $P$  to  $DC$  and label its intersection with  $DC$  by  $F$ . Construct a perpendicular from  $Q$  to  $SP$  and label its intersection with  $SP$  by  $T$  (Figure 26).

### Theorem 6.2

$$QT = QX \cdot \frac{FP}{CA}.$$

**Proof** By construction,  $QR \parallel PX$ , so by alternate interior angles  $\angle RQX = \angle QXT = \beta$ . By construction,  $QX \parallel DC$ , so by alternate interior angles  $\angle QXT = \angle PEF = \beta$ . Since  $\triangle PFE$

and  $\triangle QTX$  are right triangles with an equal acute angle  $\beta$ ,  $\triangle PFE \sim \triangle QTX$ . In the proof of Theorem 6.1 we showed that  $EP = CA$  so

$$\begin{aligned}\frac{QT}{QX} &= \frac{FP}{EP} \\ QT &= QX \cdot \frac{FP}{EP} = QX \cdot \frac{FP}{CA}. \blacksquare\end{aligned}$$

### 6.3 A formula for $QR/QT^2$

**Theorem 6.3**

$$\frac{QR}{QT^2} = \frac{CP \cdot CA}{CB^2} \cdot \frac{QV^2}{GV \cdot QX^2}. \quad (8)$$

**Proof** Let us combine the equations in Theorems 6.1 and 6.2 to get  $QR/QT^2$ .

$$\frac{QR}{QT^2} = \frac{PV \cdot \frac{CA}{CP}}{\left(QX \cdot \frac{FP}{CA}\right)^2} = \frac{PV \cdot CA^3}{QX^2 \cdot CP \cdot FP^2}. \quad (9)$$

$DC$  and  $PC$  are conjugate diameters so Theorem 7.11 gives a formula for  $PV$  that we substitute into Equation 10.

$$\frac{QR}{QT^2} = \frac{QV^2 \cdot CP^2}{GV \cdot CD^2} \cdot \frac{CA^3}{QX^2 \cdot CP \cdot FP^2} = \frac{CP \cdot CA^3}{CD^2 \cdot FP^2} \cdot \frac{QV^2}{GV \cdot QX^2}. \quad (10)$$

Next we show that  $CD \cdot FP = CA \cdot CB$ . By Theorem 7.12 the areas of the parallelograms formed by the tangents to conjugate diameters are equal. By symmetry the areas of the four small parallelograms are equal, as are the triangles formed by constructing diagonals.





Figure 27: Parallelograms formed by conjugate diameters



## 7 Ellipses

### 7.1 Fundamental properties

#### Definition 7.1 (Ellipse)

- Let  $S$  and  $H$  be two points in the plane such that  $SH = 2c \geq 0$  and choose  $a$  such that  $2a > 2c$  (Figure 29). An ellipse is the geometric locus of all points  $P$  such that  $SP + PH = 2a$ . If  $c = 0$  the geometric locus is a circle.
- Construct  $AA'$  through  $SH$ , where  $A, B$  are the intersections of the line with the ellipse.  $AA'$  is the major axis of the ellipse. Let  $O$  be the midpoint of  $SH$ .  $AO, OA'$  are the semi-major axes of the ellipse.
- Construct the perpendicular to  $AA'$  at  $O$  and let  $B, B'$  be its intersections with the ellipse.  $BB'$  is the minor axis of the ellipse and  $BO, OB'$  are the semi-minor axes of the ellipse.

#### Theorem 7.2 (Figure 30)

1.  $SB = HB = a$ .
2.  $AO = OA' = a$ .
3.  $BO = OB'$ . (Label  $BO = OB'$  by  $b$ .)

#### Proof

1.  $\triangle SBO \cong \triangle HBO$  by side-angle-side so  $SB = HB$ . Since  $B$  is on the ellipse,  $SB + HB = 2a$  and  $SB = HB = a$  follows.




Figure 29: The definition of an ellipse




Figure 30: The semi-major and semi-minor axes of an ellipse

2. Since  $A$  is on the ellipse,

$$2a = AS + HA = (AO - c) + (AO + c) = 2AO,$$

so  $AO = a$ .  $OA' = a = AO$  follows by symmetry.

3.  $BO = OB'$  follows from  $\triangle SBO \cong \triangle SB'O$ . ■

**Theorem 7.3** A point  $P = (x, y)$  on an ellipse satisfies the equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1. \quad (11)$$

**Proof** Since  $S = (-c, 0)$ ,  $H = (c, 0)$  and  $SP + PH = 2a$ ,

$$PS + PH = \sqrt{(x - (-c))^2 + y^2} + \sqrt{(x - c)^2 + y^2} = 2a.$$

Squaring twice results in

$$(x + c)^2 + y^2 = \left(2a - \sqrt{(x - c)^2 + y^2}\right)^2$$

$$4xc = 4a^2 - 4a\sqrt{(x - c)^2 + y^2}$$

$$a - \frac{c}{a}x = \sqrt{(x - c)^2 + y^2}$$

$$a^2 + \frac{c^2}{a^2}x^2 = x^2 + c^2 + y^2$$

$$\frac{x^2}{a^2} + \frac{y^2}{a^2 - c^2} = \frac{a^2 - c^2}{a^2 - c^2} = 1.$$

By Theorem 7.2 and Pythagoras's theorem,  $b^2 = a^2 - c^2$  so

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1. \quad \blacksquare$$






Figure 31: A circle circumscribing an ellipse

## 7.2 A circle circumscribing an ellipse

Consider a circle of radius  $a$  with the same center as an ellipse (Figure 31). Choose a point  $X$  on the major axis and construct a perpendicular through  $X = (x, 0)$ . Let its intersections with the ellipse and the circle be  $P_e = (x, y_e)$  and  $P_c = (x, y_c)$ , respectively.

**Theorem 7.4** *The perpendicular to the major axis through a point  $P_c = (x, y_c)$  on the circle circumscribing an ellipse intersects the ellipse at  $P_e = (x, y_e) = \left(x, \frac{b}{a}y_c\right)$ .*

**Proof** From Equation 11 and the formula  $x^2 + y^2 = a^2$  for the circle,

$$y_e = \frac{b}{a} \sqrt{(a^2 - x^2)} = \frac{b}{a} y_c. \quad \blacksquare \quad (12)$$

## 7.3 The latus rectum of an ellipse

**Definition 7.5** *Consider a line through a focus of an ellipse that is perpendicular the major axis. Let its intersections with the ellipse be  $L_1, L_2$ . Then  $L = L_1L_2$  is a latus rectum of an ellipse (Figure 32).<sup>6</sup>*

**Theorem 7.6**  *$L$ , the length of the latus rectum of an ellipse, is  $\frac{2b^2}{a}$ .*

**Proof** By Equation 12 and Pythagoras's theorem,

$$L = 2L_1 = 2 \cdot \frac{b}{a} \sqrt{a^2 - c^2} = \frac{2b^2}{a}. \quad \blacksquare$$

<sup>6</sup>Usually, lines are denoted by lower-case letters, but  $L$  for the latus rectum is the standard notation.




Figure 32: The latus rectum of an ellipse

## 7.4 The area of an ellipse

**Theorem 7.7** *The area of an ellipse is  $\pi ab$ .*

**Proof** From Equation 12

$$y_e = \frac{b}{a} \sqrt{a^2 - x^2},$$

so the area of an ellipse is

$$A_e = 2 \int_{-a}^a \frac{b}{a} \sqrt{a^2 - x^2} dx = \frac{b}{a} \cdot 2 \int_{-a}^a \sqrt{a^2 - x^2} dx = \frac{b}{a} A_c.$$

If we can show that the area of a circle is  $\pi a^2$  the theorem follows.

The proof uses polar coordinates, where  $x = a \cos \theta$  and  $y = a \sin \theta$ . First, we derive the formula for the integral of  $\sin^2 \theta$  using the double-angle identity.

$$\sin^2 \theta = \frac{1 - \cos 2\theta}{2}$$

$$\int \sin^2 \theta d\theta = \int \frac{1 - \cos 2\theta}{2} d\theta = \frac{\theta}{2} - \frac{\sin 2\theta}{4} + C.$$

Now we can compute the area of a circle as twice the area of a semicircle by changing from Cartesian to polar coordinates and integrating.

$$\begin{aligned} A_c &= 2 \int_{-a}^a \sqrt{a^2 - x^2} dx = 2 \int_{-\pi}^0 \sqrt{a^2 - (a \cos \theta)^2} d(a \cos \theta) \\ &= 2 \cdot a \cdot a \int_{-\pi}^0 \sin \theta (-\sin \theta) d\theta = -2a^2 \int_{-\pi}^0 \sin^2 \theta d\theta \\ &= -2a^2 \left( \frac{\theta}{2} - \frac{\sin 2\theta}{4} + C \right) \Big|_{-\pi}^0 = \pi a^2. \quad \blacksquare \end{aligned}$$




Figure 33: Angles at the tangent

## 7.5 The angles between a tangent and the lines to the foci

**Theorem 7.8** Let  $P$  be a point on the ellipse whose foci are  $S, H$ . Let  $PU$  be the extension of  $SP$  such that  $SU = AA' = 2a$ . Let  $RQ$  be the bisector of  $\angle HPU$ . Then  $\angle RPS = \angle QPH$  and  $RQ$  is the tangent to the ellipse at  $P$  (Figure 33).

**Proof** We prove that any point  $Q \neq P$  on the bisector is not on the ellipse, so the bisector  $RQ$  has only one point of intersection with the ellipse and it must be the tangent at  $P$ . Since  $RQ$  is the angle bisector of  $\angle HPU$  (the exterior angle of  $\angle SPH$ ),  $\angle QPH = \angle QPU = \alpha$ , and by vertical angles  $\angle QPU = \angle RPS = \alpha$ .

Construct the line  $HU$  to form the triangle  $\triangle HPU$  which intersects  $PQ$  at  $W$ . By construction  $PH = PU$  so  $\triangle HPW \cong \triangle UPW$  by side-angle-side and  $UW = HW$ . But  $HU$  is a straight line, therefore, if  $\angle HWQ = \angle UWQ$ , then they are both right angles and  $\triangle HWQ = \triangle UWQ$  by side-angle-side, so  $UQ = HQ = d$ . Suppose that  $Q$  is on the ellipse so that  $2a = SQ + QH = SQ + QU$ . By the triangle inequality  $2a = SQ + QU > SU = 2a$ , contradicting that  $Q$  is on the ellipse. ■

## 7.6 Conjugate diameters

**Definition 7.9** There are two equivalent definitions of conjugate diameters.

- Let  $P$  be a point on an ellipse,  $PG$  a diameter and let  $t$  be the tangent to the ellipse at  $P$ . Diameter  $DK$  is a conjugate diameter if it is parallel to  $t$  (Figure 34).
- Two diameters  $PG$  and  $DK$  are conjugate diameters if the midpoints of chords  $(D'K', D''K'')$  parallel to one diameter ( $DK$ ) lie on another diameter ( $PG$ ).

**Theorem 7.10** Let  $P = (x, y)$  be a point on an ellipse (not on the major axis  $AA'$ ) and construct a perpendicular  $PV$  from  $P$  to the major axis (Figure 35). Then

$$\frac{A'V \cdot AV}{PV^2} = \frac{a^2}{b^2}.$$




Figure 34: Conjugate diameters

**Proof** By Equation 12,

$$y^2 = b^2 \cdot \left(1 - \frac{x^2}{a^2}\right) = \frac{b^2(a^2 - x^2)}{a^2}$$

$$\frac{A'V \cdot AV}{PV^2} = \frac{(a+x)(a-x)}{y^2} = \frac{a^2(a^2 - x^2)}{b^2(a^2 - x^2)} = \frac{a^2}{b^2}.$$

**Theorem 7.11** *Let  $PG, DK$  be conjugate diameters of an ellipse and let  $Q$  be a point on the ellipse (Figure 36). Construct the perpendicular  $QV$  from  $Q$  to the major axis, then*

$$PV = \frac{QV^2 \cdot CP^2}{GV \cdot CD^2}.$$

**Proof** Figure 36 shows a dashed ellipse which is the original ellipse rotated about the same center  $O$ , so that  $OP$  is the semi-major axis and  $OD$  is the semi-minor axis. By Theorem 7.10,

$$\frac{GV \cdot PV}{OV^2} = \frac{a'^2}{b'^2},$$




Figure 35: Ratios on conjugate diameters




Figure 36: Ratios on conjugate diameters

where  $a', b'$  are the lengths of the semi-major and semi-minor axes of the rotated ellipse. By construction  $a' = OP$  and  $b' = OD$  so

$$\frac{GV \cdot PV}{QV^2} = \frac{CP^2}{OD^2}$$

$$PV = \frac{QV^2 \cdot CP^2}{GV \cdot OD^2} \quad \blacksquare$$

## 7.7 The parametric representation of an ellipse

Figure 37 shows an ellipse and two circles: one whose radius is the length of the semi-major axis (dotted red) and one whose radius is the semi-minor axis (dashed blue). The figure shows the *parametric representation* of a point  $P = (x, y)$  on the ellipse:

$$(x, y) = (a \cos t, b \sin t).$$

The parameter  $t$  is *not* the angle of  $P$  relative to the positive  $x$ -axis. Construct the perpendicular through  $P$  to the minor axis and let  $P_I$  be its intersection with the inner circle so that  $CP_I$  defines an angle  $t$ . Extend  $C_I$  until it intersects the outer circle at  $P_O$ . The parametric representation of  $P$  is computed from the lengths of the axes and trigonometry functions of  $t$ .

## 7.8 Areas of parallelograms

**Theorem 7.12** *The areas of the parallelograms formed by tangents to the intersections of a pair of conjugate diameters with the ellipse are equal (Figure 38).*

**Proof** We show that the area of the parallelogram  $JKLM$  is equal to the area of the parallelogram formed by tangents to the major and minor axes (dashed). By symmetry it suffices to prove that the areas of one of the quadrants of those parallelograms are equal:



triangles  $\triangle D'DC, \triangle P'PC$ . Therefore,

$$\begin{aligned}\triangle PCD &= \frac{y_p + y_d}{2}(x_p - x_d) - \frac{1}{2}x_dy_d - \frac{1}{2}x_py_p = \frac{1}{2}(x_py_d - x_dy_p) \\ &= \frac{1}{2}(a \cos t \cdot b \cos t - (-a) \sin t \cdot b \sin t) = \frac{1}{2}ab = \triangle ABC. \quad \blacksquare\end{aligned}$$

## 8 Ellipses in Euclidean geometry

The proofs of theorems about planetary freely used analytic geometry and trigonometry, but for many years after the invention of analytic geometry, mathematicians continued to limit themselves to Euclidean geometry. In this section, I present proofs in Euclidean geometry of theorems that appeared in Section 7.<sup>7</sup> The proofs are based on the definition of ellipses in terms of the geometric concepts of focus and directrix instead of the familiar analytic definition (Equation 11).

### 8.1 The definition of an ellipse using the focus and the directrix

**Definition 8.1** Let  $d$  be a line (the directrix) and  $S$  be a point (the focus) not on the directrix. Let  $0 < e < 1$  be a number (the eccentricity). An ellipse is the locus of points  $P$  such that the ratio of  $PS$  to the distance of  $P$  to the directrix is  $e$ .

All the conic sections (parabolas, ellipses and hyperbolas) are defined the same way and are distinguished by their eccentricity.

**Definition 8.2** Let  $X$  be the intersection of the perpendicular to the directrix from  $S$ . A on  $SX$  is a vertex of the ellipse if  $SA/AX = e$  (in Figure 39,  $e = 1/2$ ).

Given  $e = SA/AX$  we can locate  $A$  as follows:

$$\begin{aligned} SA + AX &= SX \\ SA &= SX - \frac{SA}{e} = SX \cdot \frac{e}{1+e} \\ AX &= SX \cdot \frac{1}{1+e}. \end{aligned}$$

<sup>7</sup>Theorems 7.8, 7.11 were already proved using Euclidean geometry and Theorem 7.7 requires taking limits.




Figure 39: The elements of the definition of an ellipse






Figure 40: Constructing points on the ellipse

Similarly, there is a second vertex  $A'$  where

$$\begin{aligned} A'X - SA' &= SX \\ SA' &= SX + \frac{SA'}{e} = SX \cdot \frac{e}{1-e} \\ A'X &= SX \cdot \frac{1}{1-e}. \end{aligned}$$

Definition 8.1 is non-constructive. It states that the ellipse is the locus of points satisfying a certain property, but aside from the vertices we have not constructed any such points. Here we show how to construct any of the points on the ellipse.

Select an *arbitrary* point  $E$  on the directrix and construct lines from  $E$  through  $A$  and  $S$ . The line through  $S$  will make an angle  $\alpha$  with  $SX$ . Construct a line from  $S$  at the *same angle*  $\alpha$  from  $ES$  and let its intersection with  $EA$  be  $P$ . Construct the perpendicular from  $P$  to the directrix and let  $K$  be its intersection with the directrix. Let  $L$  be the intersection of  $KP$  with  $ES$  (Figure 40).

**Theorem 8.3** *The point  $P$  is on the ellipse.*

**Proof**  $\angle PLS = \angle LSN = \alpha$  by alternate interior angles, so  $\triangle LPS$  is isosceles and  $PL = SP$ . Since  $PK \parallel SX$ ,  $\triangle XEA \sim \triangle KEP$  and  $\triangle AES \sim \triangle PEL$  are adjacent pairs of similar triangles, so

$$\frac{PS}{PK} = \frac{PL}{PK} = \frac{SA}{AX} = e.$$

Therefore,  $P$  is on the ellipse. ■

By choosing different points  $E$  on the directrix, any point on the ellipse can be constructed.

## 8.2 A right angle at the focus of an ellipse

**Theorem 8.4** *Let  $P, P'$  be points on the ellipse and let  $F$  be the intersection of  $PP'$  with the directrix. Then  $FS$  bisects the exterior angle of  $\angle P'SP$  (Figure 41).*




Figure 41: Bisecting the angle at the focus

**Proof** Since  $P, P'$  are on the ellipse

$$\frac{SP}{PK} = \frac{SP'}{P'K'} = e,$$

and since  $\triangle PFK \sim P'FK'$ ,

$$\frac{SP}{SP'} = \frac{PK}{P'K'} = \frac{PF}{P'F}.$$

By the exterior angle bisector theorem (Theorem A.7),  $FS$  bisects the exterior angle of  $\angle P'SP$ . ■

**Theorem 8.5** Let  $P$  be a point on the ellipse and construct lines  $PA, PA'$ . Label their intersections with the directrix by  $E$  and  $F$ , respectively. Then  $\angle FSE$  is a right angle (Figure 42).

**Proof**  $P, A, A'$  are all points on the ellipse so Theorem 8.4 applies.  $FS$  bisects  $\angle PSX = 2\gamma$  and  $ES$  bisects  $\angle P'SX = 2\delta$ , so  $2\gamma + 2\delta = 180^\circ$  and  $\angle FSE = \gamma + \delta = 90^\circ$ . ■




Figure 42: The right angle at the focus

### 8.3 Ratios of perpendiculars to the axes

This theorem proves Theorem 7.10 using Euclidean geometry.

**Theorem 8.6** *Let  $P$  be a point on an ellipse not on the major axis and construct perpendiculars  $PN, PM$  from  $P$  to the major and minor axes, respectively (Figure 43). Then*

$$\frac{PN^2}{A'N \cdot NA} = \frac{BC^2}{AC^2} = \frac{b^2}{a^2} \quad (13)$$

$$\frac{PM^2}{B'N \cdot NB} = \frac{AC^2}{BC^2} = \frac{a^2}{b^2}. \quad (14)$$

**Proof** (Equation 13)  $\triangle AXE \sim \triangle ANP$  since they are right triangles and the vertical angles at  $A$  are equal (red). Therefore,

$$\frac{PN}{AN} = \frac{EX}{AX}. \quad (15)$$

$\triangle PA'N \sim \triangle FA'X$  (blue) so

$$\frac{PN}{A'N} = \frac{FX}{A'X}. \quad (16)$$

Multiplying Equations 15 and 16 gives

$$\frac{PN^2}{AN \cdot A'N} = \frac{EX \cdot FX}{AX \cdot A'X}.$$

By Theorem 8.5  $\triangle FSE$  is a right triangle so by Theorem A.3,

$$\frac{PN^2}{AN \cdot A'N} = \frac{SX^2}{AX \cdot A'X}.$$

Since  $P$  was arbitrary this holds for any point on the ellipse, in particular, for  $B$  on the minor axis, where  $PN = BC$  and  $AN = AN' = AC$ . Therefore,

$$\begin{aligned} \frac{BC^2}{AC^2} &= \frac{SX^2}{AX \cdot A'X} \\ \frac{PN^2}{AN \cdot A'N} &= \frac{SX^2}{AX \cdot A'X} = \frac{BC^2}{AC^2} = \frac{b^2}{a^2}. \quad \blacksquare \end{aligned}$$

**Proof** (Equation 14) Since  $CM = PN, PM = CN$ , by Theorem A.2, Equation 13 becomes

$$\begin{aligned} \frac{CM^2}{AC^2 - PM^2} &= \frac{BC^2}{AC^2} \\ \frac{AC^2}{AC^2 - PM^2} &= \frac{BC^2}{CM^2}. \end{aligned} \quad (17)$$

It follows that

$$\frac{AC^2}{PM^2} = \frac{BC^2}{BC^2 - CM^2}. \quad (18)$$

By cross-multiplying the Equations 17 and 18. By Theorem A.2, Equation 18 implies

$$\frac{PM^2}{BM \cdot MB'} = \frac{AC^2}{BC^2}. \quad \blacksquare$$






Figure 45: The latus rectum of an ellipse

## 8.5 The latus rectum of an ellipse

This theorem proves Theorem 7.6 in Euclidean geometry.

**Theorem 8.8** *L, the length of the latus rectum of an ellipse (Definition 7.5), is  $\frac{2b^2}{a}$ .*

**Proof** By Theorem 8.6,

$$\frac{SL_1^2}{AS \cdot SA'} = \frac{BC^2}{AC^2}.$$

By Theorem 7.2,  $SB = AC = a$ , so by Pythagoras's theorem,

$$BC^2 = BS^2 - SC^2 = AC^2 - SC^2 = (AC - SC)(AC + SC) = AS \cdot SA'.$$

Therefore, the length of one-half the latus rectum is

$$SL_1^2 = \frac{BC^4}{AC^2}$$

$$SL_1 = \frac{BC^2}{AC} = \frac{b^2}{a}. \quad \blacksquare$$

## 8.6 Areas of parallelograms

**Theorem 8.9** *Let Y be the intersection the perpendicular through the focus S to the tangent TT' at P, and let L be the intersection of S'P and SY (Figure 46). Then Y is on the circumscribing circle and CY  $\parallel$  S'L.*

**Proof**

$\triangle STY \sim \triangle T'TC$  since they are right triangles that share the acute angle  $\angle CTT' = \angle YTS$ , so  $\angle CT'T = \angle YST = \beta$ .






Figure 48: Parallelograms formed by conjugate diameters

- $CY \parallel S'P$  (Theorem 8.9) so  $\angle CYP = \angle S'PT'$  by corresponding angles.
- $\angle S'PT' = \angle SPY$  by Theorem 7.8 since they are the angles to the foci at the tangent.
- $\angle SYP$  and  $\angle SNP$  are right angles and therefore  $SYPN$  is quadrilateral that can be circumscribed by a circle whose diameter is  $PS$ .<sup>8</sup> Therefore,  $\angle SPY = \angle SNY$  since they are subtended by the same chord  $YS$ .

Since  $\angle CYT \sim \angle CNY$  and  $\angle YCT$  is a common angle,  $\triangle CYT \sim \triangle CNY$  and

$$\frac{CN}{CY} = \frac{CY}{CT}$$

$$CN \cdot CT = CY^2 = AC^2, \quad (19)$$

since the perpendicular to the tangent from a focus is on the circumscribing circle (Theorem 8.9). Since  $NT = CT - CN$ , we have  $CN \cdot NT = CN \cdot CT - CN^2$  which equals  $AC^2 - CN^2$  by Equation 19. This in turn equals  $AN \cdot NA'$  by Theorem A.2. ■

Construct the normal to the tangent at  $P$  and let its intersection with the conjugate diameter  $DK$  be  $F$  and its intersection with the major axis be  $G$ . Construct a perpendicular from  $P$  to the major axis and let its intersection be  $N$ . Let the intersection of the tangent with the minor axis be  $T$  and its intersection with the major axis be  $T'$  (Figure 48).

### Theorem 8.11

$$PF \cdot PG = BC^2.$$

<sup>8</sup>It can be proven that a quadrilateral whose opposite angle are supplementary can be circumscribed by a circle. If two opposite angles are right angles that sum to  $180^\circ$ , then the other two angles must also sum to  $180^\circ$ .

**Proof**  $\triangle NPG \sim \triangle FPJ$  (rotate  $\triangle NPG$  to see this) so  $\angle PGN = \angle PJF = \alpha$  and

$$\begin{aligned}\frac{PF}{PN} &= \frac{PJ}{PG} \\ PF \cdot PG &= PJ \cdot PN.\end{aligned}\tag{20}$$

By vertical angles  $\angle PGN = \angle CGF = \alpha$  so  $\triangle NPG \sim \triangle FCG$  and  $\angle NPG = \angle FCG = \beta = 90^\circ - \alpha$ . By adding  $\beta$  to the right angles  $\angle BCN$  and  $\angle TPF$ , we get that  $\angle TCJ = \angle TPJ$  and therefore  $TPJC$  is a parallelogram, so  $CT = PJ$  and  $PF \cdot PG = CT \cdot PN$ . By Equation 20, the theorem will be proven if we can show that  $CT \cdot PN = BC^2$ .

$\triangle TT'C \sim \triangle PT'N$  so

$$\begin{aligned}\frac{CT}{CT'} &= \frac{PN}{NT'} \\ \frac{CT}{PN} &= \frac{CT'}{NT'}.\end{aligned}$$

Multiplying each side by fractions equal to 1 gives

$$\frac{CT \cdot PN}{PN^2} = \frac{CT' \cdot CN}{CN \cdot NT'}.$$

By Equation 19,  $CN \cdot CT' = AC^2$ , and by Theorem 8.10,  $CN \cdot NT' = AN \cdot NA'$ , so

$$\frac{CT \cdot PN}{PN^2} = \frac{AC^2}{AN \cdot NA'}.$$

Multiplying by  $PN^2 / PN^2$  and using Theorem 8.6 gives

$$\begin{aligned}\frac{CT \cdot PN}{PN^2} &= \frac{PN^2}{AN \cdot NA'} \cdot \frac{AC^2}{PN^2} = \frac{BC^2}{AC^2} \cdot \frac{AC^2}{PN^2} \\ CT \cdot PN &= BC^2. \quad \blacksquare\end{aligned}\tag{21}$$

**Theorem 8.12** *In Figure 49,*

$$\begin{aligned}CN^2 &= AM \cdot MA', \quad CM^2 = AN \cdot NA' \\ \frac{DM}{CN} &= \frac{BC}{AC} \\ \frac{CM}{PN} &= \frac{BC}{AC}\end{aligned}$$

**Proof** By Theorem 8.10,

$$\begin{aligned}CN \cdot CT &= AC^2 = CM \cdot CT' \\ \frac{CM}{CN} &= \frac{CT}{CT'}.\end{aligned}$$






Figure 49: Ratios of perpendiculars to the major axis

Since  $DK$  and  $PP'$  are conjugate diameters,  $DT' \parallel PP'$  and  $\triangle T'DC \sim \triangle CPT$ , so

$$\frac{CM}{CN} = \frac{CT}{CT'} = \frac{CN}{MT'}$$

$$CN^2 = CM \cdot MT',$$

Therefore,

$$CN^2 = CM \cdot MT' = AC^2 = AM \cdot MA' \quad (22)$$

by Theorem 8.10. By Theorem 8.6,

$$\frac{DM^2}{AM \cdot MA'} = \frac{BC^2}{AC^2},$$

and by Equation 22,

$$\frac{DM^2}{CN^2} = \frac{BC^2}{AC^2}$$

$$\frac{DM}{CN} = \frac{BC}{AC},$$

A symmetric argument shows that

$$CM^2 = AN \cdot NA'$$

$$\frac{CM}{PN} = \frac{BC}{AC} \quad \blacksquare.$$

This theorem proves Theorem 7.12 in Euclidean geometry.

**Theorem 8.13** *The area of the parallelogram formed by the tangents at the ends of the conjugate diameters  $PP'$ ,  $DK$  is equal to the area of the rectangle enclosing the ellipse at the ends of the axes (Figure 50).*

By vertical angles  $\angle DCM = \angle GCF$  and  $\angle CGF = \angle PGN$ , so  $\triangle DCM \sim \triangle PGN$

and by Theorem 8.12

By Theorem 8.11,

From Equations 23 and 24 gives  $CD \cdot PF = AC \cdot BC$ . ■



## A Theorems of Euclidean Geometry

### A.1 Constructing a circle from three points


**Theorem A.1** *Given three non-collinear points,<sup>9</sup> a circle can be constructed that goes through all three points.*

**Proof** Three non-collinear points  $A, B, C$  define a triangle  $\triangle ABC$  (Figure 51). Construct the perpendicular bisectors of any two of its three sides, say,  $AC$  and  $BC$ . By definition the perpendicular bisector is the geometric locus of points equidistant from the endpoints of the segment. Let  $O$  be the intersection of the two bisectors. Then the  $AO = CO = BO$  is the radius of a circle centered at  $O$  that goes through  $A, B, C$ . ■

### A.2 The product of two subsegments

**Theorem A.2** *Let  $AA'$  be a line segment whose midpoint is  $C$ . Then*

$$AC^2 - CN^2 = AN \cdot NA'.$$



**Proof**  $AN = AC - CN$  and  $NA' = A'C + CN = AC + CN$  since  $C$  is the midpoint of  $AA'$ . The result is obtained by multiplying the two equations. ■

**Theorem A.3** *Let  $Q$  be a point on a circle whose diameter is  $AA'$  and construct a perpendicular  $QN$  to the diameter (Figure 52). Then*

$$QN^2 = AN \cdot NA'.$$

*The equation also holds if it is given that  $\triangle AQA'$  is a right triangle.*

---

<sup>9</sup>Collinear means that the points all on the same line.




Figure 51: A circle through three arbitrary points




Figure 52: Right triangle in a circle at the diameter

**Proof** An angle that subtends a diameter is a right angle. Since the sum of the angles of a triangle is  $180^\circ$ , we can label the angles as shown in the Figure, from which follows that  $\triangle QNA \sim \triangle A'NQ$ . Therefore,

$$\frac{QN}{AN} = \frac{NA'}{QN} . \blacksquare$$

### A.3 Adjacent pairs of similar triangles

I have not encountered the following definition before but it explicitly expresses a relation among similar triangles that would have been obvious to geometers.

**Definition A.4** An adjacent pair of similar triangles is a pair of (a pair of) similar triangles that share sides. In Figure 53,  $\triangle BAC \sim \triangle EAF$  and  $\triangle CAD \sim \triangle FAG$  are an adjacent pair of similar triangles.

**Theorem A.5** For the adjacent pair of similar triangles in Figure 53,

$$\frac{AB}{AE} = \frac{AD}{AG} .$$

**Proof** By similar triangles,

$$\begin{aligned} \frac{AB}{AE} &= \frac{AC}{AF} \\ \frac{AC}{AF} &= \frac{AD}{AG} \\ \frac{AB}{AE} &= \frac{AD}{AG} . \blacksquare \end{aligned}$$

Similar ratios hold between other sides of  $\triangle BAC$  and  $\triangle CAD$  by using an intermediate step with  $AC$ . We will use the term *by an adjacent pair of similar triangles* and leave it to the reader to make the intermediate step.




Figure 53: Adjacent pairs of similar triangles

#### A.4 The angle bisector theorems

**Theorem A.6 (Interior angle bisector theorem)** In  $\triangle ABC$  let  $D$  be a point on  $BC$  (Figure 54). Then  $AD$  bisects  $\angle CAB$  if and only if

$$\frac{BD}{CD} = \frac{AB}{AC}.$$

**Proof** Suppose that  $AD$  bisects  $\angle BAC$ . Construct a line through  $C$  parallel to  $AB$  and let its intersection with  $AD$  be  $E$ . By alternate interior angles,  $\angle BAD = \angle CED$  and by vertical angles  $\angle BDA = \angle CDE$ . Therefore,  $\triangle ABD \sim \triangle EDC$  so

$$\frac{BD}{CD} = \frac{AB}{CE}.$$

$\triangle ECA$  is isosceles so  $CE = AC$  and

$$\frac{BD}{CD} = \frac{AB}{AC}.$$

To prove the converse just “run” the proof backwards. ■




Figure 54: The interior angle bisector theorem




Figure 55: The exterior angle bisector theorem

**Theorem A.7 (Exterior angle bisector theorem)** In  $\triangle ABC$  let  $D$  be a point on the extension of  $CB$  outside the triangle (Figure 55). Then  $AD$  bisects the exterior angle of  $\angle BAC$  if and only if

$$\frac{BD}{CD} = \frac{AB}{AC}.$$

**Proof** Suppose that  $AD$  bisects  $\angle BAF$ . Construct a line through  $B$  parallel to  $AD$  and let its intersection with  $AC$  be  $E$ . By alternate interior angles  $\angle BAD = \angle ABE$  and by corresponding angles  $\angle FAD = \angle AEB$ . Therefore,  $\triangle BCE \sim \triangle DCA$  so

$$\frac{BD}{CD} = \frac{AE}{AC}.$$

But  $\triangle BAE$  is isosceles so  $AE = AB$  and

$$\frac{BD}{CD} = \frac{AB}{AC}.$$

To prove the converse just “run” the proof backwards. ■

The exterior angle bisector theorem confusing to understand in a proof, because it can be hard to identify the components of a diagram. In the text the following color-coding is used: the triangle is red, the extension of one side is dashed red and the bisector is blue.

## Sources and further reading

Sections 2–5 are primarily on Hahn’s book [5]. He has written a more advanced book on the orbits of planets and spacecraft [6]. The proof in Section 6 is from [4]. Two additional articles present this aspect of Newton’s work [7, 9]. Should you wish to read Newton’s work, [2] is an up-to-date translation into English, and Cohen’s lengthy *Guide* will facilitate understanding Newton’s often terse presentation. The relevant sections of the *Guide* are 10.8–10.10. The Wikipedia entry on ellipses is very comprehensive [10].

Section 8 is based on Besant’s textbook that uses Euclidean geometry exclusively to prove theorems on conic sections [1]. Drew wrote a similar textbook, though shorter and more elementary [3]. Compare these textbooks one by one with Smith’s that uses analytic geometry [8].<sup>10</sup>

## References

- [1] W. H. Besant. *Conic Sections, Treated Geometrically (Ninth Edition Revised and Enlarged)*. George Bell and Sons, London, 1895. <https://www.gutenberg.org/ebooks/29913> and <https://archive.org/details/cu31924059322481> (Accessed 6 September 2023).
- [2] I. Bernard Cohen, Anne Whitman, and Julia Budenz. *The Principia: Mathematical Principles of Natural Philosophy*. University of California Press, 1999. Preceded by *A Guide to Newton’s Principia* by I. Bernard Cohen.
- [3] W. H. Drew. *A Geometrical Treatise on Conic Sections (Second Edition)*. Macmillan, Cambridge, 1862. <https://archive.org/details/in.ernet.dli.2015.501433> (Accessed 6 September 2023).
- [4] Graham Griffiths. The inverse square law of gravitation: An alternative to Newton’s derivation. [https://www.researchgate.net/publication/264978661\\_The\\_Inverse\\_Square\\_Law\\_of\\_Gravitation\\_An\\_Alternative\\_to\\_Newton's\\_Derivation](https://www.researchgate.net/publication/264978661_The_Inverse_Square_Law_of_Gravitation_An_Alternative_to_Newton's_Derivation), 2009. (Accessed 6 September 2023).
- [5] Alexander J. Hahn. *Calculus in Context: Background, Basics, and Applications*. Johns Hopkins University Press, 2017.
- [6] Alexander J. Hahn. *Basic Calculus of Planetary Orbits and Interplanetary Flight: The Missions of the Voyagers, Cassini, and Juno*. Springer, 2020.
- [7] Kai Hauser and Reinhard Lang. On the geometrical and physical meaning of Newton’s solution to Kepler’s problem. *The Mathematical Intelligencer*, 25(4):35–44, 2003.
- [8] Charles Smith. *An Elementary Treatise on Conic Sections*. Macmillan, New York, 1904. <https://archive.org/details/elementarytreati1905smit> (Accessed 6 September 2023).

---

<sup>10</sup>The dates given for all three books are for the editions that I used.

- [9] S. K. Stein. Exactly how did Newton deal with his planets? *The Mathematical Intelligencer*, 18(2):6–11, 1996.
- [10] Wikipedia contributors. Ellipse — Wikipedia, the free encyclopedia. <https://en.wikipedia.org/w/index.php?title=Ellipse&oldid=1168583658>, 2023. [Online; accessed 6-September-2023].