

Analysis

Gaotang Li

August 17, 2022

Contents

I	Foundations	3
1	Groups and Homomorphism	4
1.1	Basics	4
1.2	Subgroup	5
1.3	Cosets	5
1.4	Homomorphisms	6
2	Rings, Fields and Polynomials	8
2.1	Rings	8
2.2	Consequence of Ring Definitions	9
2.3	Field	9
2.4	Ordered Field	10
2.5	Formal Power Series	10
2.6	Polynomials	11
3	The Real Numbers	12
3.1	Order Completeness	12
3.2	The Consequence of Order Completeness	13
4	Vector Spaces, Affine Spaces and Algebras	14
4.1	Vector Spaces	14
II	Convergence	15
5	Convergence of Sequences	16
5.1	Sequences	16
5.2	Metric Space	16
5.3	Convergence	17
6	Normed Vector Space	19
6.1	Norms	19
6.2	The Space of Bounded Functions	20
6.3	Inner Product Spaces	21
6.4	Euclidean Spaces	22
7	Infinite Limits	24
7.1	Convergence to $\pm\infty$	24
7.2	The Limit Superior and Limit Inferior	24
8	Completeness	26
8.1	Cauchy Sequences	26
8.2	Banach Spaces	26

III	Continuous Functions	27
9	Continuity	28
9.1	Elementary Properties and Examples	28
9.2	Sequential Continuity	29
9.3	Addition and Multiplication of Continuous Functions	29
9.4	One-Sided Continuity	30
10	The Fundamentals of Topology	31
10.1	Open Sets	31
10.2	Closed Sets	31
10.3	The Closure and Interior of a Set	32
10.4	The Hausdorff Condition	33
10.5	A Characterization of Continuous Functions	33
10.6	Continuous Extensions	34
10.7	Relative Topology	34

Part I

Foundations

Chapter 1

Groups and Homomorphism

1.1 Basics

Definition 1.1.1 (Group). A pair (G, \odot) consisting of a nonempty set G and an operation \odot is called a **group** if the following holds:

- G is closed under the operation \odot
- \odot is associative
- \odot has an identity element e
- Each $g \in G$ has an **inverse** $h \in G$ such that $g \odot h = h \odot g = e$

Definition 1.1.2 (Abelian group). A group G, \odot is called **commutative** or **Abelian** if \odot is a commutative operation on G .

Remark. Let $G = (G, \odot)$

- (a) the identity element e is unique
- (b) Each $g \in G$ has a unique inverse which we denote by g^b . In particular $e^b = e$.
- (c) For each $g \in G$, we have $(g^b)^b = g$.
- (d) For arbitrary group elements g and h , $(g \odot h)^b = h^b \odot g^b$

Example. (a) Let $G := \{e\}$ be a one element set. Then $\{G, \odot\}$ is an Abelian group, the **trivial group**, with the (only possible) operation $e \odot e = e$.

(b) Let X be a nonempty set, and S_X be the set of all bijections from X to itself. Then $S_X := (S_X, \circ)$ is a group with identity element id_X when \circ denotes the composition of functions. Further, the inverse function f^{-1} is the inverse of $f \in S_X$ in the group. When X is finite, the element of S_X are called permutations and S_X is called the **permutation group** of X .

(c) Let X be a nonempty set and G, \odot a group. With the induced operation \odot , (G^X, \odot) is a group. The inverse of $f \in G^X$ is the function

$$f^b: X \rightarrow G, \quad x \mapsto (f(x))^b$$

(d) Let G_1, \dots, G_m be groups. Then $G_1 \times \dots \times G_m$ with the operation defined analogously to (d) is a group called the **direct product** of G_1, \dots, G_m .

1.2 Subgroup

Definition 1.2.1 (Subgroup). Let $G = (G, \odot)$ be a group and H a nonempty subset of G , if

- $H \odot H \subseteq H$
- $h^b \in H$ for all $h \in H$

then $H := (H, \odot)$ is itself a group and is called a **subgroup** of G .

Remark. Here we use the same symbol \odot for the restriction of the operation to H . Since H is nonempty, there is some $h \in H$ and so, from the two axioms above, $e = h^b \odot h$ is also in H .

Example. Let $G = (G, \odot)$ be a group.

- The trivial subgroup $\{e\}$ and G itself are subgroups of G , the smallest and largest subgroups with respect to inclusion
- If H_α , $\alpha \in A$ are subgroups of G , then $\bigcap_\alpha H_\alpha$ is also a subgroup of G .

1.3 Cosets

Definition 1.3.1 (Coset). Let N be a subgroup of G and $g \in G$. Then $g \odot N$ is the **left coset** and $N \odot g$ is the **right coset** of $g \in G$ with respect to N .

Remark. The definition of coset is related to the particular element.

Note. If we define

$$g \sim h \Leftrightarrow g \in h \odot N \quad (1.1)$$

Then \sim is an equivalence on G .

Proof. \sim is reflexive because $e \in N$

Let $g \in h \odot N$ and $h \in k \odot N$, then

$$g \in (k \odot N) \odot N = k \odot (N \odot N) = k \odot N$$

Let $g \in h \odot N$, then there is some $n \in N$ with $g = h \odot n$. Then it follows that $h = g \odot n^b \in N$. ■

Here 1.1 defines an equivalence relation on G . For the equivalence classes $[\cdot]$ with respect to \sim , we have

$$[g] = g \odot N, \quad g \in G. \quad (1.2)$$

For this reason, we denote G/\sim by G/N , and call G/N the **set of left cosets** of G **modulo** N . Particularly, we have subgroups N such that

$$g \odot N = N \odot g, \quad g \in G. \quad (1.3)$$

Such a subgroup 1.3 is called a **normal subgroup** of G . We call $g \odot N$ the **coset of g modulo N** since each left coset is a right coset and vice versa. We have a well-defined operation on G/N where N is the normal subgroup of G , induced from \odot , such that

$$(G/N) \times (G/N) \rightarrow G/N, \quad (g \odot N, h \odot N) \mapsto (g \odot h) \odot N \quad (1.4)$$

Proposition 1.3.1. Let G be a group and N a normal subgroup of G . Then G/N with the induced

operation is a group, the **quotient group of G modulo N** .

Proof. It is easy to check that the operation is associative. Since $(e \odot N) \odot (g \odot N) = (e \odot g) \odot N = g \odot N$, the identity element of G/N is $N = e \odot N$. Also

$$(g^b \odot N) \odot (g \odot N) = (g^b \odot g) \odot N = N$$

■

Remark. (a) In notion of 1.1, $[e] = N$ is the identity element of G/N and $[g]^b = [g^b]$ is the inverse of $[g] \in G/N$. We also have $[g] \odot h = [g \odot h]$, $g, h \in G$.

(b) Any subgroup N of an Abelian group G is normal and so G/N is a group. Meanwhile, G/N is Abelian.

1.4 Homomorphisms

Definition 1.4.1 (Homomorphism). Let $G = (G, \odot)$ and $G' = (G', \otimes)$ be groups... A function $\varphi: G \rightarrow G'$ is called a **(group) homomorphism** if

$$\varphi(g \odot h) = \varphi(g) \otimes \varphi(h), \quad g, h \in G$$

Definition 1.4.2 (Endomorphism). A homomorphism from G to itself

Remark. (a) Let e and e' be the identity elements of G and G' respectively, and let $\varphi: G \rightarrow G'$ be a homomorphism. Then

$$\varphi(e) = e' \quad \text{and} \quad (\varphi(g))^b = \varphi(g^b), \quad g \in G$$

Proof. $e' \otimes \varphi(e) = \varphi(e) = \varphi(e \odot e) = \varphi(e) \otimes \varphi(e)$
 $e' = \varphi(e) = \varphi(g^b \odot g) = \varphi(g^b) \otimes \varphi(g)$

■

(b) Let $\varphi: G \rightarrow G'$ be a homomorphism. The **kernel** of φ , $\ker(\varphi)$, defined by

$$\ker(\varphi) := \varphi^{-1}(e') = \{g \in G; \varphi(g) = e'\}$$

is a normal subgroup of G .

Proof. First, try to prove $\ker(\varphi)$ is a subgroup of G . For all $g, h \in G$,

- $\varphi(g \odot h) = \varphi(g) \otimes \varphi(h) = e' \otimes e' = e'$
- $\varphi(g^b) = (\varphi(g))^b = (e')^b = e'$

Second, try to prove it is a normal subgroup. Let $h \in g \odot \ker(\varphi)$. Then we there is some $n \in G$ such that $\varphi(n) = e'$ and $h = g \odot n$. For $m := g \odot n \odot g^b$, we have

$$\varphi(m) = \varphi(g) \otimes \varphi(n) \otimes \varphi(g^b) = \varphi(g) \otimes \varphi(g^b) = e'$$

and hence $m \in \ker(\varphi)$. Since $m \odot g = g \odot m = h$, this implies that $h \in \ker(\varphi) \odot g$. So $\ker(\varphi) \odot g \subseteq g \odot \ker(\varphi)$. Similarly one can show $g \odot \ker(\varphi) \subseteq \ker(\varphi) \odot g$. ■

(c) Let $\varphi: G \rightarrow G'$ be a homomorphism and $N := \ker(\varphi)$. Then

$$g \odot N = \varphi^{-1}(\varphi(g)), \quad g \in G,$$

and so

$$g \sim h \Leftrightarrow \varphi(g) = \varphi(h), \quad g, h \in G,$$

where \sim denotes the equivalence relation 1.1.

- (d) A homomorphism is injective if and only if its kernel is trivial, that is, $\ker(\varphi) = \{e\}$
- (e) The image $\text{im}(\varphi)$ of a homomorphism $\varphi: G \rightarrow G'$ is a subgroup of G' .

Example. (a) The constant function $G \rightarrow G', g \mapsto e'$ is a homomorphism, the **trivial** homomorphism.

(b) The identity function $\text{id}_G: G \rightarrow G$ is an endomorphism.

(c) Compositions of homomorphisms (endomorphisms) are homomorphisms (endomorphisms).

(d) If $\varphi: G \rightarrow G'$ is a bijective homomorphism, then so is $\varphi^{-1}: G' \rightarrow G$.

Definition 1.4.3 (Isomorphism). A homomorphism $\varphi: G \rightarrow G'$ is called a **(group) isomorphism** from G to G' if φ is bijective.

In this circumstance, we say that the groups G and G' are **isomorphic** and write $G \cong G'$.

Definition 1.4.4 (Automorphism). An isomorphism from G to itself.

Chapter 2

Rings, Fields and Polynomials

2.1 Rings

Definition 2.1.1 (Ring). A triple $(R, +, \cdot)$ consisting of a nonempty set R and operations, **addition** $+$ and **multiplication** \cdot , is called a **ring** if

- $(R, +)$ is an Abelian group
- Multiplication is associative
- The **distributive law** holds:

$$(a + b) \cdot c = a \cdot c + b \cdot c, \quad c \cdot (a + b) = c \cdot a + c \cdot b, \quad a, b, c \in R$$

Note. A ring is called **commutative** if multiplication is commutative.

If there is an identity element with respect to multiplication, then it is written as 1_R or simply 1 , and is called the **unity** (or **multiplicative identity**) of R , and we say $(R, +, \cdot)$ is a **ring with unity**.

When the addition and multiplication operations are clear from context, we write simply R instead of $(R, +, \cdot)$.

Example. (a) The **trivial ring** has exactly one element 0 and is itself denoted by 0 . A ring with more than one element is **nontrivial**. If R is a ring with unity, then it follows from $1_R \cdot a = a$ for each $a \in R$, that R is trivial if and only if $1_R = 0_R$.

(b) Suppose R is a ring and S is a nonempty subset of R that satisfies the following:

- S is a subgroup of $(R, +)$.
- $S \cdot S \subseteq S$

Then S itself is a ring, a **subring** of R , and R is called an **overring** of S . If R is commutative then so is S , but the converse is not true in general.

(c) Intersections of subrings are subrings.

Definition 2.1.2 (Ring Homomorphism). Let R and R' be rings. A **(ring) homomorphism** is a function $\varphi: R \rightarrow R'$ which is compatible with the ring operations, that is,

$$\varphi(a + b) = \varphi(a) + \varphi(b), \quad \varphi(ab) = \varphi(a)\varphi(b), \quad a, b \in R \quad (2.1)$$

Note. If, in addition, φ is bijective, then φ is called a **(ring) isomorphism** and R and R' are **isomorphic**.

A homomorphism φ from R to itself is a **(ring) endomorphism**. If φ is an isomorphism, then it is a **(ring) automorphism**.

Example. (a) A ring homomorphism $\varphi: R \rightarrow R'$ is, in particular, a group homomorphism from $(R, +)$ to $(R', +)$. The **kernel**, $\ker(\varphi)$, of φ is defined to be the kernel of this group homomorphism, that is,

$$\ker(\varphi) = \{a \in R; \varphi(a) = 0\} = \varphi^{-1}(0)$$

(b) The **zero function** $R \rightarrow R'$, $a \mapsto 0_{R'}$ is a homomorphism with $\ker(\varphi) = R$.

(c) Let R and R' be rings with unity and $\varphi: R \rightarrow R'$ a homomorphism. As (b) shows, it does not necessarily follow that $\varphi(1_R) = 1_{R'}$. This can be seen as a consequence of the fact that, with respect to multiplication, a ring is not a group.

2.2 Consequence of Ring Definitions

Definition 2.2.1 (The Binomial Theorem). Let a and b be two commuting elements ($ab = ba$) of a ring R with unity. Then, for all $n \in \mathbb{N}$,

$$(a + b)^n = \sum_{k=0}^n \binom{n}{k} a^k b^{n-k}$$

Lemma 2.2.1. For $m \in \mathbb{N}$ with $m \geq 2$, an element $\alpha = (\alpha_1 \dots \alpha_m) \in \mathbb{N}^m$ is called a **multi-index**. The **length** $|\alpha|$ of a multi-index $\alpha \in \mathbb{N}^m$ is defined by

$$|\alpha| := \sum_{j=1}^m \alpha_j$$

Set also

$$\alpha! := \prod_{j=1}^m (\alpha_j)!,$$

and define the **natural (partial) order** on \mathbb{N}^m by

$$\alpha \leq \beta \Leftrightarrow (\alpha_j \leq \beta_j, \ 1 \leq j \leq m).$$

for $a = (a_1, \dots, a_m) \in R^m$ and $\alpha = (\alpha_1, \dots, \alpha_m) \in \mathbb{N}^m$

Definition 2.2.2 (The Multinomial Theorem). Let R be a commutative ring with unity. Then for all $m \geq 2$,

$$\left(\sum_{j=1}^m a_j\right)^k = \sum_{|\alpha|=k} \frac{k!}{\alpha!} a^\alpha, \quad a = (a_1, \dots, a_m) \in R^m, \ k \in \mathbb{N}.$$

2.3 Field

Definition 2.3.1 (Field). K is a **field** when the following are satisfied:

- K is a commutative ring with unity.
- $0 \neq 1$
- $K^\times := K \setminus \{0\}$ is an Abelian group with respect to multiplication.

Note. The Abelian group $K^\times := (K^\times, \cdot)$ is called the **multiplicative group** of K .

Remark. Let K be a field.

- (a) For all $a \in K^\times$, $(a^{-1})^{-1} = a$
- (b) A field has no zero divisors
- (c) Let $a \in K^\times$ and $a \in K^\times$ and $b \in K$. Then there is an unique $x \in K$ with $ax = b$, namely the **quotient** $\frac{b}{a} := b/a := ba^{-1}$
- (d) Let K' be a field and $\varphi: K \rightarrow K'$ a homomorphism with $\varphi \neq 0$. Then

$$\varphi(1_K) = 1_{K'} \quad \text{and} \quad \varphi(a^{-1}) = \varphi(a)^{-1}, \quad a \in K^\times$$

2.4 Ordered Field

Definition 2.4.1 (Ordered Ring). A ring R with an ordered \leq is called an **ordered ring** if the following holds:

- (R, \leq) is totally ordered.
- $x < y \Rightarrow x + z < y + z, z \in R$
- $x, y > 0 \Rightarrow xy > 0$

Note. This leads to a series of basic arithmetic rules.
We may define absolute value function from $K \mapsto K$.

Proposition 2.4.1. Let K be an ordered field and $x, y, a, \epsilon \in K$ with $\epsilon > 0$.

- (i) $x = |x|\text{sign}(x)$, $|x| = x\text{sign}(x)$
- (ii) $|x| = |-x|, x \leq |x|$
- (iii) $|xy| = |x||y|$
- (iv) $|x| \geq 0$ and $(|x| = 0 \Leftrightarrow x = 0)$
- (v) $|x - a| < \epsilon \Leftrightarrow a - \epsilon < x < a + \epsilon$
- (vi) $|x + y| \leq |x| + |y|$ (**triangular inequality**)

Corollary 2.4.1 (reversed triangular inequality). In any ordered field K we have

$$|x - y| \geq ||x| - |y||, \quad x, y \in K.$$

2.5 Formal Power Series

Definition 2.5.1 (formal power series). Let R be a nontrivial ring with unity. On the set $R^\mathbb{N} = \text{Func}(\mathbb{N}, R)$ define addition by

$$(p + q)_n := p_n + q_n, \quad n \in \mathbb{N},$$

and multiplication by **convolution**,

$$(pq)_n := (p \cdot q)_n := \sum_{j=0}^n p_j q_{n-j} = p_0 q_n + p_1 q_{n-1} + \cdots + p_n q_0$$

for $n \in \mathbb{N}$. Here p_n denotes the value of $p \in R^{\mathbb{N}}$ at $n \in \mathbb{N}$ and is called the n^{th} **coefficient** of p . In this situation an element $p \in R^{\mathbb{N}}$ is called a **formal power series over R** , and we set $R[X] := (R^{\mathbb{N}}, +, \cdot)$

Proposition 2.5.1. $R[X]$ is a ring with unity, the **formal power series ring over R** . If R is commutative, then so is $R[X]$

2.6 Polynomials

Definition 2.6.1 (Polynomial). A **polynomial over R** is a formal power series $p \in R[X]$ such that $\{n; p_n \neq 0\}$ is finite, in other words, $p_n = 0$ "almost everywhere".

Chapter 3

The Real Numbers

Starting words: we seek an ordered **extension field** of \mathbb{Q} in which the equation $x^2 = a$ is solvable for each $a > 0$.

3.1 Order Completeness

We say a totally ordered set X is **order complete** (or X satisfies the **completeness axiom**) if every nonempty subset of X which is bounded above has a supremum.

Proposition 3.1.1. Let X be a totally ordered set. Then the followings are equivalent:

- (i) X is order complete.
- (ii) Every nonempty subset of X which is bounded below has an infimum.
- (iii) For all nonempty subsets A, B of X such that $a \leq b$ for all $(a, b) \in A \times B$, there is some $c \in X$ such that $a \leq c \leq b$ for all $(a, b) \in A \times B$ (**Dedekind cut property**)

Note. A relation \leq on X is a **partial order** on X if it is reflexive, transitive and **anti-symmetric**, that is,

$$(x \leq y)(y \leq x) \Rightarrow x = y$$

If \leq is a partial order on X , then the pair (X, \leq) is called a **partially ordered set**. If, in addition,

$$\forall x, y \in X: (x \leq y) \vee (y \leq x)$$

then \leq is called a **total order** on X and (X, \leq) is a **totally ordered set**.

Corollary 3.1.1. A totally ordered set is order complete if and only if every nonempty bounded subset has a supremum and an infimum.

Theorem 3.1.1 (Dedekind's Construction of the Real Numbers). There is, up to isomorphism, a unique order complete extension field \mathbb{R} of \mathbb{Q} . This extension is called **the field of real numbers**.

Proposition 3.1.2 (A Characterization of Supremum and Infimum). Followed from natural order defined by \mathbb{R} .

- (i) If $A \subseteq \mathbb{R}$ and $x \in \mathbb{R}$, then
 - (a) $x < \sup(A) \Leftrightarrow \exists a \in A$ such that $x < a$.
 - (b) $x < \inf(A) \Leftrightarrow \exists a \in A$ such that $x > a$.

(ii) Every subset A of \mathbb{R} has a supremum and an infimum in \mathbb{R}

3.2 The Consequence of Order Completeness

The Archimedean Property

Proposition 3.2.1 (Archimedes). \mathbb{N} is not bounded above in \mathbb{R} , that is, for each $x \in \mathbb{R}$ there is some $n \in \mathbb{N}$ such that $n > x$.

Corollary 3.2.1. Equivalent statements as the above proposition

- (a) Let $a \in \mathbb{R}$. If $0 \leq a \leq 1/n$ for all $n \in \mathbb{N}^\times$.
- (b) For each $a \in \mathbb{R}$ with $a > 0$ there is some $n \in \mathbb{N}^\times$ such that $1/n < a$.

The Density of the Rational/Irrational Numbers in \mathbb{R}

Proposition 3.2.2. For all $a, b \in \mathbb{R}$ such that $a < b$, there is some $r \in \mathbb{Q}$ such that $a < r < b$.

Proposition 3.2.3 (n^{th} Roots). For all $a \in \mathbb{R}^+$ and $n \in \mathbb{N}^\times$, there is a unique $x \in \mathbb{R}^+$ such that $x^n = a$

Proposition 3.2.4. For all $a, b \in \mathbb{R}$ such that $a < b$, there is some $\xi \in \mathbb{R} \setminus \mathbb{Q}$ such that $a < \xi < b$.

Intervals

An **interval** is a subset J of \mathbb{R} such that

$$(x, y \in J, x < y) \Rightarrow (z \in J \text{ for } x < z < y)$$

If J is a nonempty interval, then $\inf(J) \in \bar{\mathbb{R}}$ is the **left endpoint** and $\sup(J) \in \bar{\mathbb{R}}$ is the **right endpoint** of J . J is **closed on the left** if $a := \inf(J)$ is in J , and otherwise it is **open on the left**. The same applies to the other side.

An interval is **perfect** if it contains at least two points. It is **bounded** if both endpoints are in \mathbb{R} and is **unbounded** otherwise. If J is a bounded interval, then the nonnegative number $|J| := \sup(J) - \inf(J)$ is called the **length** of J .

Chapter 4

Vector Spaces, Affine Spaces and Algebras

4.1 Vector Spaces

Definition 4.1.1 (Vector Space). A **vector space over the field K** (or simply, a **K -vector space**) is a triple $(V, +, \cdot)$ consisting of a nonempty set V , an 'inner' operation $+$ on V called **addition**, and an 'outer' operation

$$K \times V \rightarrow V, \quad (\lambda, v) \mapsto \lambda \cdot v,$$

called **scalar multiplication** which satisfy the following axioms:

- $(V, +)$ is an Abelian group
- The distributive law holds:

$$\lambda \cdot (v + w) = \lambda \cdot v + \lambda \cdot w, (\lambda + \mu) \cdot v = \lambda \cdot v + \mu \cdot v, \quad \lambda, \mu \in K, \quad v, w \in V$$

- $\lambda \cdot (\mu v) = (\lambda\mu) \cdot v, 1 \cdot v = v$ for $\lambda, \mu \in K, v \in V$

A vector space is called **real** if $K = \mathbb{R}$ and **complex** if $K = \mathbb{C}$.

Note (Linear Functions). Let V and W be vector spaces over K . Then a function $T: V \mapsto W$ is **(K -)linear** if

$$T(\lambda v + \mu w) = \lambda T(v) + \mu T(w), \quad \lambda, \mu \in K, \quad v, w \in V$$

In this regard, a linear function is simply a function which is compatible with the vector space operations, in other words, it is a **(vector space) homomorphism**. The set of all linear functions from V to W is denoted by $\text{Hom}(V, W)$ or $\text{Hom}_K(V, W)$, and $\text{End}(V) := \text{Hom}(V, V)$ is the set of all (vector space) **endomorphisms**. A bijective homomorphism $T \in \text{Hom}(V, W)$ is a (vector space) **isomorphism**.

Remark. (a) A vector space homomorphism $T: V \mapsto W$ is, in particular, a group homomorphism $T: (V, +) \mapsto (W, +)$.

(b)

Part II

Convergence

Chapter 5

Convergence of Sequences

5.1 Sequences

Definition 5.1.1 (Sequence). Let X be a set. A **sequence** (in X) is simply a function from \mathbb{N} to X . If $\varphi: \mathbb{N} \mapsto X$ is a sequence, we write also

$$(x_n), (x_n)_{n \in \mathbb{N}} \text{ or } (x_0, x_1, x_2, \dots)$$

for φ , where $x_n := \varphi(n)$ is the n^{th} term of the sequence $\varphi = (x_0, x_1, x_2, \dots)$.

Remark. (a) A sequence (x_n) is different from its image $\{x_n; n \in \mathbb{N}\}$.

(b) Let (x_n) be a sequence in X and E a property. Then we say E holds for **almost all** terms of (x_n) if there is some $m \in \mathbb{N}$ such that $E(x_n)$ is true for all $n \geq m$, that is, if E holds for all but finitely many of the x_n . If there is a subset $N \subseteq \mathbb{N}$ with $\text{Num}(N) = \infty$ and $E(x_n)$ is true for each $n \in N$ then E is true for **infinitely many** terms.

(c) For $m \in \mathbb{N}^\times$, a function $\Phi: m + \mathbb{N} \mapsto X$ is also called a sequence in X .

5.2 Metric Space

Definition 5.2.1 (Metric Space). Let X be a set. A function $d: X \times X \mapsto \mathbb{R}^+$ is called a **metric** on X if the following hold:

- $d(x, y) = 0 \leftrightarrow x = y$.
- $d(x, y) = d(y, x)$, $x, y \in X$ (symmetry).
- $d(x, y) \leq d(x, z) + d(y, z)$, $x, y, z \in X$ (triangle inequality).

Note. If d is a metric on X , then (X, d) is called a **metric space**. We call $d(x, y)$ the **distance** between the **points** x and y in the metric space X .

In the metric space (X, d) , for $a \in X$ and $r > 0$, the set

$$\mathbb{B}(a, r) := \mathbb{B}_X(a, r) := \{x \in X; d(a, x) < r\}$$

is called the **open ball** with center at a and radius r , while

$$\bar{\mathbb{B}}(a, r) := \bar{\mathbb{B}}_X(a, r) := \{x \in X; d(a, x) \leq r\}$$

is called the **closed ball** with center at a and radius r .

Example. (a) \mathbb{K} is a metric space with the **natural metric**

$$\mathbb{K} \times \mathbb{K} \mapsto \mathbb{R}^+, \quad (x, y) \mapsto |x - y|$$

- (b) Let (X, d) be a metric space and Y a nonempty subset of X . Then the restriction of d to $Y \times Y$, $d_Y := d|_{Y \times Y}$, is a metric on Y , the **induced metric**, and (Y, d_Y) is a metric space, a **metric subspace** of X .
- (c) Let X be a nonempty set. Then the function $d(x, y) := 1$ for $x \neq y$ and $d(x, x) := 0$ is a metric, called the **discrete metric** on X .
- (d) Let (X_j, d_j) , $1 \leq j \leq m$, be metric spaces and $X := X_1 \times \cdots \times X_m$. Then the function

$$d(x, y) := \max_{1 \leq j \leq m} d_j(x_j, y_j)$$

for $x := (x_1, \dots, x_m) \in X$ and $y := (y_1, \dots, y_m) \in X$ is a metric on X called the **product metric**. The metric space $X := (X, d)$ is called the **product of the metric spaces** (X_j, d_j)

Proposition 5.2.1. Let (X, d) be a metric space. Then for all $x, y, z \in X$ we have

$$d(x, y) \geq |d(x, z) - d(z, y)|$$

Note. A subset U of a metric space X is called a **neighborhood** of $a \in X$ if there is some $r > 0$ such that $\mathbb{B}(a, r) \subseteq U$. The **set of all neighborhoods of the point** a is denoted by $\mu(a)$, that is,

$$\mu(a) := \mu_X(a) := \{U \subseteq X; U \text{ is a neighborhood of } a\} \subseteq P(X)$$

Cluster Point

Definition 5.2.2 (Cluster Point). We call $a \in X$ a **cluster point** of (x_n) if every neighborhood of a contains infinitely many terms of the sequence.

Proposition 5.2.2. The following are equivalent:

- (i) a is a cluster point of (x_n) .
- (ii) For each $U \in \mu(a)$ and $m \in \mathbb{N}$, there is some $n \geq m$ such that $x_n \in U$.
- (iii) For each $\epsilon > 0$ and $m \in \mathbb{N}$, there is some $n \geq m$ such that $x_n \in \mathbb{B}(a, \epsilon)$

5.3 Convergence

Definition 5.3.1 (Convergence). A sequence (x_n) **converges** (or is **convergent**) with **limit** a if each neighborhood of a contains almost all terms of the sequence. In this case we write

$$\lim_{n \rightarrow \infty} x_n = a \text{ or } x_n \rightarrow a (n \rightarrow \infty)$$

and we say that (x_n) **converges to a as n goes to ∞** . A sequence (x_n) that is not convergent is called **divergent** and we say (x_n) **diverges**.

Proposition 5.3.1. The following statements are equivalent:

- (i) $\lim_{n \rightarrow \infty} x_n = a$.

- (ii) For each $U \in \mu(a)$, there is some $N := N(U)$ such that $x_n \in U$ for all $n \geq N$.
- (iii) For each $\epsilon > 0$, there is some $N := N(\epsilon)$ such that $x_n \in \mathbb{B}(a, \epsilon)$ for all $n \geq N$.

Bounded Sets

Definition 5.3.2. A subset $Y \subseteq X$ is called **d-bounded** or **bounded in** X (with respect to the metric d) if there is some $M > 0$ such that $d(x, y) \leq M$ for all $x, y \in Y$. In this circumstance the **diameter** of Y , defined by

$$\text{diam}(Y) := \sup_{x, y \in Y} d(x, y)$$

is finite. A sequence (x_n) is **bounded** if its image $\{x_n; n \in \mathbb{N}\}$ is bounded.

Proposition 5.3.2. Any convergent sequence is bounded.

Proof. Suppose that $x_n \rightarrow a$. Then there is some N such that $x_n \in \mathbb{B}(a, 1)$ for all $n \geq N$. It follows from the triangle inequality that

$$d(x_m, x_n) \leq d(x_m, a) + d(a, x_n) \leq 2, \quad m, n \in \mathbb{N}$$

Since there is also some $M \geq 0$ such that $d(x_j, x_k) \leq M$ for all $j, k \leq N$, we have $d(x_n, x_m) \leq M + 2$ for all $m, n \in \mathbb{N}$. ■

Uniqueness of the Limit

Proposition 5.3.3. Let (x_n) be convergent with limit a . Then a is the unique cluster point of (x_n) .

Corollary 5.3.1. The limit of a convergent sequence is unique.

Subsequence

Let $\varphi = (x_n)$ be a sequence in X and $\Phi: \mathbb{N} \rightarrow \mathbb{N}$ a strictly increasing function, then $\varphi \circ \Phi \in X^{\mathbb{N}}$ is called a **subsequence** of φ . Extending the notation $(x_n)_{n \in \mathbb{N}}$ introduced above for the sequence φ , we write $(x_{n_k})_{k \in \mathbb{N}}$ for the subsequence $\varphi \circ \Phi$ where $n_k := \Phi(k)$.

Proposition 5.3.4. If (x_n) is a convergent sequence with limit a , then each subsequence $(x_{n_k})_{k \in \mathbb{N}}$ of (x_n) is convergent with $\lim_{k \rightarrow \infty} x_{n_k} = a$.

Proposition 5.3.5. A point a is a cluster point of a sequence (x_n) if and only if there is some subsequence $(x_{n_k})_{k \in \mathbb{N}}$ of (x_n) which converges to a .

Chapter 6

Normed Vector Space

6.1 Norms

Definition 6.1.1 (Norm). Let E be a vector space over \mathbb{K} . A function $\|\cdot\|: E \rightarrow \mathbb{R}^+$ is called a **norm** if the following hold:

- $\|x\| = 0 \Leftrightarrow x = 0$.
- $\|\lambda x\| = |\lambda| \|x\|$, $x \in E$, $\lambda \in \mathbb{K}$ (positive homogeneity)
- $\|x + y\| \leq \|x\| + \|y\|$, $x, y \in E$ (triangle inequality). A pair $(E, \|\cdot\|)$ consisting of a vector space E and a norm $\|\cdot\|$ is called a **normed vector space**. If the norm is clear from context, we write E instead of $(E, \|\cdot\|)$.

Remark. Let $E := (E, \|\cdot\|)$ be a normed vector space.

(a) The function

$$d: E \times E \rightarrow \mathbb{R}^+, \quad (x, y) \mapsto \|x - y\|$$

is a metric on E , the **metric induced from the norm**. Hence any normed vector space is also a metric space.

(b) The **reversed triangle inequality** holds for the norm:

$$\|x - y\| \geq \left| \|x\| - \|y\| \right|, \quad x, y \in E$$

(c) All statements from previous chapter also hold in normed vector space.

Balls

For $a \in E$ and $r > 0$, we define the **open** and **closed balls** with center at a and radius r by

$$\mathbb{B}_E(a, r) := \mathbb{B}(a, r) := \{x \in E; \|x - a\| < r\}$$

and

$$\bar{\mathbb{B}}_E(a, r) := \bar{\mathbb{B}}(a, r) := \{x \in E; \|x - a\| \leq r\}.$$

These definitions agree with those for the metric space (E, d) when d is induced from norm. We also write

$$\mathbb{B} := \mathbb{B}(0, 1) = \{x \in E; \|x\| < 1\} \quad \text{and} \quad \bar{\mathbb{B}} := \bar{\mathbb{B}}(0, 1) = \{x \in E; \|x\| \leq 1\}$$

for the **open** and **closed unit balls** in E . We have

$$r\mathbb{B} = \mathbb{B}(0, r), \quad r\bar{\mathbb{B}} = \bar{\mathbb{B}}(0, r)$$

Bounded Sets

A subset X of E is called **bounded in E** (or **norm bounded**) if it is bounded in the induced metric space.

Remark. Let $E := (E, \|\cdot\|)$ be a normed vector space

- (a) $X \subseteq E$ is bounded if and only if there is some $r > 0$ such that $X \subseteq r\mathbb{B}$, that is, $\|x\| < r$ for all $x \in X$.
- (b) If X and Y are nonempty bounded subsets of E , then so are $X \cup Y$, $X + Y$ and λX with $\lambda \in \mathbb{K}$.

Example. (a) The absolute value $|\cdot|$ is a norm on the vector space \mathbb{K} .

(b) Let F be a subspace of a normed vector space $E := (E, \|\cdot\|)$. Then the restriction $\|\cdot\|_F := \|\cdot\||_F$ of $\|\cdot\|$ to F is a norm on F . Thus $F := (F, \|\cdot\|_F)$ is a normed vector space with this **induced norm**.

(c) Let $E_j, \|\cdot\|_j, 1 \leq j \leq m$, be normed vector space over \mathbb{K} . Then

$$\|x\|_\infty := \max_{1 \leq j \leq m} \|x_j\|_j, \quad x = (x_1, \dots, x_m) \in E := E_1 \times \dots \times E_m$$

defines a norm, called the **product norm**, on the product vector space E . The metric on E induced from this norm coincides with the product metric from 5.2 example (d), where d_j is the metric induced on E_j from $\|\cdot\|_j$.

(d) For $m \in \mathbb{N}^\times$, \mathbb{K}^m is a normed vector space with the **maximum norm**

$$|x|_\infty := \max_{1 \leq j \leq m} |x_j|, \quad x = (x_1, \dots, x_m) \in \mathbb{K}^m.$$

In this case $m = 1$.

6.2 The Space of Bounded Functions

Let X be a nonempty set and $(E, \|\cdot\|)$ a normed vector space. A function $u \in E^X$ is called **bounded** if the image of u in E is bounded. For $u \in E^X$, define

$$\|u\|_\infty := \|u\|_{\infty, X} := \sup_{x \in X} \|u(x)\| \in \mathbb{R}^+ \cup \infty$$

Remark. (a) For $u \in E^X$, the followings are equivalent:

- (i) u is bounded.
- (ii) $u(X)$ is bounded in E .
- (iii) There is some $r > 0$ such that $\|u(x)\| \leq r$ for all $x \in X$.
- (iv) $\|u\|_\infty < \infty$.

(b) Clearly $\text{id} \in \mathbb{K}^\mathbb{K}$ is not bounded, that is, $\|\text{id}\|_\infty = \infty$.

(b) shows that $\|\cdot\|_\infty$ may not be a norm on the vector space E^X when E is not trivial. We therefore set

$$B(X, E) := \{u \in E^X; u \text{ is bounded}\}$$

and call $B(X, E)$ the **space of bounded functions** from X to E .

Proposition 6.2.1. $B(X, E)$ is a subspace of E^X and $\|\cdot\|_\infty$ is a norm, called the **supremum norm**, on $B(X, E)$.

Remark. (a) If $X := \mathbb{N}$, then $B(X, E)$ is the normed vector space of bounded sequences in E . In the special case $E := \mathbb{K}$, $B(\mathbb{N}, \mathbb{K})$ is denoted by ℓ_∞ , that is,

$$\ell_\infty := \ell_\infty(\mathbb{K}) := B(\mathbb{N}, \mathbb{K})$$

is the normed **vector space of bounded sequences** with the supremum norm

$$\|(x_n)\|_\infty = \sup_{n \in \mathbb{N}} |x_n|, \quad (x_n) \in \ell_\infty$$

(b) If $X = \{1, \dots, m\}$ for some $m \in \mathbb{N}^\times$, then

$$B(X, E) = (E^m, \|\cdot\|_\infty)$$

6.3 Inner Product Spaces

Definition 6.3.1 (Inner Product Space). Let E be a vector space over the field \mathbb{K} . A function

$$(\cdot|\cdot): E \times E \rightarrow \mathbb{K}, \quad (x, y) \mapsto (x|y)$$

is called a **scalar product** or **inner product** on E if the following hold:

- $(x|y) = \overline{(y|x)}$, $x, y \in E$
- $(\lambda x + \mu y|z) = \lambda(x|z) + \mu(y|z)$, $x, y, z \in E$, $\lambda, \mu \in \mathbb{K}$.
- $(x|x) \geq 0$, $x \in E$, and $(x|x) = 0 \Leftrightarrow 0$.

A vector space E with a scalar product $(\cdot|\cdot)$ is called an **inner product space** and is written in $(E, (\cdot|\cdot))$.

Remark. (a) In the real case $\mathbb{K} = \mathbb{R}$, the first point can be written as

$$(x|y) = (y|x), \quad x, y \in E$$

In other words, the function is **symmetric** when E is a real vector space. In the case $\mathbb{K} = \mathbb{C}$, the function is said to be **Hermitian** when the first point holds.

(b) From the first two points it follows that

$$(x|\lambda y + \mu z) = \bar{\lambda}(x|y) + \bar{\mu}(x|z), \quad x, y, z \in E, \quad \lambda, \mu \in \mathbb{K},$$

that is, for each fixed $x \in E$, the function $(x|\cdot) : E \rightarrow \mathbb{K}$ is **conjugate linear**. If $\mathbb{K} = \mathbb{R}$, it is **bilinear**.

(c) $(x|0) = 0$ for all $x \in E$.

Let $m \in \mathbb{N}^\times$. For $x = (x_1, \dots, x_m)$ and $y = (y_1, \dots, y_m)$ in \mathbb{K}^m , define

$$(x|y) := \sum_{j=1}^m x_j \bar{y}_j$$

to be the **Euclidean Inner product** on \mathbb{K}^m .

The Cauchy-Schwarz Inequality

Theorem 6.3.1 (Cauchy-Schwarz Inequality). Let $(E, (\cdot|\cdot))$ be an inner product space. Then

$$|(x|y)|^2 \leq (x|x)(y|y) \quad x, y \in E$$

and the equality occurs if and only if x and y are linearly dependent.

Theorem 6.3.2. Let $(E, (\cdot|\cdot))$ be an inner product space and

$$\|x\| := \sqrt{(x|x)}, \quad x \in E$$

Then $\|\cdot\|$ is a norm on E , the **norm induced from the scalar product** $(\cdot|\cdot)$. A norm which is induced from a scalar product is also called a **Hilbert norm**.

Corollary 6.3.1. Let $(E, (\cdot|\cdot))$ be an inner product space. Then

$$|(x|y)| \leq \|x\| \|y\| \quad x, y \in E$$

6.4 Euclidean Spaces

Convention: Unless otherwise stated, we consider \mathbb{K}^m to be endowed with the Euclidean inner product $(\cdot|\cdot)$ and the induced norm

$$|x| := \sqrt{(x|x)} = \sqrt{\sum_{j=1}^m |x_j|^2} \quad x = (x_1, \dots, x_m) \in \mathbb{K}^m$$

the **Euclidean norm**. In the real case, we write also $x \cdot y$ for $(x|y)$.

We further define the norm

$$|x|_1 := \sum_{j=1}^m |x_j|, \quad x = (x_1, \dots, x_m) \in \mathbb{K}^m$$

Proposition 6.4.1. Let $m \in \mathbb{N}^\times$. Then

$$|x|_\infty \leq |x| \leq \sqrt{m} |x|_\infty, \quad \frac{1}{\sqrt{m}} |x|_1 \leq |x| \leq |x|_1, \quad x \in \mathbb{K}^m$$

Equivalent Norm

Let E be a vector space. Two norms $\|\cdot\|_1$ and $\|\cdot\|_2$ on E are **equivalent** if there is some $K \geq 1$ such that

$$\frac{1}{K} \|x\|_1 \leq \|x\|_2 \leq K \|x\|_1, \quad x \in E$$

In this case we write $\|\cdot\|_1 \sim \|\cdot\|_2$.

Remark. (a) \sim is an equivalence relation on the set of all norms of a fixed vector space.

(b) $\|\cdot\|_1 \sim \|\cdot\| \sim \|\cdot\|_\infty$ on \mathbb{K}^m .

(c) We write \mathbb{B}^m for the **real open Euclidean unit ball**, that is, $\mathbb{B}^m := \mathbb{B}_{\mathbb{R}}^m$ and \mathbb{B}_1^m and \mathbb{B}_∞^m for the unit balls in $(\mathbb{R}^m, |\cdot|_1)$ and in $(\mathbb{R}^m, |\cdot|_\infty)$ respectively. We have

$$\mathbb{B}^m \subseteq \mathbb{B}_\infty^m \subseteq \sqrt{m} \mathbb{B}^m \quad \mathbb{B}_1^m \subseteq \mathbb{B}^m \subseteq \sqrt{m} \mathbb{B}_1^m$$

(d) Let $E = (E, \|\cdot\|)$ be a normed vector space and $\|\cdot\|_1$ a norm on E which is equivalent to $\|\cdot\|$. Set $E_1 := (E, \|\cdot\|_1)$. Then

$$\mathcal{U}_E(a) = \mathcal{U}_{E_1}(a), \quad a \in E$$

that is, the set of neighborhoods of a depends only on the equivalence class of the norm. Equivalent norms produce the same set of neighborhoods.

Convergence in Product Spaces

Proposition 6.4.2. Let $m \in \mathbb{N}^\times$ and $x_n = (x_n^1, \dots, x_n^m) \in \mathbb{K}^m$ for $n \in \mathbb{N}$. Then the followings are equivalent:

1. The sequence $(x_n)_{n \in \mathbb{N}}$ converges to $x = (x^1, \dots, x^m)$ in \mathbb{K}^m .
2. For each $k \in \{1, \dots, m\}$, the sequence $(x_n^k)_{n \in \mathbb{N}}$ converges to x^k in \mathbb{K} .

Chapter 7

Infinite Limits

7.1 Convergence to $\pm\infty$

Sequences in \mathbb{R} can usually be considered to converge to $+\infty$ or $-\infty$ in the extended number line $\bar{\mathbb{R}}$. A subset $U \subseteq \bar{\mathbb{R}}$ is called a **neighborhood of ∞** (or of $-\infty$) if there is some $K > 0$ such that $(K, \infty) \subseteq U$ (or such that $(-\infty, -K) \subseteq U$). The set of neighborhoods of $\pm\infty$ is denoted by $\mathcal{U}(\pm\infty)$, that is,

$$\mathcal{U}(\pm\infty) := U \subseteq \bar{\mathbb{R}}; U \text{ is neighborhood of } \pm\infty$$

Now let (x_n) be a sequence in \mathbb{R} . Then $\pm\infty$ is called a **cluster point** (or **limit**) of (x_n) , if each neighborhood U of $\pm\infty$ contains infinitely many (or almost all) terms of (x_n) . If $\pm\infty$ is the limit of (x_n) , we usually write

$$\lim_{n \rightarrow \infty} x_n = \pm\infty \quad \text{or} \quad x_n \rightarrow \pm\infty \ (n \rightarrow \infty)$$

The sequence (x_n) **converges** in $\bar{\mathbb{R}}$ if there is some $x \in \bar{\mathbb{R}}$ such that $\lim_{n \rightarrow \infty} x_n = x$. The sequence (x_n) **diverges** in $\bar{\mathbb{R}}$, if it does not converge in $\bar{\mathbb{R}}$. With this definition, any sequence which converge in \mathbb{R} , also converges in $\bar{\mathbb{R}}$, and any sequence which diverges in $\bar{\mathbb{R}}$, also diverges in \mathbb{R} . On the other hand there are divergent sequences in \mathbb{R} which converge in $\bar{\mathbb{R}}$ (to $\pm\infty$). In this case the sequence is said to converge **improperly**.

Proposition 7.1.1. Every monotone sequence (x_n) in \mathbb{R} converges in $\bar{\mathbb{R}}$, and

$$\lim x_n = \begin{cases} \sup x_n; n \in \mathbb{N}, & \text{if } (x_n) \text{ is increasing,} \\ \inf x_n; n \in \mathbb{N}, & \text{if } (x_n) \text{ is decreasing.} \end{cases}$$

7.2 The Limit Superior and Limit Inferior

Definition 7.2.1. Let (x_n) be a sequence in \mathbb{R} . We can define two new sequences (y_n) and (z_n) by

$$y_n := \sup_{k \geq n} x_k := \sup x_k; k \geq n,$$

$$z_n := \inf_{k \geq n} x_k := \inf x_k; k \geq n.$$

Clearly (y_n) is increasing and (z_n) is decreasing. By the above proposition, these sequences converge in $\bar{\mathbb{R}}$:

$$\limsup_{n \rightarrow \infty} x_n := \overline{\lim}_{n \rightarrow \infty} x_n := \lim_{n \rightarrow \infty} (\sup_{k \geq n} x_k).$$

the **limit superior**, and

$$\liminf_{n \rightarrow \infty} x_n := \underline{\lim}_{n \rightarrow \infty} x_n := \lim_{n \rightarrow \infty} (\inf_{k \geq n} x_k).$$

the **limit inferior** of the sequences (x_n) . We also have

$$\limsup x_n = \inf_{n \in \mathbb{N}} (\sup_{k \geq n} x_k) \quad \text{and} \quad \liminf x_n = \sup_{n \in \mathbb{N}} (\inf_{k \geq n} x_k).$$

Theorem 7.2.1. Any sequence (x_n) in \mathbb{R} has a smallest cluster point x_* and a greatest cluster point x^* in $\overline{\mathbb{R}}$ and these satisfy

$$\liminf x_n = x_* \quad \text{and} \quad \limsup x_n = x^*$$

Theorem 7.2.2. Let (x_n) be a sequence in \mathbb{R} . Then

$$(x_n) \text{ converges in } \overline{\mathbb{R}} \Leftrightarrow \overline{\lim} x_n \leq \underline{\lim} x_n$$

When the sequence converges, the limit x satisfies

$$x = \lim x_n = \underline{\lim} x_n = \overline{\lim} x_n.$$

Theorem 7.2.3 (Bolzano-Weierstrass). Every bounded sequence in \mathbb{K}^m has a convergent subsequence, that is, a cluster point.

Chapter 8

Completeness

8.1 Cauchy Sequences

In the following $X = (X, d)$ is a metric space.

A sequence (x_n) in X is called a **Cauchy sequence** if, for each $\epsilon > 0$, there is some $N \in \mathbb{N}$ such that $d(x_n, x_m) < \epsilon$ for all $m, n \geq N$.

Similarly, if (x_n) is a sequence in a normed vector space $E = (E, \|\cdot\|)$, then (x_n) is a Cauchy sequence if and only if for each $\epsilon > 0$ there is some N such that $\|x_n - x_m\| < \epsilon$ for all $m, n \geq N$. In particular, Cauchy sequences in E are "translation invariant", that is, if (x_n) is a Cauchy sequence and a is an arbitrary vector in E , then the 'translated' sequence $(x_n + a)$ is also a Cauchy sequence. This implies that Cauchy sequences cannot be defined using neighborhoods.

Proposition 8.1.1. Every convergent sequence is a Cauchy sequence.

Proposition 8.1.2. Every Cauchy sequence is bounded.

Proposition 8.1.3. If a Cauchy sequence has a convergent subsequence, then it is itself convergent.

8.2 Banach Spaces

A metric space X is called **complete** if every Cauchy sequence in X converges. A complete normed vector space is called a **Banach space**.

Theorem 8.2.1. \mathbb{K}^m is a Banach space.

Theorem 8.2.2. Let X be a nonempty set and $E = (E, \|\cdot\|)$ a Banach space. Then $B(X, E)$ is also a Banach space.

Remark. (a) A direct consequence of the previous two theorems is that For every nonempty set X , $B(X, \mathbb{R})$, $B(X, \mathbb{C})$, and $B(X, \mathbb{K}^m)$ are Banach spaces.

(b) The completeness of a normed vector space E is invariant under changes to equivalent norms.

(c) A complete inner product space is called a **Hilbert space**.

Part III

Continuous Functions

Chapter 9

Continuity

9.1 Elementary Properties and Examples

Let $f: X \rightarrow Y$ be a function between metric spaces (X, d_X) and (Y, d_Y) . Then f is **continuous** at $x_0 \in X$ if, for each neighborhood V of $f(x_0)$ in Y , there is a neighborhood U of x_0 in X such that $f(U) \subseteq V$.

Hence to prove the continuity of f at x_0 , one supposes that an arbitrary neighborhood V of $f(x_0)$ is given and then shows that there is a neighborhood U of x_0 such that $f(U) \subseteq V$, that is, $f(x) \in V$ for all $x \in U$.

The function $f: X \rightarrow Y$ is **continuous** if it is continuous at each point of X . We say f is **discontinuous at x_0** if f is not continuous at x_0 . f is **discontinuous** if it is discontinuous at (at least) one point of X . The set of all continuous functions from X to Y is denoted $C(X, Y)$, a subset of Y^X .

Proposition 9.1.1. A function $f: X \rightarrow Y$ is continuous at $x_0 \in X$ if and only if, for each $\epsilon > 0$, there is some $\delta := \delta(x_0, \epsilon) > 0$ with the property that

$$d(f(x_0), f(x)) < \epsilon \text{ for all } x \in X \text{ such that } d(x_0, x) < \delta$$

Corollary 9.1.1. Let E and F be normed vector spaces and $X \subseteq E$. Then $f: X \rightarrow F$ is continuous at $x_0 \in X$ if and only if, for each $\epsilon > 0$, there is some $\delta := \delta(x_0, \epsilon) > 0$ satisfying that

$$\|f(x) - f(x_0)\|_F < \epsilon \text{ for all } x \in X \text{ such that } \|x - x_0\|_E < \delta$$

Example. In the following examples, X and Y are metric spaces.

- (a) The square root function $\mathbb{R}^+ \rightarrow \mathbb{R}^+$, $x \rightarrow \sqrt{x}$ is continuous.
- (b) The floor function $\lfloor \cdot \rfloor: \mathbb{R} \rightarrow \mathbb{R}$, $x \rightarrow \lfloor x \rfloor := \max\{k \in \mathbb{Z}; k \leq x\}$ is continuous at $x_0 \in \mathbb{R} \setminus \mathbb{Z}$ and discontinuous at $x_0 \in \mathbb{Z}$.
- (c) The **Dirichlet function** $f: \mathbb{R} \rightarrow \mathbb{R}$ defined by

$$f(x) := \begin{cases} 1, & x \in \mathbb{Q}, \\ 0, & x \in \mathbb{R} \setminus \mathbb{Q}, \end{cases}$$

is nowhere continuous, that is, it is discontinuous at every $x_0 \in \mathbb{R}$.

- (d) Suppose that $f: X \rightarrow \mathbb{R}$ is continuous at $x_0 \in X$ and $f(x_0) > 0$. Then there is a neighborhood U of x_0 such that $f(x) > 0$ for all $x \in U$.
- (e) A function $f: X \rightarrow Y$ is **Lipschitz continuous** with **Lipschitz constant** $\alpha > 0$ if

$$d(f(x), f(y)) \leq \alpha d(x, y), \quad x, y \in X$$

Every Lipschitz continuous function is continuous.

- (f) Any constant function $X \rightarrow Y$, $x \mapsto y_0$ is Lipschitz continuous.
- (g) The identity function $\text{id}: X \rightarrow X$, $x \mapsto x$ is Lipschitz continuous.
- (h) If E_1, \dots, E_m are normed vector spaces, then $E := E_1 \times \dots \times E_m$ is a normed vector space with respect to the product norm $\|\cdot\|_\infty$. The canonical projections

$$\text{pr}_k: E \rightarrow E_k, \quad x = (x_1, \dots, x_m) \mapsto x_k, \quad 1 \leq k \leq m,$$

are Lipschitz continuous. In particular, the projections $\text{pr}_k: \mathbb{K}^m \rightarrow \mathbb{K}$ are Lipschitz continuous.

- (i) Let E be a normed vector space. Then the norm function

$$\|\cdot\|: E \rightarrow \mathbb{R}, \quad x \mapsto \|x\|$$

is Lipschitz continuous.

- (j) If $A \subseteq X$ and $f: X \rightarrow Y$ is continuous at $x_0 \in A$, then $f|_A: A \rightarrow Y$ is continuous at x_0 . Here A has the metric induced from X .
- (k) Let $M \subseteq X$ be a nonempty subset of X . For each $x \in X$,

$$d(x, M) := \inf_{m \in M} d(x, m)$$

is called the **distance** from x to M . The **distance function**

$$d(\cdot, M): X \rightarrow \mathbb{R}, \quad x \mapsto d(x, M)$$

is Lipschitz continuous.

- (l) For any inner product space $(E, (\cdot|\cdot))$, the scalar product $(\cdot|\cdot): E \times E \rightarrow \mathbb{K}$ is continuous.
- (m) Let E and F be normed vector spaces and $X \subseteq E$. Then the continuity of $f: X \rightarrow F$ at $x_0 \in X$ is independent of the choice of equivalent norms on E and on F .
- (n) A function f between metric spaces X and Y is **isometric** (or an **isometry**) if $d(f(x), f(x')) = d(x, x')$ for all $x, x' \in X$, that is, if f 'preserves distances'. Such a function is Lipschitz continuous and is a bijection from X to its image $f(X)$. If E and F are normed vector spaces and $T: E \rightarrow F$ is linear, then T is isometric if and only if $\|Tx\| = \|x\|$ for all $x \in E$. If, in addition, T is surjective then T is an **isometric isomorphism** from E to F , and T^{-1} is also isometric.

9.2 Sequential Continuity

Definition 9.2.1 (Sequential Continuity). A function $f: X \rightarrow Y$ between metric spaces X and Y is called **sequentially continuous** at $x \in X$, if, for every sequences (x_k) in X such that $\lim_{x_k} = x$, we have $\lim_{f(x_k)} = f(x)$.

Theorem 9.2.1 (sequence criterion). Let X, Y be metric spaces. Then a function $f: X \rightarrow Y$ is continuous at x if and only if it is sequentially continuous at x .

Let $f: X \rightarrow Y$ be a continuous function between metric spaces. Then for any convergent sequence (x_k) in X we have

$$\lim f(x_k) = f(\lim x_k)$$

That's why we say 'continuous functions respect the taking of limits'.

9.3 Addition and Multiplication of Continuous Functions

Proposition 9.3.1. Suppose that X is a metric space, F is a normed vector space, and

$$f: \text{dom}(f) \subseteq X \rightarrow F, \quad g: \text{dom}(g) \subseteq X \rightarrow F$$

are continuous at $x_0 \in \text{dom}(f) \cap \text{dom}(g)$.

- $f + g$ and λf are continuous at x_0 .
- If $F = \mathbb{K}$, then $f \cdot g$ is continuous at x_0 .
- If $F = \mathbb{K}$ and $g(x_0) \neq 0$, then f/g is continuous at x_0 .

Corollary 9.3.1. (i) Rational functions are continuous.

(ii) Polynomials in n variables are continuous (on \mathbb{K}^n).

(iii) $C(X, F)$ is a subspace of F^X , the **vector space of continuous functions** from X to F .

Theorem 9.3.1 (continuity of compositions). Let X, Y and Z be metric spaces. Suppose that $f: X \rightarrow Y$ is continuous at $x \in X$, and $g: Y \rightarrow Z$ is continuous at $f(x) \in Y$. Then the composition $g \circ f: X \rightarrow Z$ is continuous at x .

Example. Let X be a metric space and E be a normed vector space.

Let $f: X \rightarrow E$ be continuous at x_0 . Then the **norm of** f ,

$$\|f\|: X \rightarrow \mathbb{R}, \quad x \mapsto \|f(x)\|,$$

is continuous at x_0 .

9.4 One-Sided Continuity

Let X be a subset of \mathbb{R} and $x_0 \in X$. For $\delta > 0$, the set $X \cap (x_0 - \delta, x_0]$ (or $X \cap [x_0, x_0 + \delta)$) is called a **left** (or **right**) δ -**neighborhood** of x_0 .

Let Y be a metric space. Then $f: X \rightarrow Y$ is **left** (or **right**) **continuous** at x_0 , if, for each neighborhood V of $f(x_0)$ in Y , there is some $\delta > 0$ such that $f(X \cap (x_0 - \delta, x_0]) \subseteq V$ (or $f(X \cap [x_0, x_0 + \delta)) \subseteq V$).

Chapter 10

The Fundamentals of Topology

10.1 Open Sets

Definition 10.1.1. Let $X := (X, d)$ be a metric space. An element a of a subset A of X is called an **interior point** of A if there is a neighborhood U of a such that $U \subseteq A$. The set A is called **open** if every point of A is an interior point.

Example. The open ball $\mathbb{B}(a, r)$ is open.

Remark. (a) The concepts "interior point" and "open set" depend on the surrounding metric space X . It is sometimes useful to make this explicit by saying that ' a is an interior point of A with respect to X ', or ' A is open in X '.

- (b) If A is open with respect to a particular norm, it is open with respect to all equivalent norms.
- (c) Every point in a metric space has an open neighborhood

Proposition 10.1.1. Let $\tau := \{O \subseteq X; O \text{ is open}\}$ be a family of open sets.

- (i) $\emptyset, X \in \tau$
- (ii) If $O_\alpha \in \tau$ for all $\alpha \in \mathcal{A}$, then $\bigcup_\alpha O_\alpha \in \tau$. That is, arbitrary unions of open sets are open.
- (iii) If $O_0, \dots, O_n \in \tau$, then $\bigcap_{k=0}^n O_k \in \tau$. That is, finite intersections of open sets are open.

Let M be a set and $\tau \subseteq \mathcal{P}(M)$, a set of subset satisfying (i)-(iii). Then τ is called a **topology** on M , and the elements of τ are called the **open sets** with respect to τ . The pair (M, τ) is called a **topological space**.

Remark. (a) Let $\tau \in \mathcal{P}(X)$ be the family of open sets. Then τ is called the **topology on X induced from the metric d** . If X is a normed vector space with metric induced from the norm, then τ is called the **norm topology**.

- (b) Let $(X, \|\cdot\|)$ be a normed vector space, and $\|\cdot\|_1$ a norm on X which is equivalent to $\|\cdot\|$. Let $\tau_{\|\cdot\|}$ and $\tau_{\|\cdot\|_1}$ be the norm topologies induced from $(X, \|\cdot\|)$ and $(X, \|\cdot\|_1)$. Then they coincide, that is, equivalent norms induce the same topology on X .

10.2 Closed Sets

A subset A of the metric space X is called **closed** in X if A^c is open in X .

Proposition 10.2.1. (i) \emptyset and X are closed.

(ii) Arbitrary intersections of closed sets are closed.

(iii) Finite unions of closed sets are closed.

Remark. • Infinite intersections of open sets need not be open.

• Infinite unions of closed sets need not be closed.

Definition 10.2.1. Let $A \subseteq X$ and $x \in X$. We call x an **accumulation point** of A if every neighborhood of x in X has a nonempty intersection with A . The element $x \in X$ is called a **limit point** of A if every neighborhood of x in X contains a point of A other than x . We set

$$\overline{A} := \{x \in X; x \text{ is an accumulation point of } A\}.$$

Clearly any element of A and any limit point of A is an accumulation point of A .

Proposition 10.2.2. Let A be a subset of a metric space X .

(i) $A \subseteq \overline{A}$

(ii) $A = \overline{A} \Leftrightarrow A$ is closed.

Proposition 10.2.3. An element x of X is a limit point of A if and only if there is a sequence (x_k) in $A \setminus \{x\}$ which converges to x .

Corollary 10.2.1. An element x of X is an accumulation point of A if and only if there is a sequence (x_k) in A such that $x_k \rightarrow x$.

Proposition 10.2.4. For $A \subseteq X$, the followings are equivalent:

- A is closed.
- A contains all its limit points.
- Every sequence in A which converges in X , has its limits in A .

10.3 The Closure and Interior of a Set

Definition 10.3.1. Let A be a subset of a metric space X . Define the **closure** of A by

$$\text{cl}(A) := \text{cl}_X(A) := \bigcap_{B \in M} B$$

with

$$M := \{B \subseteq X; B \supseteq A \text{ and } B \text{ is closed in } X\}.$$

Remark. The closure of A is the smallest closed set which contains A . Any closed set which contains A , also contains $\text{cl}(A)$.

Proposition 10.3.1. Let A be a subset of a metric space X . Then $\overline{A} = \text{cl}(A)$.

Corollary 10.3.1. Let A and B be subsets of X .

- (i) $A \subseteq B \Rightarrow \overline{A} \subseteq \overline{B}$.
- (ii) $\overline{(\overline{A})} = \overline{A}$.
- (iii) $\overline{A \cup B} = \overline{A} \cup \overline{B}$.

Definition 10.3.2 (Interior of a Set).

$$\text{int}(A) := \text{int}_X(A) := \bigcup \{O \subseteq A; O \text{ is open in } X\}$$

$\text{int}(A)$ is the largest open subset of A .

Definition 10.3.3.

$$\mathring{A} := \{a \in A; a \text{ is an interior point of } A\}.$$

Proposition 10.3.2. Let A be a subset of a metric space X . Then $\mathring{A} = \text{int}(A)$.

Corollary 10.3.2. Let A and B be subsets of X .

- (i) $A \subseteq B \Rightarrow \mathring{A} \subseteq \mathring{B}$.
- (ii) $(\mathring{A})^\circ = \mathring{A}$.
- (iii) $A \text{ is open} \Leftrightarrow A = \mathring{A}$.

10.4 The Hausdorff Condition

Definition 10.4.1. For a subset A of a metric space X , the (topological) **boundary** of A is defined by $\partial A := \overline{A} \setminus \mathring{A}$.

Proposition 10.4.1. Let A be a subset of X .

- ∂A is closed.
- x is in ∂A if and only if every neighborhood of x has nonempty intersection with both A and A^c .

The following proposition shows that, in metric spaces, any two distinct points have disjoint neighborhoods.

Proposition 10.4.2 (Hausdorff Condition). Let $x, y \in X$ be such that $x \neq y$. Then there are a neighborhood U of x and a neighborhood V of y such that $U \cap V = \emptyset$.

Corollary 10.4.1. Any one element subset of a metric space is closed.

10.5 A Characterization of Continuous Functions

Theorem 10.5.1. Let $f: X \rightarrow Y$ be a function between metric spaces X and Y . Then the followings are equivalent:

- (i) f is continuous.
- (ii) $f^{-1}(O)$ is open in X for each open set O in Y .
- (iii) $f^{-1}(A)$ is closed in X for each closed set A in Y .

Remark. According to this theorem, a function is continuous if and only if the preimage of any open set is open, if and only if the preimage of any closed set is closed. We denote the topology of a metric space X by τ_X , that is,

$$\tau_X := \{O \subseteq X; O \text{ is open in } X\}$$

Then,

$$f: X \rightarrow Y \text{ is continuous} \iff f^{-1}: \tau_Y \rightarrow \tau_X$$

10.6 Continuous Extensions

Let X and Y be metric spaces. Suppose that $D \subseteq X$, $f: D \rightarrow Y$ is continuous and $a \in X$ is a limit point of D . If D is not closed, then a may not be in D and so f is not defined at a . In this section we consider whether $f(a)$ can be defined so that f is continuous on $D \cup \{a\}$. If such extensions exist, then, for any sequence (x_n) in D which converges to a , $(f(x_n))$ converges to $f(a)$.

Remark. (a) The followings are equivalent:

- $\lim_{x \rightarrow a} f(x) = y$.
- For each neighborhood V of y in Y , there is a neighborhood U of a in X such that $f(U \cap D) \subseteq V$.

(b) If $a \in D$ is a limit point of D , then

$$\lim_{x \rightarrow a} f(x) = f(a) \iff f \text{ is continuous at } a.$$

10.7 Relative Topology

Let X be a metric space and Y a subset of X . Then Y is itself a metric space with respect to the metric $d_Y := d|_{Y \times Y}$ induced from X , and so 'open in (Y, d_Y) ' and 'closed in (Y, d_Y) ' are well-defined concepts.

Proposition 10.7.1. Let X be a metric space and $M \subseteq Y \subseteq X$. Then M is open (or closed) in Y if and only if M is open (or closed) in (Y, d_Y) .

Corollary 10.7.1. If $M \subseteq Y \subseteq X$, then M is open in Y if and only if $Y \setminus M$ is closed in Y .