

Analysis

Gaotang Li

July 22, 2022

Abstract

Notice that since in this course, the cross-referencing between theorems, lemmas, and propositions are quite complex and hard to keep track of, hence in this note, whenever you see a **!** over $=$, like $\stackrel{!}{=}$, then that **!** is *clickable*! It will direct you to the corresponding theorem, lemma, or proposition we're using to deduce that particular equality.

Notice that there are some proofs is **intended** left as assignments, and for completeness, I put them in **??**, use it in your **own risks**! You'll lose the chance to practice and really understand the materials.

Additionally, we'll use Folland[**folland1999real**] as our main text, while using Tao[**tao2013introduction**] and Axler[**axler2019measure**] as supplementary references.

This course is taken in Winter 2022, and the date on the covering page is the last updated time.

Contents

1	Foundations	2
1.1	Basics	2
1.2	σ -algebras	2

Chapter 1

Foundations

Lecture 1: Groups and Homomorphism

1.1 Basics

22 July. 01:17

Definition 1.1.1 (Group). A pair (G, \odot) consisting of a nonempty set G and an operation \odot is called a **group** if the following holds:

- G is closed under the operation \odot
- \odot is associative
- \odot has an identity element e
- Each $g \in G$ has an **inverse** $h \in G$ such that $g \odot h = h \odot g = e$

Definition 1.1.2 (Abelian group). A group G, \odot is called **commutative** or **Abelian** if \odot is a commutative operation on G .

Remark. Let $G = (G, \odot)$

- (a) the identity element e is unique
- (b) Each $g \in G$ has a unique inverse which we denote by g^b . In particular $e^b = e$.
- (c) For each $g \in G$, we have $(g^b)^b = g$.
- (d) For arbitrary group elements g and h , $(g \odot h)^b = h^b \odot g^b$

Example. (a) Let $G := \{e\}$ be a one element set. Then $\{G, \odot\}$ is an Abelian group, the **trivial group**, with the (only possible) operation $e \odot e = e$.

- (b) Let X be a nonempty set, and S_X be the set of all bijections from X to itself. Then $S_X := (S_X, \circ)$ is a group with identity element id_X when \circ denotes the composition of functions. Further, the inverse function f^{-1} is the inverse of $f \in S_X$ in the group. When X is finite, the element of S_X are called permutations and S_X is called the **permutation group** of X .

(c)

1.2 σ -algebras

We start from the definition of the most fundamental element in measure theory.

Definition 1.2.1 (σ -algebra). Let X be a set. A collection \mathcal{A} of subsets of X , i.e., $\mathcal{A} \subset \mathcal{P}(X)$ is called a σ -algebra on X if

- $\emptyset \in \mathcal{A}$.
- \mathcal{A} is closed under complements. i.e., if $A \in \mathcal{A}$, $A^c = X \setminus A \in \mathcal{A}$.
- \mathcal{A} is closed under countable unions. i.e., if $A_i \in \mathcal{A}$, then $\bigcup_{i=1}^{\infty} A_i \in \mathcal{A}$.
- wd

Remark. There are some easy properties we can immediately derive.

- $X \in \mathcal{A}$ from $X = X \setminus \underbrace{\emptyset}_{\in \mathcal{A}}$ and \mathcal{A} is closed under complement.
- $\bigcap_{i=1}^{\infty} A_i = \left(\bigcup_{i=1}^{\infty} A_i^c \right)^c$, namely \mathcal{A} is closed under countable intersections.
- $A_1 \cup A_2 \cup \dots \cup A_n = A_1 \cup A_2 \cup \dots \cup A_n \cup \emptyset \cup \emptyset \cup \dots$, hence \mathcal{A} is closed under finite unions and intersections.

Note. The [definition of \$\sigma\$ -algebra](#) should remind us the definition of topological basis, and this is indeed the case. We can consider a topological space and put some structure on the [\$\sigma\$ -algebra](#) \mathcal{A} , which gives us the following.