

A preliminary comparison of CrIS processing algorithms after the SNPP CrIS MWIR anomaly

H. E. Motteler, L. L. Strow

UMBC Atmospheric Spectroscopy Lab
Joint Center for Earth Systems Technology

June 9, 2019


Introduction

- ▶ We present a preliminary comparison of NPP CrIS high res products for the NOAA A4, UMBC CCAST A4, and UMBC CCAST reference calibration algorithms, before and after the NPP MW anomaly.
- ▶ Complex residuals for the NOAA product are slightly larger before and significantly larger after the anomaly.
- ▶ Differences in the NOAA and UMBC CCAST processing include
 - ▶ CCAST uses cosine apodization of the extended point set outside the user-grid OPD, rather than truncation or circular shift
 - ▶ CCAST uses “small n” psinc (psinc at the decimated sensor grid) for both the ILS and SA matrix
 - ▶ CCAST spectral-space processing filters for the reference algorithm have a slightly narrower passband and more gradual rolloff than the A4 filters. The difference is greatest at the low end of the LW band.
- ▶ The CCAST reference and NOAA A4 calibration equations and CCAST apodization are described in an appendix.


Test Data

- ▶ NOAA data are GCRSO-SCRIF HDF5 files from NOAA CLASS. CCAST data are from our regular production runs, with the A4 algorithm selectable as an option.
- ▶ CCAST uses 45-scan and NOAA 60-scan granules. For these preliminary tests we simply took the intersection of scans for two overlapping granules, one pair before and one pair after the anomaly.
- ▶ The pre-anomaly test set consists of 1110 obs from relatively warm granules, midpoint time 03-Mar-2019 04:07:11. The set is divided into even and odd sweeps (even and odd FORs) and broken out by FOV.
- ▶ The post-anomaly test set consists of 1260 obs from another pair of warm granules, midpoint time 21-Apr-2019 04:08:29. The set is also divided into even and odd sweeps and broken out by FOV.
- ▶ We show representative cases for corner, side, and center FOVs, for both sweep directions.

LW FOV 1 Sweep 1 Complex Residuals




Pre-anomaly complex residual mean and standard deviation. The NOAA complex residual is slightly larger.




Post-anomaly complex residual mean and standard deviation. The NOAA complex residual is significantly larger.

LW FOV 2 Sweep 1 Complex Residuals




Pre-anomaly complex residual mean and standard deviation. The NOAA complex residual is significantly larger.




Post-anomaly complex residual mean and standard deviation. The NOAA complex residual is significantly larger.

LW FOV 5 Sweep 1 Complex Residuals




Pre-anomaly complex residual mean and standard deviation. The NOAA complex residual is slightly larger.




Post-anomaly complex residual mean and standard deviation. The NOAA residual is generally larger.

LW FOV 1 Sweep 1 BT Spectra




Pre-anomaly mean BT spectra for NOAA A4, CCAST ref, and CCAST A4. The CCAST A4 and NOAA A4 algorithms are in good agreement.




Post-anomaly mean BT spectra for NOAA A4, CCAST ref, and CCAST A4. The differences suggest high frequency components with different periods, that is.

LW FOV 1 Sweep 1 Sarta Residuals




Pre-anomaly mean BT spectra for NOAA A4, CCAST ref, and CCAST A4. The CCAST A4 and NOAA A4 algorithms are in good agreement.




Post-anomaly mean BT spectra for NOAA A4, CCAST ref, and CCAST A4. The differences suggest high frequency components with different periods, that is.

LW FOV 2 Sweep 1 BT Spectra




Pre-anomaly mean BT spectra.
The CCAST A4 and NOAA A4
algorithms are in good agreement.




Post-anomaly mean BT spectra.
The differences suggest high
frequency components with different
periods, that is, different ringing.

LW FOV 2 Sweep 1 Sarta Residuals




Pre-anomaly mean BT spectra.
The CCAST A4 and NOAA A4
algorithms are in good agreement.




Post-anomaly mean BT spectra.
The differences suggest high
frequency components with different
periods, that is, different ringing.

LW FOV 5 Sweep 1 BT Spectra




Pre-anomaly mean BT spectra.
The CCAST and NOAA algorithms
are in relatively good agreement.




Post-anomaly mean BT spectra.
The differences suggest high
frequency components with different
periods.

LW FOV 5 Sweep 1 Sarta Residuals



Pre-anomaly mean BT spectra.
The CCAST and NOAA algorithms
are in relatively good agreement.



Post-anomaly mean BT spectra.
The differences suggest high
frequency components with different
periods.

Conclusions

- ▶ Complex residuals for the NOAA product are slightly larger before and significantly larger after the anomaly.
- ▶ This may be due to the mild cosine apodization working better for the case of a significant ZPD shift.
- ▶ Because the complex residuals are smaller, we suspect the CCAST BT spectra are closer to reference truth after the anomaly. But this needs to be verified.
- ▶ The slightly larger NOAA complex residuals before the anomaly are puzzling, we do not recall seeing this in earlier tests.
- ▶ The CCAST complex residual standard deviation is larger than NOAA for the SW side and corner FOVs. This may be due to the different forms of the SA matrix.
- ▶ The next step is comparison with calculated reference truth, from clear matchups.