



ಸುಶ್ರುತ ಆಯುರ್ವೇದ ವೈದ್ಯಕೀಯ ಕಾಲೇಜು ಮತ್ತು ಆಸ್ಪತ್ರೆ

Sushrutha Ayurvedic Medical College & Hospital

(Recognized by NCISM, Ministry of AYUSH, New Delhi, Affiliated to RGUHS, Bengaluru, Karnataka)

Mr VENKATESH

Age : 54 Year(s) Male

KPME Registration : BLU01220AYHL4

APL Code : APD-KA-047

Ref. No.

LAB REPORT

Customer : sampig lab

Sample Type : CITRATED PLASMA

SlID : 8890126

Collected on : 2025-05-15 00:00

Regd on : 2025-05-15 17:31

Reported On : 2025-05-15 18:11



HAEMATOLOGY

Test Description

Prothrombin Time (PT)

(Rothmund Clotting)

Prothrombin Time - Patient Value

18.3

Sec

11 - 16

Prothrombin Time - Control Value

13.5

Sec

Prothrombin Time - Ratio

1.3

Prothrombin Time - INR

1.35

Oral Anticoagulant

Therapeutic Range: 2.0 -3.5

Interpretation of PT and PTT in Patients with a Bleeding or Clotting Syndrome

PT RESULT	PTT RESULT	COMMON CONDITION PRESENT
Prolonged	Normal	Liver disease, decreased vitamin K, decreased or defective factor VII
Normal	Prolonged	Decreased or defective factor VIII, IX, or XI, von Willebrand disease or lupus anticoagulant present
Prolonged	Prolonged	Decreased or defective factor I, II, V or X, severe liver disease, disseminated intravascular coagulation (DIC)
Normal	Normal or slightly prolonged	May indicate normal hemostasis; however PT and PTT can be normal in conditions such as mild deficiencies in other factors and mild form of von Willebrand disease. Further testing may be required to diagnose these conditions

Prothrombin Time (PT) and its derived measures of prothrombin ratio (PR) and international normalized ratio (INR) are measures of the extrinsic pathway.

P.VENKATESHWARLU

Group Leader



DR. P PRASANNA B.K

MD PATHOLOGIST

Address : Prashanti Kuteeram, Jodi Bingsipura, Jigani Hobli, Anakal Taluk, Bengaluru - 560 105

Phone Number : 8277582990/8712367629

Lab Reception : 8277582990/8712367629



ಸುಶ್ರುತ ಆಯುರ್ವೇದ ವೈದ್ಯಕೀಯ ಕಾಲೇಜು ಮತ್ತು ಆಸ್ಪತ್ರೆ
Sushrutha Ayurvedic Medical College & Hospital

(Recognized by NCISM, Ministry of AYUSH, New Delhi, Affiliated to RGUHS, Bengaluru, Karnataka)

KPME Registration : BLU01220AYHL4



LAB REPORT

Name : Mr.VENKATESH
Age : 54Years
Gender : Male
Referred By : Dr.SOWMYA S B

Patient ID : 5272
Report No : SAMC05255273
Bill Date : 15-May-2025 12:22 pm
Rep Date : 15-May-2025 3:33 pm



Test Name

Result

Unit

Reference Range

HbsAg Card

Method : Immunochromatography

SEROLOGY
NEGATIVE

Negative

Importance of Hepatitis Marker

Hepatitis B is caused by the hepatitis B virus (HBV). HBV is a noncytopathic virus. This means that the virus itself does not cause direct damage to liver cells. Instead, it is the immune systems aggressive response to the virus that usually leads to inflammation and damage to the liver (hepatitis). However, HBV can cause damage to the genetic material inside liver cells. This can lead to liver cancer which, like hepatitis, can also be fatal.

It is present in the sera of patients with viral hepatitis B (with or without clinical symptoms). Patients who developed antibodies against HBsAg (anti-HBsAg sero conversion) are usually considered non-infectious. HBsAg detection by immunoassay is used in blood screening, to establish a diagnosis of hepatitis B infection in the clinical setting (in combination with other disease markers) and to monitor antiviral treatment. Positive HBsAg tests can be due to recent vaccination against Hepatitis B virus but this positivity is unlikely to persist beyond 14 days post-vaccination.

— End Of Report —

Authorized Sign