

```
In [1]: import numpy as np
import pandas as pd
```

```
In [20]: X = [9.0,8.0,9.3,9.4,3.3,2.2,3.2,1.1]
Y = [1,1,1,1,0,0,0,0]

dataset = pd.DataFrame(data={
    'X' : X,
    'Y' : Y
})
```

```
In [21]: dataset
```

```
Out[21]:
```

	X	Y
0	9.0	1
1	8.0	1
2	9.3	1
3	9.4	1
4	3.3	0
5	2.2	0
6	3.2	0
7	1.1	0

```
In [22]: dataset['XY'] = dataset['X'] * dataset['Y']
dataset['X2'] = dataset['X'] ** 2
```

```
In [23]: dataset
```

```
Out[23]:
```

	X	Y	XY	X2
0	9.0	1	9.0	81.00
1	8.0	1	8.0	64.00
2	9.3	1	9.3	86.49
3	9.4	1	9.4	88.36
4	3.3	0	0.0	10.89
5	2.2	0	0.0	4.84
6	3.2	0	0.0	10.24
7	1.1	0	0.0	1.21

```
In [24]: n = len(dataset)
sum_x = dataset['X'].sum()
sum_y = dataset['Y'].sum()
sum_xy = dataset['XY'].sum()
sum_x2 = dataset['X2'].sum()
sum_X_h2 = sum_x ** 2
```

```
In [25]: n, sum_x , sum_y , sum_xy , sum_x2 , sum_X_h2
```

```
Out[25]: (8, 45.5, 4, 35.7, 347.03000000000003, 2070.25)
```

```
In [26]: numerator_m = (n*(sum_xy)) - (sum_x*sum_y)
numerator_m
```

```
Out[26]: 103.60000000000002
```

```
In [27]: denominator_m = (n*(sum_x2) - sum_X_h2)
denominator_m
```

```
Out[27]: 705.9900000000002
```

```
In [28]: m = numerator_m / denominator_m
```

```
In [29]: m
```

```
Out[29]: 0.14674428816272184
```

```
In [30]: numerator_b = sum_y -(m * sum_x)
denominator_b =n

b = numerator_b/denominator_b
b
```

```
Out[30]: -0.33460813892548047
```

```
In [31]: m,b
```

```
Out[31]: (0.14674428816272184, -0.33460813892548047)
```

```
In [32]: X_cap = [m*X + b for X in dataset['X']]
X_cap
```

```
Out[32]: [0.9860904545390161,
0.8393461663762942,
1.0301137409878327,
1.0447881698041048,
0.14964801201150157,
-0.011770704967492385,
0.13497358319522945,
-0.17318942194648643]
```

```
In [33]: y = [1 / (1 + np.exp(-Xcap)) for Xcap in X_cap]
```

In [34]: y

Out[34]: [0.7283150226716693,  
0.6983274932179059,  
0.7369379463021366,  
0.7397728295238472,  
0.537342340327484,  
0.4970573577331819,  
0.5336922612250837,  
0.4568105447007532]

```
In [35]: result = []
for yvals in y:
    if yvals >= 0.5:
        result.append(1)
    else:
        result.append(0)
```

In [36]: result

Out[36]: [1, 1, 1, 1, 1, 0, 1, 0]

```
In [37]: result =[1 if yvals >=0.5 else 0 for yvals in y]
```

In [38]: result

Out[38]: [1, 1, 1, 1, 1, 0, 1, 0]

```
In [39]: dataset['RESULT'] = result
```

In [40]: dataset

Out[40]:

	X	Y	XY	X2	RESULT
0	9.0	1	9.0	81.00	1
1	8.0	1	8.0	64.00	1
2	9.3	1	9.3	86.49	1
3	9.4	1	9.4	88.36	1
4	3.3	0	0.0	10.89	1
5	2.2	0	0.0	4.84	0
6	3.2	0	0.0	10.24	1
7	1.1	0	0.0	1.21	0

In [41]: #accuracy

```
In [44]: correct = 0
for key,result in zip(dataset['Y'],dataset['RESULT']):
    if key == result:
        correct += 1
    pass
```

```
In [45]: correct/n
```

```
Out[45]: 0.75
```

```
In [47]: #another code in accuracy correct
correct = 0 # Inilizing the correct with 0
values = dataset[['Y','RESULT']].values
for (key,result) in values:
    if key == result:
        correct += 1
    pass
```

```
In [48]: correct / n
```

```
Out[48]: 0.75
```

```
In [53]: #another code in accuracy correct
correct = 0 # Inilizing the correct with 0
keys = dataset['Y']
results = dataset['RESULT']
for (key,result) in zip(keys,results):
    # print(f"key {key}, result {result}")
    if key == result:
        print(f" EQUAL key {key}, result {result}")
        correct += 1
        print(f"correct {correct}")
    pass
```

```
    EQUAL key 1, result 1
correct 1
    EQUAL key 1, result 1
correct 2
    EQUAL key 1, result 1
correct 3
    EQUAL key 1, result 1
correct 4
    EQUAL key 0, result 0
correct 5
    EQUAL key 0, result 0
correct 6
```

```
In [54]: correct/n
```

```
Out[54]: 0.75
```

```
In [ ]:
```

