

Arithmetic Expression Geometry I: Foundations

Sequential Histories, Projective Semantics, and Zero Geometry

DeepSeek Harness Agent*

model `deepseek-v4-flash-vision-exp`

Mingli Yuan

Swarma Research

Beijing 100083, China

September 22, 2026

Draft manuscript under mathematical review.

Draft notice. This manuscript is a working draft in the editorial and publication sense. Here “draft” is not a mathematical claim-status label and does not alter the status of any theorem, proposition, computation, conjecture, or open problem. The geometric chapters that earlier versions of this paper contained are now developed in Paper 0; the migration is recorded in `governance/00b-paper-0-geometric-foundation-amendment.md`.

Abstract

Paper 0 constructs arithmetic expression spaces, the arithmetic motions on them, ripple geometry, the accumulative commutative space, and the contact structure of arithmetic tearing. The present paper develops the layer that those constructions presuppose: the syntax of process. We classify the finite binary expression trees whose internal operations have a unique legal evaluation order; they are exactly the trees whose internal vertices form a single spine. Marking an accumulator turns such a tree into a chronological word of one-hole contexts, with operand-slot chirality kept distinct from temporal reversal and from functional inversion. Non-degenerate bilateral contexts act projectively on $\mathbb{P}^1(K)$ and generate $\mathrm{PGL}_2(K)$ over an arbitrary field; those fixing infinity form the affine, or Borel, sector, and the same proof gives the rank-one Bruhat decomposition. A three-letter sublanguage over $\mathbb{Q}(\sqrt{2})$ realizes the $q = 4$ Hecke group and yields an explicit marked history that is projectively trivial while being ordinarily inadmissible, which is why ordinary admissibility is kept separate from projective non-degeneracy. The paper then develops the zero theory of regular arithmetic expression spaces: the zero locus is a smooth curve; a connected complete boundaryless regular space splits as $Z(a) \times \mathbb{R}$; branching zero networks are therefore excluded in the regular category; and a smooth parameter family with spatially regular zeros has a submersive total zero set, subject to a stated properness warning. A singular arithmetic expression space is defined, and one isolated-zero model is verified.

Keywords: arithmetic expressions; evaluation histories; sequential trees; operand-slot chirality; projective linear group; Hecke group; regular zero loci; singular arithmetic expression spaces

*Correspondence and responsibility for this manuscript rest with the second author, Mingli Yuan (mingli@swarma.org). The first author is an artificial-intelligence research agent (model `deepseek-v4-flash-vision-exp`, running on the DeepSeek Harness); it is not a natural person, cannot sign a copyright transfer, and cannot serve as corresponding author. The author order was fixed by the authors and is not a measure of human contribution.

Contents

Interface: From Paper 0 to Paper I	4
1 Introduction	4
1.1 Sequential histories	5
1.2 Why bilateral completion matters	5
1.3 Main results	6
1.4 Scope and relation to standard geometry	6
1.5 Organization	7
2 Sequential Arithmetic Histories	7
2.1 Expression trees and dependency	7
2.2 One-hole contexts and the marked accumulator	9
2.3 Composition and ordinary evaluation	10
2.4 Mirror, temporal reversal, and path inverse	11
2.5 Information retained and information forgotten	12
2.6 Representative chirality patterns	12
3 Projective Semantics and the Affine Sector	13
3.1 Ordinary domains and projective continuation	13
3.2 Projective evaluation of a chronological history	14
3.3 Generation of the projective group	15
3.4 A distinguished arithmetic sublanguage: the $q=4$ Hecke group	16
3.5 The affine syntactic sector	17
3.6 The positive real component	18
3.7 Bruhat placement and the projective infinitesimal direction	19
4 The Regular AES Interface	19
5 Zero Loci and Singular Expression Spaces	20
5.1 Regular zero loci	20
5.2 Complete regular splitting and a global no-go theorem	21
5.3 Singular arithmetic expression spaces	22
5.4 A verified isolated-zero singular model	23
5.5 Smooth parameter families and their total zero sets	24
6 Conclusions and Research Interfaces	25
6.1 Results proved here	25
6.2 What moved to Paper 0	26
6.3 Interface to Paper II	27
6.4 Interface to Paper III	27
6.5 Interface to Paper IV	27
6.6 Open foundational problems	27
A Composition, action, and domain conventions	28
A.1 Chronological words and functional composition	28
A.2 Column-vector projective action	29
A.3 Ordinary and projective domains	29

A.4	Mirror, reversal, and inverse notation	29
A.5	Notation conversion summary	30
B	Equality and elementary neutrality examples	30
B.1	Same value, different operators	30
B.2	Same operator, different histories	31
B.3	Same charge, nonzero relative torsion	31
B.4	Zero endpoint defect, nontrivial history	31
B.5	Open torsion and closed-loop drift	31

Interface: From Paper 0 to Paper I

Paper 0 is the geometric foundation of this series. It constructs the space in which the elementary arithmetic operations act as motions, the dual reading of those motions, the commutativization against which their order defect is measured, and the contact structure of that defect. It fixes explicit left- and right-expanded conventions, proves the distributive expansion that relates two expansions sharing an operator, and treats reciprocals, poles, the projective point at infinity, affine and reciprocal fixed points, and the common 2×2 matrix representation.

The present paper begins one level earlier and stays there. Its first problem is intrinsic: which finite binary expression trees have a unique legal order of internal evaluation, including mixed operand-slot patterns? The answer is the single-spine classification. Choosing an accumulator then gives the marked spinal histories used throughout this paper. We retain that terminology and formalism unchanged.

The division of labour is therefore

Paper 0: expansions, arithmetic motions, the models $\mathfrak{E}_0$ and $\mathfrak{E}_1$,
ripple geometry, affine cocycles, ACS and tearing, contact structure,

Paper I: intrinsic sequential histories, bilateral projective evaluation, the Hecke sublanguage, zero geometry.

This is the reverse of the arrangement that held before the amendment recorded in [governance/00b-paper-0-geomet](#): the geometric chapters that formerly stood here are now developed there, at the same labels and with the same proofs. Paper I repeats exactly one imported definition — the regular AES and its canonical frame, in Section 4 — because the zero theorems below are stated in those terms; every other result of Paper 0 is cited rather than repeated.

Two conventions are shared and used identically in both papers. A pole or projective value at infinity never certifies ordinary admissibility of a division step; that distinction remains active here. And a historical word such as “left-expanded” is meaningful only under the explicit operand-slot recursion fixed in Paper 0.

1 Introduction

An arithmetic expression has several mathematically distinct lives. It is a finite syntax tree; it may be equipped with a legal order of evaluation; a chosen running value turns that order into a marked history; the history induces an operator; and the operator sends an initial value to an endpoint. Ordinary arithmetic usually passes through these levels as quickly as possible. AEG instead asks what is lost at each passage.

The guiding structure of this paper is

$$\begin{aligned} \text{unmarked sequential tree} &\xrightarrow{\text{choose an accumulator}} \text{marked planar sequential tree}, \\ \text{marked planar sequential tree} &\longleftrightarrow \text{bounded marked history}, \\ \text{bounded marked history} &\longrightarrow \text{operator} \longrightarrow \text{endpoint value}, \\ \text{bounded marked history} &\longrightarrow \text{charge shadow}. \end{aligned} \tag{1}$$

The first arrow denotes a choice, not a canonical map. When the planar structure, mark, operation labels, constants, and slots are retained, the double arrow is the bijection proved in Section 2. The two arrows leaving a history are separate forgetful maps: charge is not a function of the evaluated operator, and an endpoint is not a function of charge alone. Different histories may nevertheless induce the same operator, and different operators may agree at one chosen initial value. The




Figure 1: The process-to-result structure. Choosing an accumulator marks an unmarked sequential tree; a fully marked planar tree and a bounded marked history determine one another. From the history, operator evaluation and the charge map are separate information-losing branches, while an endpoint is obtained by applying the operator to an initial value.

geometry belongs primarily to these intermediate levels, before endpoint evaluation has erased the order of construction.

Figure 1 renders this structure explicitly. It is a map of the distinctions used in the paper, not a claim that any forgetful arrow is injective.

1.1 Sequential histories

Not every expression tree determines a unique history. Independent subexpressions may be evaluated in either order, so a single tree can have many legal linear extensions of its dependency order. Section 2 isolates the precise exception. For a finite binary tree, uniqueness of the internal evaluation order is equivalent to the internal vertices forming one chain, or equivalently one spine. A marked accumulator on that spine determines a word of one-hole contexts

$$C_{\omega,c}^{(1)}[z] = \omega(z, c), \quad C_{\omega,c}^{(2)}[z] = \omega(c, z).$$

The superscript records the operand slot occupied by the current value. It is not a time direction. This distinction prevents three operations that are often blurred in informal tree pictures—planar mirror, temporal reversal, and path inverse—from being identified.

This use of syntax is modest. Formal generation and rewriting have long histories in logic and theoretical computer science; classical points of reference include Post’s work on combinatorial reduction and Chomsky’s generative hierarchies [3, 1]. Here the purpose is different: the sequential fragment supplies a controlled domain on which evaluation can be treated as a path without pretending that every arithmetic tree has a unique path.

1.2 Why bilateral completion matters

Earlier affine calculations use the current value in one distinguished operand slot. That fragment is closed under translations and nonzero scalings and is therefore naturally represented by the affine group of the line. Once both operand slots are allowed, the context $z \mapsto c/z$ introduces inversion. The correct algebraic ambient space is then the projective line and its fractional linear transformations.

Section 3 proves, over an arbitrary field K , that the non-degenerate bilateral contexts generate $\text{PGL}_2(K)$. The affine transformations are exactly the stabilizer of the projective point at infinity. Thus the continuous geometry is neither discarded nor promoted to the whole theory: it is identified precisely as the affine sector of a larger projective semantics. The geometric reading of these operators

— addition and multiplication as motions of an arithmetic expression space, and second-slot division as the projective step — is developed in Paper 0 and used here only at the operator level.

Inside that larger semantics, the two contexts

$$T(z) = z + \sqrt{2}, \quad J(z) = -\frac{1}{z}$$

generate the $q = 4$ Hecke group. Section 3 verifies the relation $(JT)^4 = 1$ directly from the context matrices and formulates the resulting interface from arithmetic words to group operators and then to cell cosets. This restricted alphabet is important because its hyperbolic cellulations are produced by arithmetic operator semantics rather than imposed after endpoint evaluation. The interface is deliberately not a bijection: relations identify different histories, and stabilizers identify different operators at the level of cells or endpoints.

Projective continuation does not repair an inadmissible ordinary computation. For example, c/z has a well-defined projective value at $z = 0$, namely infinity, while ordinary division by zero remains undefined. We keep the ordinary domain and the projective extension as separate data throughout.

1.3 Main results

The first group of results is combinatorial and syntactic. The trees with a unique legal internal evaluation order are exactly those whose internal vertices form a single spine; marking a leaf at the innermost vertex turns such a tree into a bounded marked spinal history, and the two constructions are mutually inverse when planar data are retained. Operand-slot chirality is part of that data and is kept distinct from chronological direction.

The second group is projective. The non-degenerate bilateral contexts generate $\mathrm{PGL}_2(K)$ for every field K , the contexts fixing infinity form the affine sector, and the proof yields the rank-one Bruhat decomposition. The $\mathbb{Q}(\sqrt{2})$ -sublanguage generated by translation by $\sqrt{2}$ and by inversion carries the order-four relation $(JT)^4 = 1$, realized by an explicit eight-step marked history that is projectively trivial and ordinarily inadmissible at one initial value. That example is the reason this paper keeps ordinary admissibility as a separate hypothesis from projective non-degeneracy.

The third group is the zero theory of regular arithmetic expression spaces. Section 4 restates the definition imported from Paper 0; Section 5 proves that the zero locus of a regular assignment is a smooth curve, that a complete connected boundaryless regular AES splits as $Z(a) \times \mathbb{R}$, that this forbids branching or multiple zero components in the regular category, and that a smooth parameter family with spatially regular zeros has a submersive total zero set — with the properness warning that prevents the last statement from being read as triviality. The section then defines singular arithmetic expression spaces and verifies one isolated-zero model in detail.

Everything else that earlier versions of this paper developed — the two affine translation coordinates, the continuous arithmetic motion, the basic hyperbolic model $\mathfrak{E}_0$, the accumulated commutative space, tearing, and the contact structure — now belongs to Paper 0, where it is proved at the same labels. This paper cites those results rather than restating them, with the sole exception of the regular-AES interface needed to state its own zero theorems.

1.4 Scope and relation to standard geometry

Several ingredients used here are standard: finite posets and their linear extensions, fractional linear transformations, and the regular-value theorem. The mathematical contribution sought in Paper I is their history-sensitive arithmetic organization and the exact interfaces among them.

Four boundaries are enforced. First, the geometric construction of arithmetic expression spaces, their motions, their commutativization, and their contact structure belongs to Paper 0. Second, a

horizontal contact distribution does not by itself select a unique complex structure, so arithmetic holomorphic and harmonic theories are deferred to Paper II; this includes any automorphic-function theory built on the G_4 action. Third, the regular-value theorem is only the starting point for singular families: pulling analytic real loci back to the Hecke cellulation, and developing their multiple zeros, discriminants, tubes, braids, and knots, belong to Paper III. Fourth, projective bivaluations, homogeneous quotient towers, and condensation belong to Paper IV, and no computational lower bound follows from noncommutativity or negative curvature alone.

1.5 Organization

Section 2 develops sequential trees and marked histories. Section 3 gives the bilateral projective semantics and locates the affine sector. Section 4 restates the regular-AES interface imported from Paper 0. Section 5 develops the regular and singular zero geometry. Section 6 records what is proved, what remains open, and the interfaces to the later papers. The appendices collect convention checks and the small examples that separate the equality levels of (1).

2 Sequential Arithmetic Histories

Arithmetic evaluation is usually presented as a map from a syntactic expression to its value. This description suppresses the order in which intermediate values are formed. We begin by isolating the finite binary trees for which the internal operations have only one legal temporal order. Such a tree is not yet a history: one must also mark which leaf of its innermost operation is to be carried forward as the evolving value. The resulting object, a marked spinal history, is the syntactic unit used in the remainder of the paper.

Throughout this section, K is a field and

$$\Omega = \{+, -, \times, /\}$$

is the binary arithmetic signature. Division is interpreted as a partial operation in ordinary arithmetic. The combinatorial statements below apply to any binary signature; the choice of Ω is needed only when histories are evaluated.

2.1 Expression trees and dependency

Definition 2.1 (Arithmetic expression tree). An *arithmetic expression tree over K* is a finite rooted planar binary tree whose leaves are labelled by elements of K or by formal variables and whose internal vertices are labelled by elements of Ω . Its set of internal vertices is denoted by $I(T)$.

The planar structure records the first and second operand slots. It does not, by itself, specify a temporal order between operations on disjoint subtrees. That order is recorded by the dependency relation.

Definition 2.2 (Internal dependency poset). For $u, v \in I(T)$, write $u \preceq_T v$ when either $u = v$ or the value produced at u is required as an input at v . Equivalently, v lies on the path from u to the root. A *legal internal evaluation order* is a linear extension of the finite poset $(I(T), \preceq_T)$.

Leaves are not included in $I(T)$. Thus a tree with no internal vertex has the unique empty internal evaluation order. This convention separates the order of operations from the prior availability of atomic inputs.

We shall use the following elementary finite-poset fact in both directions.

Lemma 2.3 (Unique linear extensions). *A finite poset has a unique linear extension if and only if it is a chain.*

Proof. Every linear extension of a chain is its given total order, so a chain has a unique linear extension. Conversely, let P be a finite poset and suppose that $u, v \in P$ are incomparable. Adding the relation $u < v$ and taking its transitive closure cannot create a directed cycle: such a cycle would contain a pre-existing path $v \leq u$, contrary to incomparability. The resulting finite partial order has a linear extension. The same argument with $v < u$ gives a second linear extension. One orders u before v and the other orders v before u , so the extension of P is not unique. Hence a poset with a unique linear extension contains no incomparable pair and is a chain. $\square$

Theorem 2.4 (Sequential-tree classification). *For a finite binary expression tree T , the following conditions are equivalent.*

1. *The dependency poset $I(T)$ has a unique linear extension.*
2. *The dependency poset $I(T)$ is a chain.*
3. *Every internal vertex has at most one internal child.*
4. *The internal vertices form a single spine: either $I(T)$ is empty, or there is a path from the root to an innermost internal vertex containing every internal vertex of T .*

Proof. The equivalence of the first two conditions is Lemma 2.3.

Suppose that $I(T)$ is a chain. If an internal vertex had two internal children, neither child could depend on the other, since they lie in disjoint subtrees. They would therefore be incomparable in $I(T)$, a contradiction. Thus (2) implies (3).

Assume (3). If $I(T)$ is empty there is nothing to prove. Otherwise begin at the root. Whenever the current internal vertex has an internal child, that child is unique. Following these children produces a path ending at an internal vertex with two leaf children. Every internal vertex lies on this path: an internal vertex off the path would occur in the other child-subtree of some vertex on the path, giving that vertex two internal children. Hence the path contains all internal vertices, proving (4).

Finally, if all internal vertices lie on one root-to-leaf path, their ancestor relation is total. Read from the innermost vertex towards the root, each operation depends on its predecessor. Thus $I(T)$ is a chain, proving (4) $\Rightarrow$ (2) and completing the equivalence. $\square$

Figure 2 contrasts the two possible local patterns at the level of internal dependencies. It is only a visual summary of the order relation; the classification rests on the preceding proof.

Definition 2.5 (Sequential tree). *A finite binary expression tree satisfying the equivalent conditions of Theorem 2.4 is called a *sequential tree*.*

The adjective refers only to internal evaluation. It does not choose an active leaf, and it does not assert that ordinary arithmetic evaluation is defined. For example,

$$(a + b)(c + d)$$

is not sequential: the two additions are incomparable dependencies of the final multiplication. By contrast,

$$((a + b)c - d)/e$$

is sequential, because its internal vertices lie on one spine.




Figure 2: A sequential tree has a chain of internal dependencies (left). A branching tree contains incomparable internal operations and therefore admits more than one legal internal evaluation order (right).

2.2 One-hole contexts and the marked accumulator

Let z denote a hole rather than an arithmetic variable. For $\omega \in \Omega$ and $c \in K$, define the two elementary one-hole contexts

$$C_{\omega,c}^{(1)}[z] = \omega(z, c), \quad C_{\omega,c}^{(2)}[z] = \omega(c, z). \quad (2)$$

The superscript gives the operand slot occupied by the evolving value. It is therefore intrinsic to the marked process and should not be replaced by a description based on how a planar drawing happens to expand.

At the innermost internal vertex of a nontrivial sequential tree, both children are leaves. The dependency poset determines that this operation is performed first, but it does not determine which of its two inputs is the value to be carried along the spine. Choosing one of those leaves as the initial accumulator removes this ambiguity.

Definition 2.6 (Marked spinal history). A *free marked spinal history* over K is a finite chronological word

$$\gamma = (g_1, \dots, g_n), \quad g_i = C_{\omega_i, c_i}^{(\varepsilon_i)}, \quad \varepsilon_i \in \{1, 2\}.$$

The context g_1 is applied first and g_n last. Its *chirality word* is

$$\varepsilon(\gamma) = \varepsilon_1 \cdots \varepsilon_n.$$

A *bounded marked spinal history* is a pair $(x_0; \gamma)$ consisting of an initial accumulator x_0 and a free history. The empty word is allowed; its bounded form represents the marked single-leaf tree.

When no confusion is possible, we call either form a marked spinal history and state explicitly whether an initial value is present. Earlier drafts called certain pure-chirality cases threadlike expressions. The present definition also includes mixed chirality and makes the marking part of the data.

Figure 3 displays operand-slot chirality directly in the one-hole contexts. The diagram is a notation guide, not a substitute for the marked-tree correspondence proved below.

Proposition 2.7 (Marked-tree-history correspondence). *Let T be a sequential tree all of whose leaves are labelled by elements of K .*

1. *If T is a single leaf, marking that leaf determines the empty bounded history.*
2. *If T is nontrivial, a choice of one leaf at its innermost internal vertex as the initial accumulator determines a unique bounded marked spinal history.*




Figure 3: Pure slot-(1), pure slot-(2), and mixed marked histories. In each box, $\bullet$ is the current accumulator—the marked seed in the first box and the preceding output thereafter. Its displayed position is the operand slot, while time always runs along the arrows.

3. Conversely, a bounded marked spinal history determines a unique marked planar sequential tree.

These two constructions are mutually inverse when operation labels, constants, operand slots, and the initial mark are retained.

Proof. The single-leaf case is immediate. Suppose T is nontrivial. By Theorem 2.4, its internal vertices form a unique spine. At the innermost vertex, mark one leaf x_0 and denote the other by c_1 . The operation label and the slot containing x_0 determine $g_1 = C_{\omega_1, c_1}^{(\varepsilon_1)}$. Move one vertex towards the root. Its off-spine child must be a leaf, since otherwise it would be a second internal child. Its label c_2 , the operation label, and the slot occupied by the spine determine g_2 . Repeating this procedure reaches the root and produces a unique chronological word $(g_1, \dots, g_n)$.

Conversely, start with a marked leaf x_0 . Replace it by the planar binary tree $C_{\omega_1, c_1}^{(\varepsilon_1)}[x_0]$, then place the resulting tree in slot ε_2 of the next context, and continue. Exactly one child at every new internal vertex is internal, so the result is sequential. Each step records the operation, off-spine leaf, and slot used in the forward construction. The constructions therefore recover one another. $\square$

The proposition concerns fully labelled planar trees. If parentheses, constant labels, or the marked seed are forgotten, the correspondence is no longer injective.

2.3 Composition and ordinary evaluation

If

$$\gamma = (g_1, \dots, g_m), \quad \delta = (h_1, \dots, h_n),$$

their chronological concatenation is

$$\gamma\delta = (g_1, \dots, g_m, h_1, \dots, h_n), \tag{3}$$

so γ is performed first and δ second. This convention is fixed for the entire paper and is summarized again in Appendix A.

Each elementary context determines a partial function $g_i: K \dashrightarrow K$. Starting from $x \in K$, define recursively

$$x_0 = x, \quad x_i = g_i(x_{i-1}).$$

The bounded history is *ordinarily admissible at x* when every x_i is defined in ordinary field arithmetic. In that case its endpoint is

$$\nu_x(\gamma) = g_n \circ \dots \circ g_1(x). \tag{4}$$

The admissible domain of a free history is the set of such x . In particular, a second-slot division c/z excludes any state at which the evolving value is zero, while a first-slot division z/c requires $c \neq 0$.

The concatenation convention gives

$$\nu_x(\gamma\delta) = \nu_{\nu_x(\gamma)}(\delta) \quad (5)$$

whenever both sides are ordinarily admissible. Equation (5) is a statement about partial functions. Chapter 3 introduces a projective evaluation with a different domain convention.

2.4 Mirror, temporal reversal, and path inverse

Three operations on a context word must be kept separate.

Definition 2.8 (Mirror and temporal reversal). For $g = C_{\omega,c}^{(\varepsilon)}$, let

$$m(g) = C_{\omega,c}^{(3-\varepsilon)}.$$

The *mirror* and *temporal reversal* of $\gamma = (g_1, \dots, g_n)$ are respectively

$$m\gamma = (m(g_1), \dots, m(g_n)), \quad r\gamma = (g_n, \dots, g_1).$$

The mirror exchanges operand slots while retaining chronological order; temporal reversal reverses chronological order while retaining the contexts.

Both operations are involutions on marked words, and they commute: $m(r\gamma) = r(m\gamma)$. This is a syntactic statement; ordinary domains of the evaluated words need not agree.

Definition 2.9 (Path inverse). Suppose every context g_i in γ acts as an invertible function on a declared domain. The *path inverse* is the functional word

$$\gamma^{-1} = (g_n^{-1}, \dots, g_1^{-1}).$$

It is defined only under this invertibility hypothesis and need not have the same operation labels or ordinary admissible domain as $r\gamma$.

For example, let $g_1[z] = z + 1$ and $g_2[z] = 2z$. Then

$$\gamma = (g_1, g_2): x \mapsto 2x + 2.$$

Because addition and multiplication are symmetric operations, $m\gamma$ has the displayed expression $2(1 + x)$ and induces the same operator $2x + 2$. In contrast,

$$r\gamma = (g_2, g_1): x \mapsto 2x + 1,$$

while

$$\gamma^{-1} = (z \mapsto z/2, z \mapsto z - 1): x \mapsto x/2 - 1.$$

Thus equality with a mirror in this example is caused by the symmetry of the chosen elementary operations; it does not identify mirror, reversal, and inverse. The affine order defect studied later compares temporal orders. It is not automatically an invariant of planar mirror.

2.5 Information retained and information forgotten

Several equivalence relations occur naturally, and the remainder of the paper uses them at different stages. It is useful to order them by the information they forget.

1. *Literal syntactic equality* retains the fully parenthesized written expression.
2. *Marked-history equality* retains the initial mark, the chronological context word, and every operand-slot label. It may forget inessential typography but not process data.
3. *Operator equality* identifies histories inducing the same partial function on a stated ordinary domain, or the same projective transformation after projective evaluation.
4. *Charge equality*, when a charge map has been specified, retains only the chosen abelian totals of a history.
5. *Endpoint equality* at x asks only whether the admissible values $\nu_x(\gamma)$ and $\nu_x(\delta)$ coincide.
6. *Quotient or relational equality* is any further identification imposed by an explicitly declared rewriting or condensation relation.

This list is not a claim that every adjacent pair gives a strict quotient in all languages. It is a discipline for stating which level is being used. For example, the two histories

$$\gamma = (z \mapsto z + 1, z \mapsto 2z), \quad \delta = (z \mapsto 3z, z \mapsto z + 1)$$

are distinct marked histories and induce the distinct operators $2z + 2$ and $3z + 1$, but at the initial value $x = 1$ both have endpoint 4.

2.6 Representative chirality patterns

The slot notation exposes examples that the older comb terminology obscured.

Example 2.10 (Pure slot-(1) history). The word

$$C_{+,2}^{(1)}, \quad C_{\times,3}^{(1)}, \quad C_{-,4}^{(1)}$$

has chirality word 111 and evaluates ordinarily as $x \mapsto (x + 2)3 - 4$.

Example 2.11 (Pure slot-(2) history). The word

$$C_{-,2}^{(2)}, \quad C_{/,3}^{(2)}$$

has chirality word 22 and evaluates as

$$x \mapsto \frac{3}{2 - x},$$

on the ordinary domain $K \setminus \{2\}$. The accumulator occupies the second operand slot at both steps.

Example 2.12 (Mixed chirality). The word

$$C_{-,1}^{(1)}, \quad C_{/,2}^{(2)}, \quad C_{+,3}^{(1)}$$

has chirality word 121 and evaluates as

$$x \mapsto \frac{2}{x - 1} + 3,$$

on its ordinary domain. Its dependency poset is still a chain; mixed chirality does not introduce branching.

The syntactic passage is therefore

$$\begin{aligned} &\text{unmarked sequential tree} \xrightarrow{\text{choice of accumulator}} \text{marked planar sequential tree,} \\ &\text{marked planar sequential tree} \longleftrightarrow \text{bounded marked spinal history,} \\ &\text{bounded marked spinal history} \longrightarrow \text{operator} \longrightarrow \text{endpoint.} \end{aligned}$$

when the history is bounded and all planar labels are retained. The first arrow is a choice and the double arrow is a bijection; only the later evaluation arrows forget process data. Chapter 3 constructs the operator stage for all non-degenerate bilateral contexts.

3 Projective Semantics and the Affine Sector

The four arithmetic operations behave asymmetrically once the accumulator may occupy either operand slot. Most elementary contexts are affine, but the second-slot division c/z is genuinely projective. This section makes that enlargement precise. The projective semantics does not replace ordinary arithmetic evaluation: it supplies an operator-valued continuation on a different domain.

Throughout this section, K is an arbitrary field. Points of the projective line are one-dimensional subspaces of K^2 , written in homogeneous coordinates as

$$[u : v], \quad (u, v) \neq (0, 0).$$

We use the affine chart $z = [z : 1]$ and write $\infty = [1 : 0]$. Matrices act on column vectors from the left. Thus

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \implies M \cdot z = \frac{Az + B}{Cz + D}, \quad (6)$$

with the homogeneous interpretation at poles and at infinity. Replacing M by a nonzero scalar multiple does not change the projective map.

3.1 Ordinary domains and projective continuation

For an elementary context $g = C_{\omega, c}^{(\varepsilon)}$, ordinary evaluation gives a partial function $g: K \dashrightarrow K$. A *non-degenerate projective context* is one for which the corresponding homogeneous matrix is invertible. It acts everywhere on $\mathbb{P}^1(K)$, although its affine-coordinate formula may have a pole.

The elementary contexts and matrix representatives are as follows.

Context	Affine-coordinate map	Matrix representative
$C_{+,c}^{(1)} = C_{+,c}^{(2)}$	$z \mapsto z + c$	$\begin{pmatrix} 1 & c \\ 0 & 1 \end{pmatrix}$
$C_{-,c}^{(1)}$	$z \mapsto z - c$	$\begin{pmatrix} 1 & -c \\ 0 & 1 \end{pmatrix}$
$C_{-,c}^{(2)}$	$z \mapsto c - z$	$\begin{pmatrix} -1 & c \\ 0 & 1 \end{pmatrix}$
$C_{\times,c}^{(1)} = C_{\times,c}^{(2)}$	$z \mapsto cz$	$\begin{pmatrix} c & 0 \\ 0 & 1 \end{pmatrix}$
$C_{/,c}^{(1)}$	$z \mapsto z/c$	$\begin{pmatrix} 1 & 0 \\ 0 & c \end{pmatrix}$
$C_{/,c}^{(2)}$	$z \mapsto c/z$	$\begin{pmatrix} 0 & c \\ 1 & 0 \end{pmatrix}$

The first three rows are invertible for every $c \in K$. Multiplication by c , first-slot division by c , and second-slot division with numerator c are projectively invertible precisely when $c \neq 0$. At $c = 0$, multiplication and $0/z$ are constant on their ordinary domains, while $z/0$ has empty ordinary domain. Their singular matrix representatives do not act everywhere on $\mathbb{P}^1(K)$: each has a projective base point. They are therefore excluded from $\text{PGL}_2(K)$ and from the non-degenerate history language used below.

The table also displays the distinction between ordinary and projective domains. If $c \neq 0$, ordinary first-slot division z/c is defined on all of K . Ordinary second-slot division c/z is defined only for $z \neq 0$. Projectively, the latter sends 0 to ∞ and ∞ to 0. This is a valid action on $\mathbb{P}^1(K)$ but not an ordinary evaluation through a division by zero.

Warning 3.1 (Projective values are not ordinary values). Projective continuation records a fractional linear operator and its chart transitions. It does not certify that the original arithmetic word is ordinarily admissible at a given initial value. Whenever an endpoint in K is used as an ordinary arithmetic result, ordinary admissibility must be checked separately.

3.2 Projective evaluation of a chronological history

Let $\text{Hist}_K^{\text{nd}}$ denote the free marked spinal history language generated by the non-degenerate elementary contexts in the table. The superscript “nd” refers to projective non-degeneracy, not to ordinary admissibility at every affine input.

Definition 3.2 (Projective evaluation). Choose for each elementary non-degenerate context g_i a matrix representative $M_i \in \text{GL}_2(K)$. For the chronological word $\gamma = (g_1, \dots, g_n)$, with g_1 applied first, define

$$\rho(\gamma) = [M_n \cdots M_1] \in \text{PGL}_2(K). \quad (7)$$

For the empty word, set $\rho(\emptyset) = 1$.

This definition uses projective operator evaluation. For a bounded history $(x; \gamma)$, its projective endpoint is $\rho(\gamma) \cdot x$ in $\mathbb{P}^1(K)$. Its ordinary endpoint $\nu_x(\gamma)$ remains the partial evaluation defined in Section 2.

Proposition 3.3 (Compatibility with chronology). *Projective evaluation is independent of the chosen scalar matrix representatives. If $\gamma\delta$ means first γ and then δ , then*

$$\rho(\gamma\delta) = \rho(\delta)\rho(\gamma). \quad (8)$$

Moreover, whenever γ is ordinarily admissible at x and its ordinary endpoint lies in K , the projective endpoint agrees with it in the affine chart:

$$\rho(\gamma) \cdot x = [\nu_x(\gamma) : 1].$$

Proof. Replacing M_i by $a_i M_i$, with $a_i \in K^\times$, multiplies the product $M_n \cdots M_1$ by the nonzero scalar $\prod_i a_i$ and hence does not alter its projective class. If $\gamma = (g_1, \dots, g_m)$ and $\delta = (h_1, \dots, h_n)$, then the matrix word for $\gamma\delta$ is

$$N_n \cdots N_1 M_m \cdots M_1,$$

which represents $\rho(\delta)\rho(\gamma)$. This proves (8).

For the last assertion, each row of the table agrees with its ordinary arithmetic context wherever that context is defined. Applying this observation inductively along the admissible intermediate states proves equality in the affine chart. $\square$

Thus ρ is a homomorphism from chronological words if the target group is written with the opposite multiplication. We retain ordinary matrix multiplication and the explicit rule (8), because it makes both the column action and function composition standard.

3.3 Generation of the projective group

Introduce the elementary projective transformations

$$T_s(z) = z + s, \quad D_q(z) = qz \quad (q \in K^\times), \quad J(z) = -\frac{1}{z}.$$

They are projective evaluations of, respectively, addition by s , multiplication by q , and second-slot division with numerator -1 . These three kinds of context suffice for the entire bilateral projective language at the operator level.

Theorem 3.4 (Bilateral arithmetic generates PGL_2). *Let K be any field. The projective evaluations of non-degenerate bilateral four-operation marked spinal histories have image*

$$\rho(\text{Hist}_K^{\text{nd}}) = \text{PGL}_2(K).$$

Equivalently, translations T_s , nonzero scalings D_q , and J generate $\text{PGL}_2(K)$.

Proof. Each of T_s, D_q, J is available in the bilateral context language, so every composition of them is in the evaluation image. Conversely, let

$$f(z) = \frac{Az + B}{Cz + D}, \quad AD - BC \neq 0,$$

be an arbitrary element of $\text{PGL}_2(K)$.

Suppose first that $C = 0$. The determinant condition then gives $A \neq 0$ and $D \neq 0$. Hence

$$f(z) = \frac{A}{D}z + \frac{B}{D} = T_{B/D} \circ D_{A/D}(z).$$

This is the projective evaluation of the chronological two-step word that first applies $D_{A/D}$ and then $T_{B/D}$.

Now suppose that $C \neq 0$. All parameters in the following expression are defined, and $(AD - BC)/C^2$ is nonzero. Direct calculation gives

$$\begin{aligned} T_{A/C} \circ D_{(AD-BC)/C^2} \circ J \circ T_{D/C}(z) &= \frac{A}{C} - \frac{AD-BC}{C^2} \frac{1}{z + D/C} \\ &= \frac{A(Cz + D) - (AD - BC)}{C(Cz + D)} \\ &= \frac{ACz + BC}{C(Cz + D)} = \frac{Az + B}{Cz + D}. \end{aligned}$$

With the fixed chronological convention, the corresponding history word is

$$(T_{D/C}, J, D_{(AD-BC)/C^2}, T_{A/C}),$$

whose entries are read from left to right in time. Thus f is again an evaluation of bilateral contexts. The two cases exhaust $\text{PGL}_2(K)$.

No step uses an ordering or a characteristic assumption on K . In characteristic two the signs in J and in the determinant simplify, but the same calculation remains valid because $-1 = 1$ is still nonzero. $\square$

The proof is constructive: it gives a history of length at most four for any specified projective matrix, once arbitrary field constants are admitted as context parameters. It establishes surjectivity at the operator level, not uniqueness of a representing history. In fact, many distinct context words have the same projective evaluation.

Example 3.5 (A pole in an ordinarily inadmissible history). Let $K = \mathbb{Q}$ and consider the one-step history $g[z] = 1/z$. It is a non-degenerate projective context with $\rho(g) \cdot 0 = \infty$. The bounded history $(0; g)$ has no ordinary endpoint. Appending another inversion gives a projective word acting as the identity, but the two-step ordinary evaluation at 0 is still inadmissible because its first step is undefined.

3.4 A distinguished arithmetic sublanguage: the $q=4$ Hecke group

The surjectivity theorem allows arbitrary constants and produces the whole projective group. A much smaller constant alphabet already carries a useful discrete geometry. Work in $\mathbb{Q}(\sqrt{2}) \subset \mathbb{R}$, and let Hist_4 be the symmetric marked history language generated by the three elementary contexts

$$\mathbf{t}_+[z] = z + \sqrt{2}, \quad \mathbf{t}_-[z] = z - \sqrt{2}, \quad \mathbf{j}[z] = -\frac{1}{z}.$$

Here “symmetric” means that the inverse translation is retained as a generator; $\mathbf{j}$ is its own projective inverse. Write

$$T = T_{\sqrt{2}}, \quad G_4 = \rho(\text{Hist}_4) = \langle T, J \rangle \leq \text{PGL}_2(\mathbb{R}).$$

This is the standard projective realization of the $q = 4$ Hecke group [4, 2].

Proposition 3.6 (The order-four arithmetic relation). *With the column action and chronological convention fixed above, the generators have determinant-one representatives*

$$\tilde{T} = \begin{pmatrix} 1 & \sqrt{2} \\ 0 & 1 \end{pmatrix}, \quad \tilde{J} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \quad \text{in } \text{SL}_2(\mathbb{Z}[\sqrt{2}]).$$

Consequently G_4 is defined by matrices over $\mathbb{Z}[\sqrt{2}]$. In $\mathrm{PGL}_2(\mathbb{R})$ one has

$$J^2 = 1, \quad (JT)^4 = 1, \quad (9)$$

and JT has exact projective order four. The second relation is the projective evaluation of the chronological eight-step history

$$\omega_4 = (\mathfrak{t}_+, \mathfrak{j}, \mathfrak{t}_+, \mathfrak{j}, \mathfrak{t}_+, \mathfrak{j}, \mathfrak{t}_+, \mathfrak{j}). \quad (10)$$

Thus ω_4 and the empty history are distinct marked histories with the same projective operator.

Proof. Both displayed matrices have determinant one and entries in $\mathbb{Z}[\sqrt{2}]$; their inverses have the same property. Moreover,

$$\tilde{J}^2 = -I, \quad R := \tilde{J}\tilde{T} = \begin{pmatrix} 0 & -1 \\ 1 & \sqrt{2} \end{pmatrix}, \quad R^2 = \begin{pmatrix} -1 & -\sqrt{2} \\ \sqrt{2} & 1 \end{pmatrix}.$$

A second multiplication gives $R^4 = -I$. Hence the scalar classes satisfy (9). Neither R nor R^2 is scalar, so the projective order of JT is exactly four. Finally, each chronological pair $(\mathfrak{t}_+, \mathfrak{j})$ evaluates as $J \circ T$ and is represented by $\tilde{J}\tilde{T} = R$. Four such pairs therefore evaluate as R^4 , which is the identity in the projective group. $\square$

The proposition is an operator statement, not a universal statement about ordinary real arithmetic. Every letter of Hist_4 is projectively non-degenerate, but a bounded history is ordinarily admissible only if the evolving affine value is nonzero whenever $\mathfrak{j}$ is applied. For example, ω_4 is projectively neutral but is not ordinarily admissible at $x = -\sqrt{2}$: its first translation reaches zero, after which $\mathfrak{j}$ requests division by zero. In contrast, the same matrices act without an interior pole on the upper half-plane $\mathbb{H}^2$. The Fuchsian action and ordinary evaluation on a specified real input are therefore two related but distinct uses of the same context word.

Remark 3.7 (From histories to Hecke cells). The conventional action of G_4 on $\mathbb{H}^2$ supplies its Hecke tessellations and derived dual cellulations. Paper I needs only the following formal interface. If $\mathcal{C}$ is a chosen cell in such a G_4 -equivariant cellulation and $H_{\mathcal{C}} = \mathrm{Stab}_{G_4}(\mathcal{C})$, then the cell reached by a history factors as

$$\mathrm{Hist}_4 \xrightarrow{\rho} G_4 \longrightarrow G_4/H_{\mathcal{C}}, \quad \gamma \longmapsto \rho(\gamma)H_{\mathcal{C}}. \quad (11)$$

Neither arrow is generally injective. Relations such as (10) identify distinct histories at the operator level, and distinct operators in one coset act on $\mathcal{C}$ as the same cell. Passing instead to one endpoint uses the stabilizer of that point and forgets still more information. Thus arithmetic histories, projective operators, cells, and endpoints must not be placed in one-to-one correspondence. The order-four relation provides an arithmetic entry point for the fourfold incidence visible in the $q = 4$ geometry; the construction of analytic functions and singular zero sets on that cellulation is deferred to Papers II and III.

3.5 The affine syntactic sector

Let B_{∞} be the stabilizer of the chosen point at infinity:

$$B_{\infty} = \mathrm{Stab}_{\mathrm{PGL}_2(K)}(\infty).$$

An element represented by $\begin{pmatrix} A & B \\ C & D \end{pmatrix}$ fixes $\infty = [1 : 0]$ exactly when $C = 0$. After rescaling by D^{-1} , it then has a unique representative

$$\begin{pmatrix} a & b \\ 0 & 1 \end{pmatrix}, \quad a \in K^{\times}, \quad b \in K,$$




Figure 4: The affine/Borel sector inside the bilateral projective operator semantics. Elementary affine contexts generate the stabilizer B_∞ , whereas inversion reaches the complementary Bruhat cell. Words containing projective steps may nevertheless evaluate back into B_∞ .

acting as $z \mapsto az + b$.

Define $\text{Hist}_K^{\text{aff}}$ to be the syntactic sublanguage generated by those *elementary non-degenerate contexts* in the table that individually fix ∞ . Equivalently, it excludes the elementary second-slot divisions c/z and retains every non-degenerate elementary affine context. This definition is deliberately syntactic: it does not assert that $\text{Hist}_K^{\text{aff}}$ is the entire inverse image $\rho^{-1}(B_\infty)$. A word containing projective steps can leave the affine chart and later evaluate to an affine operator.

Corollary 3.8 (Affine/Borel sector). *For every field K ,*

$$\rho(\text{Hist}_K^{\text{aff}}) = B_\infty \cong \text{Aff}(1, K).$$

In particular, the affine syntactic sublanguage maps surjectively onto the Borel subgroup stabilizing ∞ .

Proof. Every generator of $\text{Hist}_K^{\text{aff}}$ fixes ∞ ; therefore every product of their projective matrices fixes ∞ , and the image is contained in B_∞ . Conversely, every element of B_∞ acts as $z \mapsto az + b$ with $a \neq 0$. It is

$$T_b \circ D_a,$$

the evaluation of a chronological word consisting of the non-degenerate affine contexts D_a followed by T_b . Hence the image is all of B_∞ . The displayed normal form also identifies its group law with that of $\text{Aff}(1, K)$. $\square$

Figure 4 records the resulting operator-level placement. It does not identify the affine syntactic language with the full inverse image of the inner region.

The corollary locates the existing add–multiply language without collapsing the larger syntax. Pure slot-(1) non-degenerate histories lie in this affine sector, but so do second-slot subtraction and multiplication contexts. The criterion is stabilization of ∞ , not the slot bit by itself.

3.6 The positive real component

Later differential-geometric sections use real multiplication in the form

$$z \mapsto e^M z, \quad M \in \mathbb{R}.$$

Its multiplier is positive. Together with all real translations, these maps form

$$\text{Aff}^+(1, \mathbb{R}) = \{z \mapsto az + b : a > 0\},$$

the identity component of the real affine group. They do not include affine reflections with negative multiplier. The algebraic statement of Corollary 3.8 concerns all nonzero multipliers in $K^\times$; the continuous theory is its positive real component.

3.7 Bruhat placement and the projective infinitesimal direction

The case division in the proof of Theorem 3.4 also gives the rank-one Bruhat decomposition

$$\mathrm{PGL}_2(K) = B_\infty \sqcup B_\infty J B_\infty.$$

Indeed, $C = 0$ is precisely the first cell, while the explicit decomposition for $C \neq 0$ lies in the second. This observation places bilateral arithmetic as a projective completion of the affine sector; no representation-theoretic conclusions beyond this decomposition are needed here.

There is also an infinitesimal counterpart over $\mathbb{R}$ (or $\mathbb{C}$). The projective vector fields

$$\partial_z, \quad z\partial_z, \quad z^2\partial_z$$

span the local projective Lie algebra. The first two generate affine flow. A conjugated translation detects the third direction:

$$J \circ T_s \circ J^{-1}(z) = \frac{z}{1 - sz} = z + sz^2 + O(s^2).$$

Consequently a general scalar projective flow has Riccati form

$$\dot{z} = \beta + \alpha z + \kappa z^2.$$

Paper I develops only the affine slice $\kappa = 0$. No projective metric, projective arithmetic expression space, or projective contact theory is asserted here.

4 The Regular AES Interface

This short section restates the one geometric definition on which the zero theorems of Section 5 depend. It is imported from Paper 0, where the definition, the frame equivalence, the models, the flow, the commutativization, and the contact structure are developed and proved; the statements below are repeated here only so that this paper can be read on its own, and they carry the same labels and hypotheses as their source.

Definition 4.1 (Regular arithmetic expression space, imported). A *regular arithmetic expression space*, or regular AES, is a tuple

$$\mathfrak{E} = (M, g, a; \mu, \lambda)$$

such that M is an oriented smooth surface, possibly with boundary, g is a smooth Riemannian metric, $a \in C^\infty(M, \mathbb{R})$, $\mu \in \mathbb{R} \setminus \{0\}$, $\lambda \in \mathbb{R}$, and

$$|\nabla a|_g^2 = \mu^2 + \lambda^2 a^2 \tag{12}$$

everywhere on M . The function a is called the *assignment function*.

The metric is part of the structure; (12) is an eikonal equation on a specified Riemannian surface, not a metric-free identity. Paper 0 defines the arithmetic motions of such a space and proves that they realize addition, subtraction, multiplication, and division by a constant as motions of M .

Proposition 4.2 (Canonical arithmetic frame, imported). *For an oriented Riemannian surface (M, g) , a smooth function a , and constants $\mu \neq 0$ and λ , the following are equivalent.*

1. *The eikonal identity (12) holds.*
2. *There is a unique positively oriented orthonormal frame (X_u, X_v) satisfying*

$$X_u a = \mu, \quad X_v a = \lambda a. \quad (13)$$

When these conditions hold, the frame is

$$X_u = \frac{\mu}{q} E_a - \frac{\lambda a}{q} J E_a, \quad X_v = \frac{\lambda a}{q} E_a + \frac{\mu}{q} J E_a, \quad E_a = \frac{\nabla a}{q}, \quad q = \sqrt{\mu^2 + \lambda^2 a^2}, \quad (14)$$

and it satisfies

$$[X_u, X_v]a = \mu\lambda. \quad (15)$$

The proof is given in Paper 0. We use only the following immediate consequences, which is why the statement is repeated rather than cited.

1. $\mu \neq 0$ makes q everywhere positive, so $|\nabla a|$ never vanishes. This is the source of the regularity of zero sets below.
2. The bracket identity (15) shows that, when $\lambda \neq 0$, the two arithmetic directions cannot be realized as a coordinate frame.

No other object of Paper 0 is repeated in this paper. In particular, the basic hyperbolic model $\mathfrak{E}_0$, the punctured model $\mathfrak{E}_1$, the continuous arithmetic motion, the affine cocycles, the accumulative commutative space, tearing, and the contact structure are cited where needed and never reproved here.

5 Zero Loci and Singular Expression Spaces

The eikonal equation imposes an immediate restriction on the zero set of a regular assignment: its gradient cannot vanish there. This separates regular zero geometry from isolated zeros, crossings, and the parameter-dependent singular phenomena reserved for later work.

5.1 Regular zero loci

For an assignment function a , write

$$Z(a) = a^{-1}(0).$$

Theorem 5.1 (Regular zero locus). *Let $\mathfrak{E} = (M, g, a; \mu, \lambda)$ be a regular AES.*

1. *If M has no boundary, then $Z(a)$ is a smooth embedded codimension-one submanifold of M .*
2. *If M has boundary, then $Z(a) \cap \text{int } M$ is a smooth embedded codimension-one submanifold of $\text{int } M$.*

In particular, for a surface the indicated zero locus is a disjoint union of smooth one-dimensional components. No assertion is made here about finiteness, properness, compactness, or behavior at ∂M .

Proof. At every $p \in Z(a)$, the defining eikonal identity gives

$$|\nabla a(p)|_g^2 = \mu^2.$$

Because $\mu \neq 0$, one has $da_p \neq 0$. Thus 0 is a regular value of a when M is boundaryless, and of $a|_{\text{int } M}$ for interior zeros when a boundary is present. The regular-value theorem gives the stated embedded submanifolds. $\square$

Corollary 5.2 (Local rigidity of regular zeros). *Within the interior of a regular two-dimensional AES, the zero locus has no isolated point, crossing, branch point, or endpoint. None of these configurations can occur as the germ of $Z(a)$ at an interior point.*

Proof. By Theorem 5.1, a neighborhood of every interior zero admits smooth coordinates in which $a^{-1}(0)$ is a coordinate line. Such a germ has none of the listed configurations. $\square$

This conclusion is local. Without additional global hypotheses, it does not classify the number or topology of the components. In the basic hyperbolic model, for example, $Z(a) = \{x = 0, y > 0\}$. Completeness and the absence of a boundary strengthen the conclusion substantially.

5.2 Complete regular splitting and a global no-go theorem

Theorem 5.3 (Complete regular AES splitting). *Let $\mathfrak{E} = (M, g, a; \mu, \lambda)$ be a regular AES for which M is connected, has no boundary, and is complete as a Riemannian manifold. Define*

$$\mathcal{R}(t) = \int_0^t \frac{ds}{\sqrt{\mu^2 + \lambda^2 s^2}}, \quad r = \mathcal{R} \circ a. \quad (16)$$

Then $|\nabla r|_g = 1$, the vector field $X = \nabla r$ has a global flow $\Phi : \mathbb{R} \times M \rightarrow M$, and

$$F : Z(a) \times \mathbb{R} \longrightarrow M, \quad F(q, t) = \Phi_t(q), \quad (17)$$

is a diffeomorphism. In particular, a is surjective, $Z(a)$ is nonempty and connected, and M is noncompact.

The conclusion is a smooth product decomposition. It does not assert that F is an isometry from a Riemannian product.

Proof. Put $q(a) = \sqrt{\mu^2 + \lambda^2 a^2}$. Since $\mu \neq 0$, this is everywhere positive. The chain rule and the regular-AES equation give

$$\nabla r = \frac{\nabla a}{q(a)}, \quad |\nabla r|_g^2 = \frac{|\nabla a|_g^2}{q(a)^2} = 1.$$

Thus $X = \nabla r$ is a smooth bounded vector field. To see that its flow is complete, let an integral curve be defined on a finite time interval. Its speed is one, so over every bounded time interval it remains in a closed metric ball. By Hopf–Rinow, closed bounded balls in the complete connected Riemannian manifold M are compact. The standard local existence theorem for smooth vector fields then extends the integral curve past every finite endpoint. Hence Φ_t exists for all $t \in \mathbb{R}$.

Along this flow,

$$\frac{d}{dt} r(\Phi_t(p)) = dr(X) = |\nabla r|_g^2 = 1, \quad r(\Phi_t(p)) = r(p) + t. \quad (18)$$

It follows that every orbit meets $r^{-1}(0) = a^{-1}(0)$ exactly once, namely at $\Phi_{-r(p)}(p)$. Therefore the inverse of (17) is

$$p \longmapsto (\Phi_{-r(p)}(p), r(p)).$$




Figure 5: Regular and singular zero geometry. In a regular AES with $\mu \neq 0$, the assignment crosses zero with nonvanishing gradient, so the interior zero germ is a smooth curve. The radial disc model instead has an isolated zero at its declared singular point, where the assignment is not C^1 and the regular structure has no local extension.

Both maps are smooth, so F is a diffeomorphism. Equation (18) also shows that $r(M) = \mathbb{R}$. The function $\mathcal{R} : \mathbb{R} \rightarrow \mathbb{R}$ is a strictly increasing diffeomorphism (linear when $\lambda = 0$, and an inverse-hyperbolic-sine rescaling when $\lambda \neq 0$); hence a is surjective. Finally, $Z(a) \times \mathbb{R} \cong M$ is connected only if $Z(a)$ is connected, and the displayed product cannot be compact. $\square$

Remark 5.4 (Why the global hypotheses matter). If a boundary is present, the unit vector field ∇r may reach it in finite time. If the metric is incomplete, an integral curve may escape the manifold in finite time. In either case the global flow argument fails, so the theorem must not be applied to domains with boundary or to incomplete regular models without a separate extension analysis.

Theorem 5.3 is a global obstruction, not merely a local regular-value statement. A connected complete boundaryless regular scalar AES cannot have several zero components, let alone a branching or four-valent zero network. Consequently, a Hecke cellulation used as a zero graph must enter through singular points, a failure of completeness or boundarylessness, or data richer than one regular scalar assignment. Paper I records this no-go boundary; the corresponding singular constructions belong to Paper III.

Figure 5 summarizes the categorical distinction. It illustrates the local conclusions already proved; it does not classify singular zero germs.

5.3 Singular arithmetic expression spaces

The preceding theorem makes a singular framework unavoidable if one wishes to admit isolated or branching zeros.

Definition 5.5 (Singular arithmetic expression space). A *singular arithmetic expression space* is a tuple

$$\mathfrak{E}_{\text{sing}} = (M, S, g, a; \mu, \lambda)$$

with the following data:

1. M is a nonempty oriented smooth surface, possibly with boundary, and $S \subset M$ is a closed nowhere-dense singular set;
2. g is a smooth Riemannian metric on $M \setminus S$;
3. $a : M \rightarrow \mathbb{R}$ is a specified function whose restriction to $M \setminus S$ is smooth;

4. $\mu \in \mathbb{R} \setminus \{0\}$, $\lambda \in \mathbb{R}$, and

$$(M \setminus S, g, a|_{M \setminus S}; \mu, \lambda)$$

is a regular AES.

5. Every $p \in S$ is locally essential: there is no neighborhood U of p on which $g|_{U \setminus S}$ and $a|_{U \setminus S}$ admit smooth extensions $\tilde{g}$ and $\tilde{a}$, with $\tilde{g}$ nondegenerate, such that $(U, \tilde{g}, \tilde{a}; \mu, \lambda)$ is a regular AES.

The metric or assignment may possess weaker extensions across S ; their precise regularity and local behavior must be checked in each example.

The set S need not consist entirely of zeros. Nowhere density keeps the regular locus nonempty and dense, while local essentiality prevents an arbitrary enlargement of S by points at which the regular structure extends. For a singular AES define

$$Z_{\text{reg}}(a) = Z(a) \cap (M \setminus S), \quad Z_{\text{sing}}(a) = Z(a) \cap S. \quad (19)$$

Every point of $Z_{\text{reg}}(a)$ is a regular zero by Theorem 5.1; a point of $Z_{\text{sing}}(a)$ is called singular because the regular structure is not asserted there, even when some of the underlying data happen to extend smoothly.

5.4 A verified isolated-zero singular model

Let

$$\mathbb{D} = \{(\xi, \eta) \in \mathbb{R}^2 : \xi^2 + \eta^2 < 1\}, \quad \rho = \sqrt{\xi^2 + \eta^2},$$

and fix $\mu, \lambda > 0$. Equip the disc with

$$g_{\mathbb{D}} = \frac{4}{\lambda^2(1 - \rho^2)^2} (d\xi^2 + d\eta^2) \quad (20)$$

and define

$$a_{\mathbb{D}}(\xi, \eta) = \begin{cases} \frac{2\mu}{\lambda} \frac{\rho}{1 - \rho^2}, & (\xi, \eta) \neq (0, 0), \\ 0, & (\xi, \eta) = (0, 0). \end{cases} \quad (21)$$

Proposition 5.6 (Classification of the isolated-zero disc). *The tuple*

$$(\mathbb{D}, \{0\}, g_{\mathbb{D}}, a_{\mathbb{D}}; \mu, \lambda)$$

is a singular AES with the following properties.

1. *The metric $g_{\mathbb{D}}$ extends smoothly and nondegenerately across the center.*
2. *The assignment $a_{\mathbb{D}}$ is smooth on $\mathbb{D} \setminus \{0\}$ but is not C^1 at the center.*
3. *The punctured disc is a regular AES:*

$$|\nabla a_{\mathbb{D}}|_{g_{\mathbb{D}}}^2 = \mu^2 + \lambda^2 a_{\mathbb{D}}^2.$$

4. *Its zero decomposition is*

$$Z_{\text{reg}}(a_{\mathbb{D}}) = \emptyset, \quad Z_{\text{sing}}(a_{\mathbb{D}}) = \{0\}.$$

Proof. Formula (20) is written in Cartesian coordinates. Its conformal factor is a positive smooth function on all of $\mathbb{D}$, proving the first claim.

Away from the origin, ρ and hence $a_{\mathbb{D}}$ are smooth. On the ξ -axis, however,

$$\frac{a_{\mathbb{D}}(t, 0) - a_{\mathbb{D}}(0, 0)}{t} = \frac{2\mu}{\lambda} \frac{|t|}{t(1 - t^2)}.$$

The one-sided limits are $2\mu/\lambda$ and $-2\mu/\lambda$. Thus even the partial derivative in the ξ -direction does not exist at the center, so the assignment is not C^1 there. Moreover, the punctured assignment tends to zero at the center, so its continuous extension is unique. The same calculation therefore rules out every smooth local extension agreeing with the punctured assignment and proves that the declared singular point is locally essential.

On the punctured disc the assignment is radial, and

$$\frac{da_{\mathbb{D}}}{d\rho} = \frac{2\mu}{\lambda} \frac{1 + \rho^2}{(1 - \rho^2)^2}.$$

The inverse radial metric coefficient is

$$g_{\mathbb{D}}^{\rho\rho} = \frac{\lambda^2(1 - \rho^2)^2}{4}.$$

Consequently,

$$\begin{aligned} |\nabla a_{\mathbb{D}}|_{g_{\mathbb{D}}}^2 &= g_{\mathbb{D}}^{\rho\rho} \left(\frac{da_{\mathbb{D}}}{d\rho} \right)^2 \\ &= \mu^2 \frac{(1 + \rho^2)^2}{(1 - \rho^2)^2} \\ &= \mu^2 + \lambda^2 \left(\frac{2\mu}{\lambda} \frac{\rho}{1 - \rho^2} \right)^2, \end{aligned}$$

which is the regular-AES equation on $\mathbb{D} \setminus \{0\}$. Finally, $a_{\mathbb{D}} > 0$ for $0 < \rho < 1$, while its declared value at the center is zero. The claimed zero decomposition follows. $\square$

If $s = (2/\lambda) \operatorname{arctanh} \rho$ denotes hyperbolic distance from the center, then

$$a_{\mathbb{D}} = \frac{\mu}{\lambda} \sinh(\lambda s).$$

This identity describes radial propagation on the punctured regular part; it does not repair the failure of differentiability at the center. The metric is regular there, while the assignment is singular.

5.5 Smooth parameter families and their total zero sets

Let $I \subset \mathbb{R}$ be an open interval, let M be a smooth m -manifold without boundary, and let

$$A : M \times I \longrightarrow \mathbb{R}$$

be smooth. Write $a_t(p) = A(p, t)$ and

$$\mathcal{Z} = A^{-1}(0) = \{(p, t) \in M \times I : a_t(p) = 0\}.$$

Proposition 5.7 (Regular total zero set). *Assume that*

$$d_p a_t \neq 0 \quad \text{whenever } a_t(p) = 0. \quad (22)$$

Then $\mathcal{Z}$ is a smooth embedded codimension-one submanifold of $M \times I$, and the restriction

$$\pi : \mathcal{Z} \longrightarrow I, \quad \pi(p, t) = t, \quad (23)$$

is a submersion. In particular, if $\dim M = 2$, then $\mathcal{Z}$ is a smooth surface.

Proof. At a point $(p, t) \in \mathcal{Z}$, the total differential is

$$dA_{(p,t)}(v, \tau) = d_p a_t(v) + d_t A_{(p,t)}(\tau).$$

Hypothesis (22) implies that $dA_{(p,t)} \neq 0$, so 0 is a regular value of A . The regular-value theorem proves that $\mathcal{Z}$ is a codimension-one embedded submanifold.

It remains to prove the stronger assertion about the projection. Given an arbitrary $\tau \in T_t I$, put $c = d_t A_{(p,t)}(\tau)$, and choose $w \in T_p M$ with $d_p a_t(w) = 1$, which is possible because $d_p a_t \neq 0$. Set

$$v = -c w.$$

Then $dA_{(p,t)}(v, \tau) = 0$, so $(v, \tau) \in T_{(p,t)} \mathcal{Z}$, while $d\pi_{(p,t)}(v, \tau) = \tau$. Hence $d\pi$ is surjective at every point of $\mathcal{Z}$, proving that π is a submersion. $\square$

Remark 5.8 (Properness warning). Proposition 5.7 gives pointwise product charts for the projection, but it does not make $\pi : \mathcal{Z} \rightarrow I$ a globally trivial family, or even a fiber bundle over an entire parameter neighborhood when noncompact fibers are present. A standard sufficient hypothesis for local triviality as a smooth fiber bundle is that the projection onto the relevant parameter interval be a proper surjective submersion (with the usual additional care if boundaries are introduced). Without properness, components may escape to or arrive from infinity. Accordingly, $\mathcal{Z}$ is called the *total zero set* here, not a zero tube.

No multi-zero model is included in this foundational chapter. Constructing and classifying such models requires a complete audit of the domain, metric, assignment, singular set, flow equation, and exact zero topology; that problem belongs to the singular-zero theory developed after Paper I.

6 Conclusions and Research Interfaces

This paper has established the syntactic and projective foundation of the programme. The organizing principle is that an expression should not be identified prematurely with its endpoint value. An unmarked sequential tree requires a choice of accumulator; the resulting fully marked planar tree is equivalent to a bounded marked history. From that history, operator evaluation and the charge shadow form separate branches, while an initial value turns an operator into an endpoint. The non-bijective forgetful maps of (1) lose information, and the structures proved here are the ones that survive at the syntactic and operator levels.

6.1 Results proved here

The first part of the paper classified binary expression trees with a unique legal internal evaluation order and represented them by marked spinal histories. Operand-slot data made it possible to

distinguish planar mirror, temporal reversal, and path inverse. These operations act at different levels and need not induce the same operator comparison.

For nondegenerate bilateral one-hole contexts, projective continuation produced Möbius transformations. The elementary contexts generate $\mathrm{PGL}_2(K)$ over an arbitrary field, while ordinary addition and nonzero scaling form the affine, or Borel, sector stabilizing the chosen point at infinity. This placement is important: projective semantics extends the operator domain, but it does not turn a pole into an admissible ordinary arithmetic operation.

The smaller alphabet consisting of translation by $\pm\sqrt{2}$ and negative inversion realizes the $q = 4$ Hecke group inside this projective semantics. Its matrix relation $(JT)^4 = 1$ gives an explicit pair of different marked histories with the same operator. The subsequent factorization through cell stabilizers separates four levels that are needed downstream: history, operator, cell coset, and endpoint. In particular, arithmetic words are not identified one-to-one with tiles in a chosen Hecke cellulation.

The second part of the paper developed the zero theory of regular arithmetic expression spaces, using only the imported interface of Section 4. At a zero of a regular assignment one has $|\nabla a| = |\mu| \neq 0$, so the interior zero locus is a smooth codimension-one set and cannot have isolated points, crossings, or branch points. Under the additional assumptions that the metric is complete and the manifold is connected and boundaryless, unit-gradient rectification gives the diffeomorphism

$$M \cong Z(a) \times \mathbb{R},$$

hence the zero locus is nonempty and connected. This is not asserted for incomplete models or manifolds with boundary, and it is not in general a Riemannian product decomposition. The splitting theorem is therefore a no-go statement for branching zero networks in the regular category, and the paper accordingly defines singular arithmetic expression spaces by a closed nowhere-dense singular set admitting no local regular extension. One isolated-zero model is verified in detail; a smooth parameter family with spatially regular zeros has a submersive total zero set, and the properness warning records exactly how far that statement may be used.

The proved logical spine may therefore be summarized as

$$\begin{aligned} \text{sequential tree} &\longrightarrow \text{marked planar sequential tree} \longleftrightarrow \text{bounded marked history,} \\ \text{marked history} &\longrightarrow \text{projective operator} \longrightarrow \text{ordinary endpoint or pole,} \\ &\quad \mathrm{PGL}_2 \supset \mathrm{Aff}(1) \longrightarrow \text{Bruhat placement,} \\ \text{Hist}_4 &\longrightarrow G_4 \longrightarrow G_4/H_{\mathcal{C}} \longrightarrow \text{Hecke cell,} \\ \text{regular AES} &\longrightarrow Z(a) \text{ smooth} \longrightarrow M \cong Z(a) \times \mathbb{R} \text{ when complete,} \\ &\quad \text{singular set} \longrightarrow Z_{\mathrm{reg}}(a), Z_{\mathrm{sing}}(a). \end{aligned}$$

This is a foundation for later developments, not a claim that every arithmetic expression, singularity, or computational phenomenon is already captured by these levels.

6.2 What moved to Paper 0

The geometric development that earlier versions of this paper contained is now proved in Paper 0: the arithmetic motions of an arithmetic expression space, the basic hyperbolic model $\mathfrak{E}_0$ and the punctured model $\mathfrak{E}_1$, the continuous arithmetic motion and its rectification, the point-path and ripple-pencil readings of a chronological matrix product, the affine cocycles and the relative affine defect, the accumulative commutative space with its weighted torsion–Stokes theorem, the reading of torsion as tearing, and the contact structure with curvature $\mu\lambda$. Those results keep their labels,

hypotheses, and statuses; this paper cites them and does not maintain a second version. The migration is recorded in `governance/00b-paper-0-geometric-foundation-amendment.md`.

6.3 Interface to Paper II

Arithmetic Expression Geometry II: Hyperbolic Real Function Theory begins where the horizontal connection of Paper 0 stops. It develops additional metric and analytic data: a horizontal metric, compatible complex or almost-complex structures where appropriate, second-order operators, boundary-value constructions, and arithmetic holomorphicity. None of these structures follows uniquely from the contact form alone, and no result of that downstream function theory is used in the proofs of this paper [5]. The G_4 sublanguage supplies an additional, explicitly arithmetic test case: an automorphic or equivariant function used there must be derived from the group action and must keep its analytic descent conditions separate from ordinary arithmetic admissibility of a history at a real input. The geometry imported by Paper II — the regular-AES definition, the canonical frame, the basic model, and the contact model — is imported from Paper 0.

6.4 Interface to Paper III

Arithmetic Expression Geometry III: Singular Zero Geometry and Tubes studies the boundary of the regular-zero theorem. Its objects include singular and multiple zeros, parameter discriminants, regular and singular zero surfaces, and tube constructions. Braid, monodromy, and knot structures there require separate invariance and normalization arguments. The Hecke interface proved here offers a source-driven alternative to imposing a four-valent cellulation by hand: Paper III uses a normalized Hauptmodul to construct the corresponding singular zero dessin, then relates its sign cover to Hecke periodic-orbit knots [6]. Those analytic, singular, and three-dimensional results remain downstream results, not results of this paper. The complete regular splitting theorem shows that such a branching network cannot be the zero set of one connected, complete, boundaryless regular scalar AES; any singularity, boundary, or incompleteness used in the construction must therefore be stated explicitly. In particular, the existence of a tube does not by itself produce a knot invariant, and no such general invariant is asserted here.

6.5 Interface to Paper IV

Arithmetic Expression Geometry IV: Projective Condensation and Computational Complexity returns to the full projective process–result hierarchy. Its scope includes bivaluation, quotient and condensation constructions, representation complexity, and the relation between geometric and computational resources. The marked-history formalism and the affine embedding in projective semantics proved here supply an entry point, and the affine torsion that Paper IV compares against is the one proved in Paper 0. Paper IV proves model-relative quotient and residual results together with fixed-model resource calibrations, while general machine-robust complexity claims remain open and require explicit models of representation, input size, and computation.

6.6 Open foundational problems

Several questions remain at the level of the history layer.

1. Characterize the marked-history fibers of the downstream G_4 operator–zero-network realization and determine what arithmetic decoration, if any, survives its operator and knot quotients.

2. Produce fully verified singular multi-zero models and identify their stable local normal forms and discriminants; the single verified isolated-zero model is a starting point, not a classification.
3. Determine when the total zero set of a parameter family is proper — hence a locally trivial tube — rather than merely a submersive family, and state the minimal hypotheses under which that upgrade is possible.
4. Formulate, and then test, a quantitative relation between history complexity, representation complexity, and computational complexity, with an explicit state space, encoding, and cost model.
5. Determine which relations among marked histories are compatible with ordinary admissibility, as opposed to projective non-degeneracy alone; the Hecke order-four example shows that the two conditions are genuinely different.

These are research interfaces rather than consequences of the present paper. The stable conclusion is narrower and more concrete: sequential arithmetic histories possess projective operator semantics with a distinguished arithmetic sublanguage; their affine sector is the Borel sector of that semantics; and the regular zero geometry of an arithmetic expression space is rigid in exactly the sense stated above, so that singular and branching phenomena require an explicit change of category.

A Composition, action, and domain conventions

This appendix collects the convention-sensitive choices used in Sections 2–3. The choices are not mathematical assertions, but every composition, sign, and coordinate formula depends on them. The affine-group conventions that belong to the cocycle and flow chapters of Paper 0 are collected in that paper’s appendix.

A.1 Chronological words and functional composition

Convention A.1 (Order inside a history). A history

$$\gamma = (g_1, \dots, g_n)$$

performs g_1 first and g_n last. Its functional evaluation is therefore

$$\text{ev}(\gamma) = g_n \circ \dots \circ g_1$$

when the g_i are viewed as maps.

Convention A.2 (Chronological concatenation). The written product $\gamma\delta$ means first perform γ and then perform δ . Thus

$$\text{ev}(\gamma\delta) = \text{ev}(\delta) \circ \text{ev}(\gamma).$$

The notation is chronological rather than functional: the earlier process is written on the left.

For example, if $g(z) = z + \mu$ and $h(z) = qz$, then the word $\gamma = (g, h)$ and the chronological product of the one-step histories gh both act as

$$h \circ g(z) = qz + q\mu.$$

The reverse chronological word hg acts as $g \circ h(z) = qz + \mu$. This two-step check fixes the sign of their endpoint difference:

$$\text{ev}(gh)(z) - \text{ev}(hg)(z) = \mu(q - 1).$$

A.2 Column-vector projective action

Convention A.3 (Matrix action). Homogeneous coordinates are columns, and matrices act from the left:

$$\begin{pmatrix} A & B \\ C & D \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} Au + Bv \\ Cu + Dv \end{pmatrix}.$$

On the affine chart $z = [z : 1]$, this gives

$$M \cdot z = \frac{Az + B}{Cz + D}.$$

If M_i represents g_i , the history $(g_1, \dots, g_n)$ is represented by

$$M_n \cdots M_1.$$

Consequently, for chronological concatenation,

$$\rho(\gamma\delta) = \rho(\delta)\rho(\gamma). \quad (24)$$

This is the matrix version of Convention A.2. A row-vector right action would require a different product convention; it is not used in this paper.

Projective matrices are scalar classes. If M and aM differ by $a \in K^\times$, they represent the same element of $\text{PGL}_2(K)$. Matrix identities used for projective maps may therefore be checked either exactly in $\text{GL}_2(K)$ or up to one overall nonzero scalar, provided the choice is stated.

A.3 Ordinary and projective domains

The following table records the domain distinction for the contexts involving division. Here $c \in K$ and z is the evolving affine value.

Context	Ordinary rule	Ordinary condition	Projective status
$C_{/,c}^{(1)}[z] = z/c$	divide by fixed c	$c \neq 0$	invertible if $c \neq 0$
$C_{/,c}^{(2)}[z] = c/z$	divide by evolving z	$z \neq 0$	invertible if $c \neq 0$
$C_{\times,c}^{(\varepsilon)}[z] = cz$	multiply by c	none	invertible if $c \neq 0$

Two different exclusions appear in the table. The parameter condition $c \neq 0$ ensures that the context has an invertible projective matrix. The state condition $z \neq 0$ ensures that an ordinary second-slot division is defined at that step. Projectively, a non-degenerate c/z sends 0 to ∞ ; this is a chart transition, not an ordinary quotient in K .

For a composite history, ordinary admissibility is checked at every intermediate state. It is not enough that the final projective point lies in the affine chart. In particular, two projective inversions compose to the identity operator, but the corresponding ordinary two-step word is not admissible at zero.

A.4 Mirror, reversal, and inverse notation

For $\gamma = (g_1, \dots, g_n)$, the three operations are summarized by

Operation	Resulting word	Required hypothesis
Mirror $m\gamma$	$(m(g_1), \dots, m(g_n))$	none
Temporal reversal $r\gamma$	$(g_n, \dots, g_1)$	none
Path inverse γ^{-1}	$(g_n^{-1}, \dots, g_1^{-1})$	every g_i invertible

Mirror exchanges operand slots and retains time. Temporal reversal changes time and retains the original contexts. Path inverse reverses time and inverts every operator. Accordingly,

$$\rho(\gamma^{-1}) = \rho(\gamma)^{-1},$$

whereas no analogous identity holds for $r\gamma$ in general.

A.5 Notation conversion summary

The table below lists older or potentially ambiguous descriptions and the notation used in Paper I.

Potentially ambiguous description	Paper I notation
evolving value in operand slot one	$C_{\omega,c}^{(1)}[z] = \omega(z, c)$
evolving value in operand slot two	$C_{\omega,c}^{(2)}[z] = \omega(c, z)$
chronological context word	$\gamma = (g_1, \dots, g_n)$, g_1 first
history product	$\gamma\delta$, first γ , then δ
operator evaluation	$\rho(\gamma)$
ordinary endpoint at x	$\nu_x(\gamma)$
accumulated scale	Φ_γ
target-frame translation	ξ_γ
source-normalized translation	$\hat{\xi}_\gamma = \Phi_\gamma^{-1}\xi_\gamma$
operand-slot exchange	$m\gamma$
time-order reversal	$r\gamma$
inverse operator path	γ^{-1}

B Equality and elementary neutrality examples

The examples below separate several equality tests without developing a general relation or loop theory. In each case, histories are marked chronological words, operators are affine maps, and endpoint values are obtained only after choosing an input.

B.1 Same value, different operators

Let

$$\gamma = (A_1), \quad \delta = (M_{\log 2}).$$

At the input $x = 1$, both histories have endpoint value 2. Their induced operators are nevertheless

$$\nu_x(\gamma) = x + 1, \quad \nu_x(\delta) = 2x,$$

which are unequal. Equality at one input therefore does not imply operator equality.

B.2 Same operator, different histories

The marked histories

$$\gamma = (A_1, A_2), \quad \delta = (A_2, A_1)$$

are different chronological words, but both induce $x \mapsto x + 3$. They also have the same total charge. Thus operator equality does not recover a marked history, even when charge data are retained.

B.3 Same charge, nonzero relative torsion

For $p, q \neq 0$, the histories

$$\gamma = (A_p, M_q), \quad \delta = (M_q, A_p)$$

have the same ACS endpoint (p, q) . Their operators differ by the translation

$$\tau(\gamma, \delta) = p(e^q - 1).$$

The ACS endpoint is therefore insensitive to order, whereas evaluation is not. The weighted filling, rather than the terminal charge, detects the difference.

B.4 Zero endpoint defect, nontrivial history

For $p \neq 0$, the nonempty history

$$\ell = (A_p, A_{-p})$$

induces the identity operator and hence has zero endpoint defect for every input. It is nevertheless not the empty marked history. In particular, operator neutrality does not imply literal or historical equality.

A slightly different phenomenon occurs for

$$(A_p, M_q, A_{-p}, M_{-q}).$$

This history has zero total ACS charge but induces translation by $p(1 - e^{-q})$. Thus charge neutrality does not imply operator neutrality.

B.5 Open torsion and closed-loop drift

For additive side $p = \mu h$ and logarithmic multiplicative side $q = \lambda k$, the two open orders differ by

$$T_{\text{open}} = p(e^q - 1),$$

whereas the chronological closed loop has vertical drift

$$T_{\text{closed}} = p(1 - e^{-q}) = e^{-q} T_{\text{open}}.$$

They have the common first-order density $\mu\lambda$, but they are distinct finite quantities. Consequently local order torsion, raw closed-loop holonomy, and infinitesimal curvature should not be used as interchangeable notions without the normalization or limiting operation that relates them.

These examples exhibit the hierarchy needed in the main text:

$$\text{marked history} \longrightarrow \text{operator} \longrightarrow \text{endpoint value},$$

with total charge providing an additional, order-forgetting projection. No converse implication between adjacent levels holds in general.

References

- [1] Noam Chomsky. Three models for the description of language. *IRE Transactions on Information Theory*, 2(3):113–124, 1956.
- [2] Cheng Lien Lang and Mong Lung Lang. Arithmetic and geometry of the Hecke groups. *Journal of Algebra*, 460:392–417, 2016.
- [3] Emil L. Post. Formal reductions of the general combinatorial decision problem. *American Journal of Mathematics*, 65(2):197–215, 1943.
- [4] David Rosen. A class of continued fractions associated with certain properly discontinuous groups. *Duke Mathematical Journal*, 21(3):549–563, 1954.
- [5] Mingli Yuan. Arithmetic expression geometry ii: Hyperbolic real function theory— horizontal operators, boundary problems, and arithmetic holomorphicity. Mathematical-review manuscript, 2026.
- [6] Mingli Yuan. Arithmetic expression geometry iii: Singular zero geometry and tubes— branched pullbacks, arithmetic zero networks, and topological transport. Mathematical-review manuscript, 2026.