

Data for Heroin Model

Tricia Phillips

January 11, 2019

- We will run our model from 2013-2017, and initial conditions for the classes will be estimated, as we do not have data for the number of individuals in each class at the beginning of 2013.
- The following are the **number of individuals in each class** for Tennessee in 2015:

***Susceptibles**: (Total population in 2015-size of four other classes= 6,590,726 – 1,819,581 – 48,000 – 14,000 – R_0 =FILL IN) [19]

***Prescription opioid users**, 2015: 1,819,581 [16]

Although this number does not explicitly state it is for individuals 12 and older, we assume it is since it comes from the Tennessee Department of Health; if it does include individuals under 12, we assume that number is negligible.

***Opioid addicts**, 2015/2016 average for “Pain Reliever Use Disorder” for individuals 12 and older: 48,000 [18]

We note that their definition of pain reliever use disorder includes those who meet the American Psychiatric Association criteria for dependence or abuse. Here, opioid dependence is classified as having “signs and symptoms that reflect compulsive, prolonged self-administration of opioid substances that are used for no legitimate medical purpose or...are used in doses that are greatly in excess of the amount needed for pain relief...regular patterns of compulsive drug use that daily activities are typically planned around obtaining and administering opioids.” This definition falls under our definition of opioid addiction. Opioid abuse, on the other hand, they consider to be less severe than dependence, and would not lead to the development of withdrawal symptoms. This latter definition does not fall under our characterization of addiction, but we make a note of this to say that this estimate for those with a pain reliever use disorder may be overestimated for what we are concerned with, but is an acceptable approximation [1].

***Heroin/fentanyl addicts**, 2015/2016 average for “Past Year Heroin Use” for individuals 12 and older: 14,000 [18]

Although this number includes those who may have used heroin once or twice in the past year, we are under the assumption that the majority of these individuals are addicts and that very few, if any, individuals use heroin recreationally. In addition, the number of heroin users does not include fentanyl users explicitly, but we are under the assumption that those who take fentanyl are a subset of those who use heroin, and therefore, would mostly be included in these numbers. We admit the values may be slightly too low, for the cases of individuals who do fentanyl and not

heroin, but data has not been found for fentanyl addicts only. Therefore, we are working under the assumption that it would be a negligible population that takes fentanyl without heroin. Overall, these two assumptions may work to balance one another out.

**Recovered addicts: won't be able to find because we do not know the total number of individuals that have been in treatment, successfully finished, and not relapsed in our time frame*

- For Tennessee, the **total number of individuals taking prescription opioids** for pain [16]:
2013: 1,845,144
2014: 1,824,342
2015: 1,819,581
2016: 1,761,363
2017: 1,636,374

Although this number does not explicitly state it is for individuals 12 and older, we assume it is since it comes from the Tennessee Department of Health; if it does include individuals under 12, we assume that number is negligible.

- For Tennessee individuals 12 and older, **the number of treatment admissions for non-heroin opiates/synthetics** as the primary substance of abuse to facilities that receive state/public funding (generally referring to funding by the state substance abuse agency) [11]:
2005: 1,578
2006: 1,529
2007: 1,743
2008: 2,022
2009: 2,464
2010: 3,384
2011: 3,884
2012: 4,203
2013: 4,485
2014: 4,530
2015: 4,326

We are under the assumption that if one were addicted to heroin in addition to prescription opioids, their heroin problem would be the primary reason for going to treatment and would be included in the following numbers.

- For Tennessee individuals 12 and older, **the number of treatment admissions for heroin** as the primary substance of abuse to facilities that receive state/public funding (generally referring to funding by the state substance abuse agency) [11]:
2010: 199
2011: 240
2012: 390

2013: 555
2014: 743
2015: 1,083

Again, these numbers do not include fentanyl users explicitly, but we are under the assumption that those who take fentanyl are a subset of those who use heroin, and therefore, would mostly be included in these numbers. We admit the values may be slightly too low, for the cases of individuals who do go to treatment with the primary substance of abuse being fentanyl, but there is not data available for those numbers currently.

- We use data on the number of prescription opioid overdose deaths which include natural, semi-synthetic, and synthetic opioids; however, we subtract out the number of fentanyl overdoses (fentanyl is classified as a synthetic prescription opioid), since those overdoses are counted for in their own category, listed below. This results in the following **total number of prescription opioid overdose deaths** [17]:

2013: (637-53=) 584
2014: (697-69=) 628
2015: (848-169=) 679
2016: (1,009-294=) 715

Although this number does not explicitly state it is for individuals 12 and older, we assume it is since it comes from the Tennessee Department of Health; if it does include individuals under 12, we assume that number is negligible.

- We add together the heroin and fentanyl overdoses from the years 2013-2016 for the state of Tennessee. The **total number of heroin and fentanyl overdoses** for these four years are:

2013: (63+53=) 116
2014: (147+69=) 216
2015: (205+169=) 374
2016: (260+294=) 554 [17].

Although this number does not explicitly state it is for individuals 12 and older, we assume it is since it comes from the Tennessee Department of Health; if it does include individuals under 12, we assume that number is negligible.

We may calculate the overdose death rate in the year 2015, since we know the number of addicted individuals in that year. To find the continuous-time rate at which individuals are dying from the addicted class, we consider the equation $kA_0 = A_0 e^{-\mu A t}$, where A_0 is the number of individuals addicted to opioids in 2015 and k is the proportion of these individuals in the addicted class at the start of 2016 (when $t = 1$). In 2015, there were 679 individuals out of the entire Tennessee population 12 and older that overdosed on prescription opioids [17]. However, it is estimated that only 54.6% of these individuals were actually at an increased risk for an opioid-related overdose death; we will assume that if an individual met

the criteria for at least one high-risk factor, that they were considered addicted to opioids [2]. Therefore, we will assume 54.6% of the 679 individuals that overdosed were addicted. With a total of 48,000 opioid addicts in 2015, this means that $(48,000 - 0.546 \cdot 679) / 48,000 \approx 0.992$ is the proportion of addicted individuals that remain by the beginning of the next year. This implies $0.992A_0 = A_0e^{-\mu_A(1)}$, and solving results in $\mu_A \approx 0.00775$.

Similarly, we may calculate μ_H . There were 14,000 heroin/fentanyl addicts in 2015 and 374 heroin-related overdoses. We make the assumption that if an individual died of a heroin overdose they were addicted in line with our previous assumptions that if one is using heroin, they are considered addicted due to the nature of the drug. This means that $(14,000 - 374) / 14,000 \approx 0.973$ is the proportion of heroin users that remain at $t = 1$, which implies $0.973H_0 = H_0e^{-\mu_H(1)}$, which results in $\mu_H \approx 0.0271$, the continuous-time rate at which individuals are dying from the heroin class.

We make a note that individuals that do not have an opioid use disorder and die because of an opioid overdose are counted in the “natural mortality rate.”

- Total population estimated in Tennessee each year [19]:
 2013: 6,490,795
 2014: 6,540,007
 2015: 6,590,726
 2016: 6,649,404
 2017: 6,715,984
 2018: **FILL IN**

There were an estimated 1,073,214 individuals Tennessee in 2018 aged 12 and under. To figure out those who are 12 years old, we take approximately 1/8th of the individuals that are in the age group 5-12, which is approximately 83,175 individuals [3]. Thus, an estimated 990,039 individuals are *under* the age of 12 in Tennessee. Given the last total population estimate for 2017 being 6,715,984 from above, this means that approximately 15% of the population is under the age of 12. **FIX when get updated 2018 total population.** Since we do not see a reason for this percentage to be significantly different from year to year, we assume that this percentage is constant throughout the time period we are looking at. Then, we are able to consider the following **Tennessee population estimates for individuals 12 and older** in order to align with the rest of the data that is in this age range by taking off 15% of the above total population estimates.

2013: 5,517,176
 2014: 5,559,006
 2015: 5,602,117
 2016: 5,651,993
 2017: 5,708,586

- The **age-adjusted death rate** for Tennessee in 2016 was calculated to be 886.3 out of 100,000 individuals, or approximately 50,094 people out of a total population 12 and older of 5,651,993 [14]. Subtracting off the number of people who died from

a prescription opioid or heroin/fentanyl overdose in 2016 results in 48,825 people who died that year. This implies that $(5,651,993 - 48,825) / 5,651,993 \approx 0.991$ is the proportion of the population that remains by the beginning of 2017. If we consider T_0 to be the total population in 2016, we can find the continuous-time rate at which individuals die naturally from the equation $0.991T_0 = T_0e^{-\mu t}$, which results in the natural death rate $\mu \approx 0.00868$.

- **Relationship among θ_1 , θ_2 , and θ_3 :** Since we could not find data for values of θ_1 , θ_2 , or θ_3 for Tennessee, we consider a national study of individuals 12 and older to establish a relationship among these three rates. For a national study consisting of 609,000 participants, “the recent heroin incidence rate was 19 times higher among those who reported prior non-medical pain reliever (NMPR) use (0.39%) than among those who did not report NMPR use (0.02%) [15]. Thus, we will extrapolate this information to say that the rate that prescription opioid users and opioid addicts move to heroin use is 19 times greater than the rate at which susceptibles move to heroin use (i.e. $\theta_2 + \theta_3 > 19\theta_1$). Although θ_2 is going from P to H and consists of prescription opioid users who do not misuse their prescription, we will make the assumption that those who misuse are much more likely to be the ones to move to heroin use.

Ignore for now, later tell story about what model tells us regarding this.

Information for moving from recovery to opioid addiction or heroin/fentanyl addiction, σ_A and σ_H :

We consider here the number of individuals in Tennessee age 12 and older who dropped out of treatment and assume that dropping out would result in an individual going back into addiction since they did not successfully complete treatment. The number of drop-outs from medication-assisted opioid detox programs and outpatient medication-assisted opioid programs was reported as essentially zero for Tennessee in 2012, 2013, and 2015, and 3 or fewer for 2011 and 2014, which does not seem realistic [6, 9, 10, 12, 13]. Therefore, to get an estimate for what this value may look like, we took the total number of admissions into all drug treatment programs for each year and calculated the drop-out rates for each of those years:

2011: $2,910/13,422=.217$ [4, 6]

2012: $3,127/13,525=.231$ [5, 9]

2013: $3,273/14,476=.226$ [7, 10]

2014: $3,164/14,909=.212$ [8, 12]

2015: $3,039/14,916=.204$ [11, 13]

We take the average of these drop out rates and extrapolate this to be an estimate for those dropping out of opioid-related therapies: .218. (SEEMS WAY TOO LOW).

References

- [1] American Psychiatric Association (2000). American psychiatric association: *Diagnostic and Statistical Manual of Mental Disorders Fourth Edition*. Available at <https://dsm.psychiatryonline.org/doi/pdf/10.1176/appi.books.9780890420249.dsm-iv-tr> (accessed 10/10/2018).
- [2] Baumblatt, J. A. G., Wiedeman, C., Dunn, J. R., Schaffner, W., Paulozzi, L., and Jones, T. F. (2014). High-risk use by patients prescribed opioids for pain and its role in overdose deaths. *JAMA Internal Medicine*, 175(5):796–801.
- [3] Department of Health and Human Services: Administration for Children and Families, Office of Child Care (2018). State/territory profile-tennessee. Available at: <https://childcareta.acf.hhs.gov/state-profiles/profiles/TN/pdf> (accessed 11/5/18).
- [4] Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (2013). Treatment episode data set (teds) 2001-2011 state admissions to substance abuse treatment services. Available at https://www.dasis.samhsa.gov/dasis2/teds-pubs/2011_teds_rpt_st.pdf (accessed 10/10/2018).
- [5] Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (2014b). Treatment episode data set (teds) 2002-2012 state admissions to substance abuse treatment services. Available at https://www.dasis.samhsa.gov/dasis2/teds-pubs/2012_teds_rpt_st.pdf (accessed 10/10/2018).
- [6] Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (2014a). Treatment episode data set (teds) 2011: Discharges from substance use treatment services. Available at https://www.dasis.samhsa.gov/dasis2/teds-pubs/2011_teds_rpt_d.pdf (accessed 10/10/2018).
- [7] Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (2015). Treatment episode data set (teds) 2003-2013 state admissions to substance abuse treatment services. Available at https://www.dasis.samhsa.gov/dasis2/teds-pubs/2013_teds_rpt_st.pdf (accessed 10/10/2018).
- [8] Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (2016c). Treatment episode data set (teds) 2004-2014 state admissions to substance abuse treatment services. Available at https://www.dasis.samhsa.gov/dasis2/teds-pubs/2014_teds_rpt_st.pdf (accessed 10/10/2018).
- [9] Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (2016a). Treatment episode data set (teds) 2012: Discharges from substance use treatment services. Available at https://www.dasis.samhsa.gov/dasis2/teds-pubs/2012_teds_rpt_d.pdf (accessed 10/10/2018).

- [10] Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (2016b). Treatment episode data set (teds) 2013: Discharges from substance use treatment services: Tables. Available at https://www.dasis.samhsa.gov/dasis2/teds_pubs/2013_teds_rpt_d.pdf (accessed 10/10/2018).
- [11] Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (2017a). Treatment episode data set (teds) 2005-2015 state admissions to substance abuse treatment services. Available at: https://www.dasis.samhsa.gov/dasis2/teds_pubs/2015_teds_rpt_st.pdf (accessed 9/24/18).
- [12] Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (2017b). Treatment episode data set (teds) 2014: Discharges from substance use treatment services. Available at https://www.dasis.samhsa.gov/dasis2/teds_pubs/2014_teds_rpt_d.pdf (accessed 10/10/2018).
- [13] Department of Health and Human Services: Substance Abuse and Mental Health Services Administration (2018). Treatment episode data set (teds) 2015: Discharges from substance use treatment services: Tables. Available at https://www.dasis.samhsa.gov/dasis2/teds_pubs/2015_teds_rpt_d.pdf.
- [14] Henry J Kaiser Family Foundation (2018). Number of deaths per 100,000 population. Available at [https://www.kff.org/other/state-indicator/death-rate-per-100000/?currentTimeframe=0&sortModel=\(accessed 11/5/18\)](https://www.kff.org/other/state-indicator/death-rate-per-100000/?currentTimeframe=0&sortModel=(accessed 11/5/18)).
- [15] Muhuri, P. K., Gfroerer, J. C., and Davies, M. C. (2013). Associations of nonmedical pain reliever use and initiation of heroin use in the United States.
- [16] Office of Information and Analytics Tennessee Department of Health. Data dashboard. Available at: <https://www.tn.gov/health/health-program-areas/pdo/pdo/data-dashboard.html> (accessed 8/14/18).
- [17] Office of Information and Analytics: Tennessee Department of Health (2018). Prescription drug overdose program 2018 report. Available at https://www.tn.gov/content/dam/tn/health/documents/pdo/PDO_2018_Report_02.06.18.pdf (accessed).
- [18] Substance Abuse and Mental Health Services Administration. Selected drug use, perceptions of great risk, past year substance use disorder and treatment, and past year mental health measures in tennessee, by age group: Estimated numbers (in thousands), annual averages based on 2015-2016 nsduhs. Available at <https://www.samhsa.gov/data/report/2015-2016-nsduh-state-specific-tables> (accessed 11/27/17).
- [19] United States Census Bureau (2018). National, state, and puerto rico commonwealth totals datasets: Population, population change, and estimated components of population change: April 1, 2010 to july 1, 2017. Available at: <https://www.census.gov/data/datasets/2017/demo/popest/state-total.html#partextimage500989927> (accessed 9/6/18).