

Solutions: Proving The Double Angle Identities with Euclidean Transformations in Two-Dimensional Space

Abstract

Although the angle addition rule suffices to prove the Double Angle Identities for sine and cosine, we invoke some linear algebra to find an alternate proof using the composite map of rotations and a reflection in \mathbb{R}^2 . The resultant identities can be used to derive the matrix representation of a reflection about an arbitrary line L .

Goals

1. Using rotation matrices, prove the following double angle identities

$$\cos(2\theta) = \cos^2(\theta) - \sin^2(\theta)$$

$$\sin(2\theta) = 2\sin(\theta)\cos(\theta)$$

2. Let L be a line through the origin in \mathbb{R}^2 and let θ be the angle between L and the x-axis measured counter-clock wise from the x-axis to L . Argue geometrically that the matrix representing a reflection about L is equal to the composite map,

$$Ref_L = R_\theta \circ Ref_{x-axis} \circ R_{-\theta}$$

where R_θ is a clockwise rotation by θ , $R_{-\theta}$ is a counter clockwise rotation by θ and Ref_{x-axis} represents a reflection about the x-axis.

3. Compute the matrix for Ref_L in terms of θ

Solutions

Clockwise and counter-clockwise rotations by θ in \mathbb{R}^2 can be described by the following matrices, respectively by

$$R_\theta = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}$$

and

$$R_{-\theta} = \begin{bmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{bmatrix}$$

Now if we rotate a vector by an angle θ and then rotate it again by θ , composing the two transformations, the resultant vector would be the same as if we rotated the original vector once by 2θ , thus


$$R_{2\theta} = \begin{bmatrix} \cos(2\theta) & -\sin(2\theta) \\ \sin(2\theta) & \cos(2\theta) \end{bmatrix} = R_\theta \circ R_\theta = \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix}^2$$

which yields


$$\begin{bmatrix} \cos^2(\theta) - \sin^2(\theta) & -2\cos(\theta)\sin(\theta) \\ 2\sin(\theta)\cos(\theta) & \cos^2(\theta) - \sin^2(\theta) \end{bmatrix}$$

This shows that the first and third matrices must be equal and, matching the elements of the matrices, we see that $\cos(2\theta)$ is indeed equal to $\cos^2(\theta) - \sin^2(\theta)$ and that $\sin(2\theta) = 2\cos(\theta)\sin(\theta)$


Now let's take a look at these vectors. It is given that θ is the measure of the angle from L to the X-axis.




Our definition of Ref_L is that if we take an arbitrary vector in \mathbb{R}^2 (call it V_1), rotate it counterclockwise by θ to get a vector V_2 ,



then reflect it about the x-axis to get a vector V_3 ...



and then finally rotate it clockwise by θ , we will get a vector V_4 which is identical to V_L



therefore

$$Ref_L = R_\theta \circ Ref_{x-axis} \circ R_{-\theta}$$

.

And as a function of θ , Ref_L could be calculated as

$$\begin{aligned} Ref_L &= \begin{bmatrix} \cos(\theta) & -\sin(\theta) \\ \sin(\theta) & \cos(\theta) \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \begin{bmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{bmatrix} \\ &= \begin{bmatrix} \cos(\theta) & \sin(\theta) \\ \sin(\theta) & -\cos(\theta) \end{bmatrix} \begin{bmatrix} \cos\theta & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{bmatrix} \\ &= \begin{bmatrix} \cos^2\theta - \sin^2\theta & 2\sin(\theta)\cos(\theta) \\ 2\sin(\theta)\cos(\theta) & \sin^2(\theta) - \cos^2(\theta) \end{bmatrix} \end{aligned}$$

We can further simplify this matrix by employing the double angle formulas, reducing our result to

$$Ref_L(\theta) = \begin{bmatrix} \cos(2\theta) & \sin(2\theta) \\ \sin(2\theta) & -\cos(2\theta) \end{bmatrix}$$

, a reflection about the line L in \mathbb{R}^2