

# IN THE CRUCIBLE: MASSIVE COMPACT GALAXIES WITH HIGH-VELOCITY OUTFLOWS AND EDDINGTON-LIMITED STAR FORMATION

ALEKSANDAR M. DIAMOND-STANIC<sup>1,2</sup>, JOHN MOUSTAKAS<sup>1</sup>, CHRISTY A. TREMONTI<sup>3</sup>, ALISON L. COIL<sup>1</sup>, RYAN C. HICKOX<sup>4</sup>, ADAY ROBAINA<sup>5</sup>, GREGORY H. RUDNICK<sup>6</sup>, & PAUL SELL<sup>3</sup>

*Draft version April 20, 2012*

## ABSTRACT

We present the discovery of compact, obscured star formation in galaxies that exhibit  $\gtrsim 1000 \text{ km s}^{-1}$  outflows. Using optical morphologies from the Hubble Space Telescope and infrared photometry from the Wide-field Infrared Survey Explorer, we estimate star formation rate (SFR) surface densities that approach  $\Sigma_{\text{SFR}} \approx 5000 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$ , which is comparable to the Eddington limit from radiation pressure on dust grains. We argue that feedback associated with a compact starburst in the form of radiation pressure from massive stars and ram pressure from supernovae and stellar winds is sufficient to produce the high-velocity outflows we observe, without the need to invoke feedback from an active galactic nucleus.

*Subject headings:* galaxies: evolution — galaxies: kinematics and dynamics — galaxies: ISM — galaxies: starburst

## 1. INTRODUCTION

The central regions of elliptical galaxies are thought to form in compact starbursts (e.g., Kormendy et al. 2009; Hopkins et al. 2009). Feedback associated with such starbursts can produce outflows driven by thermal energy from supernova explosions (e.g., Chevalier & Clegg 1985), stellar winds (e.g., Leitherer et al. 1992), and momentum input from both supernova ram pressure and radiation pressure on dust grains (e.g., Murray et al. 2005). It has been argued that such feedback imposes a limit on the maximum star-formation rate (SFR) surface density ( $\Sigma_{\text{SFR}}$ ) for starbursts (e.g., Lehnert & Heckman 1996; Meurer et al. 1997; Murray et al. 2005; Thompson et al. 2005) and the maximum stellar surface density for elliptical galaxies and star clusters (e.g., Hopkins et al. 2010).

Galactic winds are ubiquitous in star-forming galaxies at all redshifts and generally exhibit outflow velocities in the  $100\text{--}500 \text{ km s}^{-1}$  range (e.g., Heckman et al. 2000; Shapley et al. 2003; Martin 2005; Rupke et al. 2005; Weiner et al. 2009; Rubin et al. 2010; Steidel et al. 2010). Thus, the discovery of  $|v| > 1000 \text{ km s}^{-1}$  outflows in a sample of massive ( $M_{*} \approx 10^{11} \text{ M}_{\odot}$ ) post-starburst galaxies at  $z \sim 0.6$  by Tremonti et al. (2007) suggested that a more energetic source, such as feedback from an accreting supermassive black hole (e.g., Silk & Rees 1998; Di Matteo et al. 2005), could have been responsible for launching the winds (see Fabian 2012, for a recent review).

However, it also plausible that feedback from a compact starburst could expel gas with such large velocities. Indeed, there is evidence for a positive correlation

between outflow velocity and starburst luminosity (e.g., Martin 2005; Rupke et al. 2005; Tremonti et al. 2007), albeit with significant scatter. Furthermore, Heckman et al. (2011) recently found outflows with maximum velocities reaching  $\sim 1500 \text{ km s}^{-1}$  in a sample of local starbursts with compact nuclei, and argued that such velocities could be explained by feedback from massive stars.

In this Letter, we present results for a sample of massive galaxies at  $z \sim 0.6$  that exhibit  $\gtrsim 1000 \text{ km s}^{-1}$  outflows, expanding on the sample from Tremonti et al. (2007). We seek to test whether the energetic outflows in these galaxies could have been driven by feedback from starbursts with very large SFR surface densities. Our analysis combines galaxy sizes obtained with the Hubble Space Telescope (HST) with SFRs and stellar masses estimated using photometry from the Wide-field Infrared Survey Explorer (WISE), the Spitzer Space Telescope, the Sloan Digital Sky Survey (SDSS), and the Galaxy Evolution Explorer (GALEX).

## 2. ANALYSIS

### 2.1. Morphologies and Sizes

Our sample derives from 29 galaxies targeted for HST/WFC3 imaging in programs 12019 and 12272 (PI: C. Tremonti). Using the F814W filter on the UVIS channel, which has  $0.04''$  pixels and  $\text{FWHM} = 0.074''$ , we obtained  $4 \times 10$  min exposures in a single orbit for each galaxy. The dithered images were processed with MultiDrizzle<sup>7</sup> to produce science mosaics with  $0.02''$  pixels. For each galaxy, we use GALFIT (Peng et al. 2002) to model the two-dimensional surface brightness profile with a single Sersic component, using stars in the images to construct the model point-spread function (PSF). The Sersic index ( $n$ ) and effective radius ( $r_e$ ) are free parameters in the model. In cases where the best-fit model returns  $n > 4$ , we also fit an  $n = 4$  de Vaucouleurs model, yielding a larger  $r_e$  value (due to the covariance between  $n$  and  $r_e$ ), and we use these these larger effective radii in our analysis.

<sup>1</sup> Center for Astrophysics and Space Sciences, University of California, San Diego, La Jolla, CA 92093, USA

<sup>2</sup> Center for Galaxy Evolution Fellow

<sup>3</sup> Department of Astronomy, University of Wisconsin-Madison, Madison, WI 53706, USA

<sup>4</sup> Department of Physics and Astronomy, Dartmouth College, Hanover, NH 03755, USA

<sup>5</sup> Institut de Ciències del Cosmos, University of Barcelona, 08028 Barcelona, Spain

<sup>6</sup> Department of Physics and Astronomy, University of Kansas, Lawrence, KS 66045, USA

<sup>7</sup> <http://stdas.stsci.edu/multidrizzle/>




FIG. 1.— Left: Observed UV-IR SEDs ( $\lambda_{\text{rest}} = 0.1\text{--}15\ \mu\text{m}$ ) for three galaxies with large SFR surface densities ( $\Sigma_{\text{SFR}} > 3000\ \text{M}_{\odot}\ \text{yr}^{-1}\ \text{kpc}^{-2}$ ). We show stellar population fits to the  $\lambda_{\text{rest}} = 0.1\text{--}3\ \mu\text{m}$  emission (black solid line) and two templates for dust emission (M82 starburst, Arp 220 starburst; Polletta et al. 2007) scaled to match the  $\lambda_{\text{obs}} = 4.6\ \mu\text{m}$  band. The starburst templates provide reasonable fits to the  $\lambda_{\text{obs}} = 12$  and  $22\ \mu\text{m}$  WISE photometry, while a quasar template does not. Right: HST/WFC3 F814W images (probing  $\lambda_{\text{rest}} \approx 5000\ \text{\AA}$ ) showing that these galaxies are dominated by a compact nucleus.

For this paper, we are most interested in the galaxies with the largest  $\Sigma_{\text{SFR}}$  values, which also have the smallest effective radii. We show the HST images for three high- $\Sigma_{\text{SFR}}$  galaxies in Figure 1. In all three cases, the single-component model GALFIT accounts for  $> 85\%$  of the total flux. The residuals show diffuse emission that is consistent with these systems being late-stage galaxy mergers after final coalescence, although we defer a detailed comparison of the observed morphologies with expectations from merger simulations to future work.

For the most compact galaxy (J0905+5759,  $r_e = 0.013''$  or  $100\ \text{pc}$ ), we also show the observed one-dimensional surface brightness profile in the bottom-right panel of Figure 2.1. We compare to the profiles of six stars in the same image, the best-fit de Vaucouleurs model, and a de Vaucouleurs model with a larger effective radius ( $r_e = 0.04''$ , the native WFC3/UVIS pixel size, which corresponds to a physical scale of  $290\ \text{pc}$ ). This comparison illustrates that this galaxy, while only marginally resolved with an effective radius that is  $\sim 20\%$  of the image FWHM, is clearly more extended than a point source.

For such a compact source, there is significant uncertainty in our  $r_e$  measurement given uncertainties in the model PSF. To quantify this uncertainty, we used TinyTim to generate a model PSF that is artificially narrower than the stars in the image (convolving the output of tiny2 with a FWHM =  $0.04''$  Gaussian, whereas the image FWHM is  $0.074''$ ) and found that this increased the effective radius in the GALFIT model by a factor of two. We also fit a two-component PSF+Sersic model, but found that the Sersic component dominates the fit, yielding a similar effective radius. Furthermore, the spectrum of the galaxy shows no evidence for an AGN contribution to the  $I$ -band continuum (see Figure 3), so




FIG. 2.— One-dimensional surface brightness profile for J0905+5759, which has the smallest effective radius in the sample. The observed profile is shown as the solid black line, and the profiles of six stars in the same image are shown in blue. The best-fit de Vaucouleurs profile with  $r_e = 100\ \text{pc}$  is shown as a dashed red line, and for comparison a broader profile with  $r_e = 0.04'' = 290\ \text{pc}$  is shown as the dotted green line. This galaxy is quite compact, but is more extended than a point source.

there is no clear motivation for including an unresolved, point-source component in the model. We conclude that this galaxy is quite compact and that our  $r_e$  estimate of  $100\ \text{pc}$ , while uncertain, is likely accurate within a factor of two.

## 2.2. Star Formation Rates and Stellar Masses

We gathered photometry from the WISE All-Sky Release, the SDSS Seventh Data Release, and GALEX General Release 6. We also obtained  $5 \times 30\ \text{sec}$  dithered exposures at  $3.6\ \mu\text{m}$  and  $4.5\ \mu\text{m}$  for all sources with the Infrared Array Camera on the Spitzer Space Telescope as part of General Observer program 60145 (PI: C. Tremonti). We used the post-basic calibrated data to perform aperture photometry on all sources and point-source photometry on sources in crowded fields. We show spectral energy distributions (SEDs) for the three high- $\Sigma_{\text{SFR}}$  galaxies in Figure 1.

We estimate IR-based SFRs for the 25/29 galaxies with WISE 12 or  $22\ \mu\text{m}$  detections by fitting Chary & Elbaz (2001) templates to their 12 and  $22\ \mu\text{m}$  fluxes. For the 14/29 galaxies with  $22\ \mu\text{m}$  detections, this yields SFRs that agree with those obtained from the Rujopakarn et al. (2012) method based on  $24\ \mu\text{m}$  luminosity with a scatter of  $0.05\ \text{dex}$ . We note that several authors have shown that the shape of the IR SED for star-forming galaxies depends on the surface density of star formation (e.g., Rujopakarn et al. 2011; Elbaz et al. 2011), with more compact starbursts having larger total-IR ( $8\text{--}1000\ \mu\text{m}$ ) to mid-IR ( $8\text{--}24\ \mu\text{m}$ ) ratios. The extreme  $\Sigma_{\text{SFR}}$  values of our sample imply large total-IR to mid-IR ratios, characteristic of the most luminous galaxies in the local universe (e.g., Rieke et al. 2009). If we used the most luminous local templates for the 8/29 sources with SFRs in the ULIRG regime ( $\text{SFR}_{\text{IR}} > 200\ \text{M}_{\odot}\ \text{yr}^{-1}$ ), we would obtain SFRs that are larger by  $0.5\ \text{dex}$  than the values we adopt for this paper.




FIG. 3.— Spectra covering  $\lambda_{\text{rest}} = 2500\text{--}5200 \text{ \AA}$  for the three galaxies shown in Figure 1. These spectra are dominated by light from the young stellar population and exhibit [O II]  $\lambda 3727$ , H $\beta$   $\lambda 4861$ , and [O III]  $\lambda 5007$  emission lines. The panels on the right highlight the Mg II  $\lambda\lambda 2796, 2803$  spectral region in both wavelength and velocity space to illustrate the outflow kinematics. The spectrum in the bottom panel has sufficient spectral resolution (FWHM  $\approx 8 \text{ km s}^{-1}$ ) to resolve the intrinsic shape of the absorption-line profile, revealing that the gas near the centroid velocity ( $v = -2470 \text{ km s}^{-1}$ ) covers the entire galaxy.

We also estimate SFRs and stellar masses based on stellar population fits to the  $\lambda_{\text{rest}} = 0.1\text{--}3 \text{ }\mu\text{m}$  SEDs using the method of Moustakas et al. (2011). For galaxies with  $\text{SFR}_{\text{IR}} > 50 \text{ M}_{\odot} \text{ yr}^{-1}$ , there is agreement between these UV-based SFR estimates and  $\text{SFR}_{\text{IR}}$  with a scatter of 0.32 dex. For an SMC dust law, we find a median attenuation of  $A_V = 0.4 \text{ mag}$ .

### 2.3. Outflow Kinematics and Covering Factors

We present  $\lambda_{\text{rest}} = 2500\text{--}5200 \text{ \AA}$  spectroscopy for three high- $\Sigma_{\text{SFR}}$  sources in Figure 3 based on data from MMT/Blue Channel and SDSS (J1506+5402), Magellan/MagE (J1341-0321), and Keck/LRIS and Keck/HIRES (J0905+5759). The spectra are dominated by light from the young ( $t < 50 \text{ Myr}$ ) stellar population. We highlight the Mg II  $\lambda\lambda 2796, 2803$  absorption lines, which are used to measure outflow velocities. At low spectral resolution (e.g., the top right panel of Figure 3) it is not possible to determine the intrinsic shape of the absorption line profile and therefore the covering factor of the outflowing gas. However, the Keck/HIRES spectrum of J0905+5759 (FWHM  $\approx 8 \text{ km s}^{-1}$ ) reveals that the gas covers the entire continuum source near the velocity centroid ( $v = -2470 \text{ km s}^{-1}$ ) indicating a galaxy-wide outflow.

## 3. DISCUSSION

The compact sizes ( $r_e \approx 100 \text{ pc}$ ) and large SFRs ( $\text{SFR} \approx 300 \text{ M}_{\odot}$ ) for the galaxies described above implies extremely large SFR surface densities ( $\Sigma_{\text{SFR}} \approx 5000 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$ ). To place these galaxies in context, we plot their  $\Sigma_{\text{SFR}}$  values as a function of stellar mass in Figure 4. We include comparison samples of  $\sim 10^5$  star-forming galaxies at  $0.5 < z < 1.5$  from Wuyts et al. (2011) and a few dozen gas-rich mergers

at  $z < 0.3$  including 32 ULIRGs from Veilleux et al. (2006), five Lyman break analogs with dominant central objects from Overzier et al. (2009), and the local compact starburst Arp 220 (Scoville et al. 1997; Kennicutt 1998; Rodríguez Zaurín et al. 2008). We also mark the empirical threshold for launching winds ( $\Sigma_{\text{SFR}} \approx 0.1 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$ , Heckman 2002), the 90th percentile limit for the surface brightness of starbursts measured using UV, H $\alpha$ , far-IR, and radio continuum emission ( $\Sigma_{\text{SFR}} \approx 45 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$  Meurer et al. 1997), and the theoretical limit for a starburst limited by feedback from radiation pressure ( $\Sigma_{\text{SFR}} \approx 2000 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$ , Murray et al. 2005; Thompson et al. 2005; Hopkins et al. 2010). The most luminous, compact starbursts in our sample exhibit SFR surface densities that reach the Eddington limit, suggesting that their growth is being regulated by momentum input from massive stars.

### 3.1. Constraints on Ongoing AGN Activity

While the SEDs (Figure 1) and optical spectra (Figure 3) for our sample indicate that their bolometric output is dominated by star formation, it is worthwhile to consider the level of ongoing AGN activity and its effect on our measurements. Among the high- $\Sigma_{\text{SFR}}$  galaxies, the strongest case for detectable AGN emission can be made for J1506+5402, which has the most luminous [O III]  $\lambda 5007$  line among the 29 galaxies observed by HST. This source also has a weak [Ne V]  $\lambda 3426$  emission line, which is associated with AGN activity (e.g., Gilli et al. 2010). It was observed with the Chandra X-ray Observatory (proposal ID 11700896, PI: C. Tremonti) and had four detected counts, corresponding to a 0.5–8.0 keV X-ray luminosity of  $10^{42.7} \text{ erg s}^{-1}$  (see P. Sell et al., in preparation for more details on the 12/29 galaxies with Chandra observations). Using the relationship between 2–10 keV X-ray and 12.3  $\mu\text{m}$  mid-IR luminosity for AGNs from Gandhi et al. (2009), one would expect a source with  $L_X \approx 5 \times 10^{42} \text{ erg s}^{-1}$  to have  $L_{\text{MIR}} \approx 7 \times 10^{42} \text{ erg s}^{-1}$ , which is  $400\times$  smaller than observed luminosity for J1506+5402 ( $L_{\text{MIR}} \approx 3 \times 10^{45} \text{ erg s}^{-1}$ ). Based on its [O III] luminosity ( $10^{42.1} \text{ erg s}^{-1}$ , a significant fraction of which could be attributed to star formation) and the calibration for type 1 AGNs from Heckman et al. (2005), one would expect an intrinsic 2–10 keV luminosity of  $L_X \approx 5 \times 10^{43} \text{ erg s}^{-1}$ , suggesting absorption by a factor of  $\sim 10$  in the X-rays. However, even if the X-ray attenuation were a factor of 100, which is the typical value for local Compton-thick AGNs (Diamond-Stanic et al. 2009), the expected mid-IR AGN contribution would only be relevant at the  $\lesssim 30\%$  level. We conclude that the bolometric output of the galaxies in our sample is dominated by star formation and that our results regarding large  $\Sigma_{\text{SFR}}$  values are not affected by AGN contamination.

### 3.2. The Outflow Launching Mechanism

It is worth considering whether the high-velocity outflows we observe could have been produced by the compact starburst. Murray et al. (2011) argued that massive star clusters with large gas surface densities are the ideal launching point for galactic-scale outflows driven by radiation pressure, and that the outflow velocity should scale with escape velocity of the most massive star clusters in a galaxy. The  $\gtrsim 1000 \text{ km s}^{-1}$  outflow velocities




FIG. 4.— SFR surface densities and stellar masses for the HST-WISE sample described in this paper (black circles, with symbol size proportional to outflow velocity), along with samples of  $z < 0.3$  gas-rich mergers (orange squares) and  $z \sim 1$  star-forming galaxies (shown with 68%, 95%, and 99.7% contours). We mark the empirical threshold for launching winds (dotted line,  $\Sigma_{\text{SFR}} \approx 0.1 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$ ; Heckman 2002), the 90th-percentile starburst intensity limit from Meurer et al. (1997) (dashed line,  $\Sigma_{\text{SFR}} \approx 45 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$ ), and the Eddington limit from radiation pressure on dust grains (solid line,  $\Sigma_{\text{SFR}} \approx 2000 \text{ M}_{\odot} \text{ yr}^{-1} \text{ kpc}^{-2}$ ; Murray et al. 2005; Thompson et al. 2005; Hopkins et al. 2010). Our HST-WISE sample overlaps with the region characterized by gas-rich mergers, and extends to very large SFR surface densities near the Eddington limit, suggesting growth that is limited by momentum injection from massive stars.

we observe correspond approximately to the escape velocities for a compact stellar population with the sizes and masses that we’ve measured.

$$v_{\text{esc}} = \sqrt{2GM_*/r} \\ = 2100 \left( \frac{M_*}{10^{11} M_{\odot}} \right)^{1/2} \left( \frac{r}{200 \text{ pc}} \right)^{-1/2} \text{ km s}^{-1} \quad (1)$$

This argument, combined with the fact that we observe galaxies with significant dust-obscured star formation and  $\Sigma_{\text{SFR}}$  values near the Eddington limit, indicates that momentum input from massive stars in the form of radiation pressure is a viable mechanism for launching these outflows.

In addition to radiation pressure, we also expect significant momentum flux from stellar winds and supernovae. For example, a starburst with  $\text{SFR} \approx 200 \text{ M}_{\odot} \text{ yr}^{-1}$  would be associated with radiation pressure  $L_{\text{bol}}/c \approx 2 \times 10^{35} \text{ dyne}$  and ram pressure  $\dot{p} \approx 5\text{--}10 \times 10^{35} \text{ dyne}$  from stellar winds and supernovae (e.g., Leitherer et al. 1992,

1999; Veilleux et al. 2005). Furthermore, Heckman et al. (2011) noted that such a large momentum injection ( $\dot{p} \approx 10^{35} \text{ dyne}$ ) from a small initial radius  $r_0 \approx 100 \text{ pc}$  could accelerate a cloud with column density  $N_H \approx 10^{21} \text{ cm}^{-2}$  to a terminal velocity  $v_{\infty} \approx 1800 \text{ km s}^{-1}$ . Thus, the energetics of compact starbursts are sufficient to produce the high-velocity outflows we observe, and it is plausible that both radiation pressure on dust grains and supernova ram pressure contribute to driving the winds.

### 3.3. Placing These Galaxies in Context

It is clear from Figure 4 that the high- $\Sigma_{\text{SFR}}$  galaxies in our sample constitute a rare population, suggesting that they represent an unusual or short-lived phase. Mergers of gas-rich galaxies are a viable mechanism for producing compact starbursts (e.g., Mihos & Hernquist 1996), and such gas-rich major mergers are rare at  $z \sim 0.6$  due to the decline in both the merger rate (e.g., Lotz et al. 2011) and the gas fraction (e.g., Tacconi et al. 2010) of

galaxies since  $z \sim 2$ . Furthermore, the high- $\Sigma_{\text{SFR}}$  galaxies in our sample are caught in a particular time interval where there is both vigorous star formation and strong feedback. The length of this phase may be set by the gas consumption timescale or the timescale for feedback to suppress subsequent star formation. Based on the Kennicutt–Schmidt (K-S) relation (Kennicutt 1998), a compact starburst with  $r_e \approx 100$  pc,  $\text{SFR} \approx 300 M_\odot$ , and  $\Sigma_{\text{SFR}} \approx 5000 M_\odot \text{ yr}^{-1} \text{ kpc}^{-2}$  would have a gas surface density of  $\Sigma_{\text{gas}} \sim 2 \times 10^{11} M_\odot \text{ kpc}^{-2}$  corresponding to  $M_{\text{gas}} \sim 5 \times 10^9 M_\odot$  inside 100 pc, which would be consumed on a timescale  $\sim 30$  Myr. This scenario could be tested with CO observations of molecular gas masses and kinematics for these extreme galaxies.

Considering our full sample of galaxies in Figure 4, a vertical  $\Sigma_{\text{SFR}}$  sequence is apparent. This could be explained as an evolutionary sequence where the high- $\Sigma_{\text{SFR}}$  galaxies represent the peak of the starburst when the high-velocity outflows are launched, while the lower  $\Sigma_{\text{SFR}}$  galaxies represent a subsequent, post-starburst phase. It is interesting to note that all 12/29 galaxies above the  $\Sigma_{\text{SFR}} \approx 45 M_\odot \text{ yr}^{-1} \text{ kpc}^{-2}$  limit from Meurer et al. (1997) exhibit outflows (with median centroid velocity  $v = -1500 \text{ km s}^{-1}$ ), while all 7/29 galaxies without detected outflows have  $\Sigma_{\text{SFR}} < 30 M_\odot \text{ yr}^{-1} \text{ kpc}^{-2}$ . If a compact starburst is a requirement for the production of high-velocity outflows, it may be that the sources without outflows have not gone through such a phase (see A. Robaina, et al., in preparation for a discussion of the relationship between galaxy morphology, outflow velocity, and stellar population age in this sample).

Finally, it is worthwhile to consider the implications of our results for models of massive galaxy formation. Simulations of major galaxy mergers with large gas fractions ( $f_{\text{gas}} \sim 50\%$ ) can produce  $M_* \sim 10^{11} M_\odot$  remnants with effective radii  $r_e \sim 1$  kpc (e.g., Wuyts et al. 2010), but our sample includes galaxies of similar mass that are smaller by almost an order of magnitude. If

these systems do evolve into quiescent elliptical galaxies, it would be extremely challenging for them to grow in size from  $r_e \sim 0.1$  kpc to  $r_e \sim 5$  kpc to reach the local size–mass relation (Shen et al. 2003) in the  $t \sim 6$  Gyr since  $z = 0.6$ . For comparison the compact, quiescent galaxies observed at  $z \sim 2$  (e.g., Trujillo et al. 2007; van Dokkum et al. 2008) have  $t \sim 10$  Gyr to grow by a factor of  $\sim 5$ . In this context, it is worth noting that the half-light radius (ignoring dust attenuation) at rest-frame  $V$  band (where our HST observations probe) can be a factor of 5–10 smaller than the half-mass radius for a gas-rich merger at final coalescence near the peak of starburst activity (Wuyts et al. 2010). It would be worthwhile to test for such size discrepancies and to probe the radial dependence of the mass-to-light ratio for these galaxies by measuring sizes at rest-frame near-IR wavelengths.

#### 4. CONCLUSIONS

We have measured large SFR surface densities for galaxies that exhibit  $\gtrsim 1000 \text{ km s}^{-1}$  outflows. The largest  $\Sigma_{\text{SFR}}$  values are comparable to the Eddington limit from radiation pressure on dust grains, and such compact starburst are expected to have substantial momentum input from massive stars and supernovae. There is some evidence for AGN activity in a fraction of these systems, but the AGN is generally not relevant in a bolometric sense. While high-velocity outflows have been previously interpreted as a signpost of AGN feedback, we conclude that stellar feedback from a compact starburst is equally capable of producing such a signature.

We acknowledge useful discussions with and assistance from several people. AMD acknowledges support from the Southern California Center for Galaxy Evolution, a multi-campus research program funded by the University of California Office of Research.

#### REFERENCES

- Chary, R., & Elbaz, D. 2001, *ApJ*, 556, 562  
 Chevalier, R. A., & Clegg, A. W. 1985, *Nature*, 317, 44  
 Diamond-Stanic, A. M., Rieke, G. H., & Rigby, J. R. 2009, *ApJ*, 698, 623  
 Di Matteo, T., Springel, V., & Hernquist, L. 2005, *Nature*, 433, 604  
 Elbaz, D., Dickinson, M., Hwang, H. S., et al. 2011, *A&A*, 533, A119  
 Fabian, A. C. 2012, arXiv:1204.4114  
 Gandhi, P., Horst, H., Smette, A., et al. 2009, *A&A*, 502, 457  
 Gilli, R., Vignali, C., Mignoli, M., et al. 2010, *A&A*, 519, A92  
 Heckman, T. M., Lehnert, M. D., Strickland, D. K., & Armus, L. 2000, *ApJS*, 129, 493  
 Heckman, T. M. 2002, *Extragalactic Gas at Low Redshift*, 254, 292  
 Heckman, T. M., Ptak, A., Hornschemeier, A., & Kauffmann, G. 2005, *ApJ*, 634, 161  
 Heckman, T. M., et al. 2011, *ApJ*, 730, 5  
 Hopkins, P. F., Cox, T. J., Dutta, S. N., et al. 2009, *ApJS*, 181, 135  
 Hopkins, P. F., Murray, N., Quataert, E., & Thompson, T. A. 2010, *MNRAS*, 401, L19  
 Kennicutt, R. C., Jr. 1998, *ApJ*, 498, 541  
 Kormendy, J., Fisher, D. B., Cornell, M. E., & Bender, R. 2009, *ApJS*, 182, 216  
 Lehnert, M. D., & Heckman, T. M. 1996, *ApJ*, 472, 546  
 Leitherer, C., Robert, C., & Drissen, L. 1992, *ApJ*, 401, 596  
 Leitherer, C., Schaerer, D., Goldader, J. D., et al. 1999, *ApJS*, 123, 3  
 Lotz, J. M., Jonsson, P., Cox, T. J., et al. 2011, *ApJ*, 742, 103  
 Martin, C. L. 2005, *ApJ*, 621, 227  
 Meurer, G. R., Heckman, T. M., Lehnert, M. D., Leitherer, C., & Lowenthal, J. 1997, *AJ*, 114, 54  
 Mihos, J. C., & Hernquist, L. 1996, *ApJ*, 464, 641  
 Moustakas, J., Zaritsky, D., Brown, M., et al. 2011, arXiv:1112.3300  
 Murray, N., Quataert, E., & Thompson, T. A. 2005, *ApJ*, 618, 569  
 Murray, N., Ménard, B., & Thompson, T. A. 2011, *ApJ*, 735, 66  
 Overzier, R. A., Heckman, T. M., Tremonti, C., et al. 2009, *ApJ*, 706, 203  
 Peng, C. Y., Ho, L. C., Impey, C. D., & Rix, H.-W. 2002, *AJ*, 124, 266  
 Polletta, M., Tajer, M., Maraschi, L., et al. 2007, *ApJ*, 663, 81  
 Rodríguez Zaurín, J., Tadhunter, C. N., & González Delgado, R. M. 2008, *MNRAS*, 384, 875  
 Rieke, G. H., Alonso-Herrero, A., Weiner, B. J., et al. 2009, *ApJ*, 692, 556  
 Rubin, K. H. R., Weiner, B. J., Koo, D. C., Martin, C. L., Prochaska, J. X., Coil, A. L., & Newman, J. A. 2010, *ApJ*, 719, 1503  
 Rujopakarn, W., Rieke, G. H., Eisenstein, D. J., & Juneau, S. 2011, *ApJ*, 726, 93  
 Rujopakarn, W., Rieke, G. H., Weiner, B. J., et al. 2011, arXiv:1107.2921  
 Rupke, D. S., Veilleux, S., & Sanders, D. B. 2005, *ApJS*, 160, 115  
 Scoville, N. Z., Yun, M. S., & Bryant, P. M. 1997, *ApJ*, 484, 702  
 Shapley, A. E., Steidel, C. C., Pettini, M., & Adelberger, K. L. 2003, *ApJ*, 588, 65  
 Shen, S., Mo, H. J., White, S. D. M., et al. 2003, *MNRAS*, 343, 978  
 Silk, J., & Rees, M. J. 1998, *A&A*, 331, L1  
 Steidel, C. C., Erb, D. K., Shapley, A. E., Pettini, M., Reddy, N., Bogosavljević, M., Rudie, G. C., & Rakic, O. 2010, *ApJ*, 717, 289  
 Tacconi, L. J., Genzel, R., Neri, R., et al. 2010, *Nature*, 463, 781

- Thompson, T. A., Quataert, E., & Murray, N. 2005, *ApJ*, 630, 167
- Tremonti, C. A., Moustakas, J., & Diamond-Stanic, A. M. 2007, *ApJ*, 663, L77
- Trujillo, I., Conselice, C. J., Bundy, K., et al. 2007, *MNRAS*, 382, 109
- van Dokkum, P. G., et al. 2008, *ApJ*, 677, L5
- Veilleux, S., Cecil, G., & Bland-Hawthorn, J. 2005, *ARA&A*, 43, 769
- Veilleux, S., Kim, D.-C., Peng, C. Y., et al. 2006, *ApJ*, 643, 707
- Weiner, B. J., et al. 2009, *ApJ*, 692, 187
- Wuyts, S., Cox, T. J., Hayward, C. C., et al. 2010, *ApJ*, 722, 1666
- Wuyts, S., Förster Schreiber, N. M., van der Wel, A., et al. 2011, *ApJ*, 742, 96