

# Practice Questions

April 16, 2024

**Question 1.** Consider a Markov chain with state space  $\{1, 2, 3, 4, 5\}$ , with transition matrix given by

$$P = \begin{bmatrix} 0.3 & 0.7 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 & 0 \\ 0.1 & 0 & 0.6 & 0.2 & 0.1 \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{bmatrix}$$

Compute  $f_{32}$ .

**Question 2.** Consider a Markov chain on the state space  $S = \{0, 1, 2, \dots\}$ . For any  $i \geq 1$ , we define the transition from the state  $i$  as

$$p_{i,i+1} = \frac{i}{2i+1}, \quad \text{and} \quad p_{i,i-1} = \frac{i+1}{2i+1},$$

and  $p_{i,j} = 0$  for  $j \notin \{i-1, i+1\}$ . Show that the Markov chain is null recurrent.

**Question 3.** Let  $(B_t : t \geq 0)$  be a standard Brownian motion.

- If the process  $M_t := \sin(tB_t) - \int_0^t f(B_s)ds$  is a martingale. Write down the function form of  $f$ , and express  $M_t$  in the form of an Itô integral.
- Find the probability  $\mathbb{P}(B_1 > -1 \text{ and } \max_{0 \leq t \leq 1} M_t > 1)$ .
- Apply Itô's formula to the process  $(e^{\lambda B_t - \lambda^2 t/2})_{t \geq 0}$ , and use it to compute the moment generating function of  $\tau$ , where  $\tau := \inf \{t > 0 : |B_t| = 1\}$ .

**Question 4.** Let  $(X_t)_{t \geq 0}$  be a recurrent Markov chain on the state space  $S$ , and let  $V : S \rightarrow \mathbb{R}$  be a real-valued function, such that

$$\sum_{j \in S} p_{i,j} V(j) = V(i), \quad \text{for } i \in S.$$

- If  $V$  is uniformly bounded in  $[0, 1]$ , show that  $V$  is a constant for all states.
- Let the Markov chain be simple symmetric random walk on  $\mathbb{Z}$ . Find a non-constant and unbounded function  $V$  such that the above equation is true.