

## Original Investigation

# Cortical Representation of Afferent Bodily Signals in Borderline Personality Disorder

## Neural Correlates and Relationship to Emotional Dysregulation

Laura E. Müller, MSc; André Schulz, PhD; Martin Andermann, Dipl-Psych; Andrea Gäbel, PhD; Dorothee Maria Gescher, MD; Angelika Spohn, Dipl-Psych; Sabine C. Herpertz, MD; Katja Bertsch, PhD

**IMPORTANCE** The ability to perceive and regulate one's own emotions has been tightly linked to the processing of afferent bodily signals (interoception). Thus, disturbed interoception might contribute to the core feature of emotional dysregulation in borderline personality disorder (BPD), as increased levels of depersonalization, body image disturbances, and reduced sensitivity to physical pain suggest poor body awareness in BPD.

**OBJECTIVE** To determine neural correlates of disturbed body awareness in BPD and its associations with emotional dysregulation and to explore improvements in body awareness with BPD symptom remission.

**DESIGN, SETTING, AND PARTICIPANTS** Case-control study performed at Heidelberg University Hospital, Heidelberg, Germany. Heartbeat evoked potentials (HEPs), an indicator of the cortical representation of afferent signals from the cardiovascular system, were investigated in 34 medication-free patients with BPD, 31 healthy volunteers, and 17 medication-free patients with BPD in remission. The HEPs were assessed using 5-minute resting-state electroencephalograms and parallel electrocardiograms. Core BPD symptoms, history of childhood traumatization, and psychiatric disorders were assessed by means of self-reports and structured interviews. To measure neural correlates of disturbed body awareness, high-resolution T1-weighted structural magnetic resonance imaging scans were collected and analyzed using voxel-based morphometry and region-of-interest-based approaches. The study was performed between 2012 and 2014, and data analysis was performed in 2014.

**MAIN OUTCOMES AND MEASURES** Mean HEP amplitudes in resting-state electroencephalograms and their correlation with self-reported emotional dysregulation, as well as with gray matter volume.

**RESULTS** Patients with BPD had significantly reduced mean HEP amplitudes compared with healthy volunteers ( $F_{1,61} = 11.32, P = .001$ ), whereas the mean HEP amplitudes of patients with BPD in remission lie somewhere in between these 2 groups of participants ( $P > .05$ ). The HEP amplitudes were negatively correlated with emotional dysregulation ( $R = -0.30, P = .01$ ) and positively associated with gray matter volume in the left anterior insula ( $R = 0.53, P < .05$ ) and the bilateral dorsal anterior cingulate cortex ( $R = 0.47, P < .05$ ), 2 structures that have been identified as core regions for interoception.

**CONCLUSIONS AND RELEVANCE** The results indicate state-dependent deficits in the cortical processing of bodily signals in patients with BPD, which appear to be associated with core features of BPD. The analysis of patients with BPD in remission suggests an improvement in cortical representation of bodily signals with symptom remission. Results recommend the integration of techniques to strengthen bodily awareness in psychotherapeutic interventions of BPD.

*JAMA Psychiatry*. doi:10.1001/jamapsychiatry.2015.1252  
Published online September 16, 2015.

← Editorial

+ Supplemental content at  
jamapsychiatry.com

**Author Affiliations:** Department of General Psychiatry, Center for Psychosocial Medicine, University of Heidelberg, Heidelberg, Germany (Müller, Gäbel, Gescher, Herpertz, Bertsch); Institute for Health and Behaviour, Integrative Research Unit on Social and Individual Development, University of Luxembourg, Walferdange, Luxembourg (Schulz); Section of Biomagnetism, Department of Neurology, University of Heidelberg, Heidelberg, Germany (Andermann); Department of Psychosomatic Medicine, Central Institute of Mental Health Mannheim, Medical Faculty Mannheim/Heidelberg University, Heidelberg, Germany (Spohn).

**Corresponding Author:** Katja Bertsch, PhD, Department for General Psychiatry, Center of Psychosocial Medicine, University of Heidelberg, Voßstraße 4, 69115 Heidelberg, Germany (katja.bertsch@med.uni-heidelberg.de).

Emotional dysregulation is a core feature of borderline personality disorder (BPD).<sup>1</sup> Patients with BPD are highly reactive to other people's emotions and show increased activation of the amygdala and anterior insula<sup>2-4</sup> and reduced prefrontal inhibition of these limbic regions for emotional stimuli.<sup>5,6</sup> According to prominent theories,<sup>7,8</sup> these alterations in the perception and regulation of emotions may stem from consistent invalidation (ie, emotional experiences that were ignored, dismissed, or punished during childhood). Growing up in such an invalidating environment may lead individuals with BPD to focus more on external cues to guide their behavior than on internal experiences.

Since the James-Lange theory,<sup>9,10</sup> the ability to perceive and regulate emotions has been tightly linked to the perception of afferent bodily signals (interoception). This has been confirmed by results indicating a positive relationship between interoceptive awareness and the ability to perceive one's own emotions<sup>11-14</sup> and altered processing of interoceptive signals in mental disorders characterized by emotional dysregulation, such as major depression<sup>15,16</sup> or depersonalization.<sup>17,18</sup> Alterations in the processing and regulation of emotions in a patient with BPD might hence be related to deficits in the representation of the patient's own bodily signals.<sup>1,19,20</sup> Indications of interoceptive deficits in patients with BPD come from studies showing increased levels of alexithymia,<sup>21</sup> depersonalization,<sup>22</sup> severe body image disturbance,<sup>23,24</sup> and reduced sensitivity to physical pain.<sup>25,26</sup>

However, the 2 studies<sup>27,28</sup> that investigated the actual interoceptive accuracy (IA) of patients with BPD in interoceptive tasks yielded mixed results. Although reduced IA in a group of patients with personality disorders, including BPD, compared with patients with somatoform disorders and healthy volunteers was reported in Mussgay et al,<sup>27</sup> in a more recent study,<sup>28</sup> no significant differences in IA were found between patients with BPD and healthy volunteers. In these studies,<sup>27,28</sup> IA was assessed with heartbeat perception tasks, which are commonly used, and valid methods.<sup>29-32</sup> Interestingly, IA in these tasks has been associated with volume and activity in the anterior insular, dorsal anterior cingulate, and somatomotor cortices.<sup>33-35</sup> However, these "conventional" methods have one important shortcoming: they are unable to differentiate between actual afferent bodily signals and the active process of attention focusing and perception.<sup>36,37</sup>

By contrast, heartbeat evoked potentials (HEPs),<sup>38-40</sup> cortical potentials measured in the resting electroencephalogram (EEG), are considered as indicators of the cortical representation of afferent signals from the cardiovascular system independent of an attentional focus on the heartbeat.<sup>41-43</sup> This is of importance for the assessment of interoception in patients with BPD because manipulating the attentional focus alters the patients' emotional reactivity.<sup>44</sup> With regard to the neural correlates of HEPs, the results of a dipole localization EEG study<sup>45</sup> indicate that the insular and anterior cingulate cortices, key regions for IA,<sup>33-35</sup> are generators of HEPs. Neuroimaging studies on neural correlates of HEPs are still lacking, and little is known about the clinical relevance of interoception in mental disorders with emotional dysregulation, such as BPD. This is surprising because several successful psychotherapeutic treat-

ments incorporate interventions that focus on the improvement of a patient's awareness of bodily signals<sup>8,46</sup> and may decrease insular hyperreactivity to emotional stimuli (R. Schmitt, PhD, D. Winter, Dipl-Psych, I. Niedtfeld, PhD, S. C. Herpertz, MD, and C. Schmahl, MD, unpublished data, April 2015). Structural and functional alterations in insular and anterior cingulate cortices have been related to emotional dysfunction in patients with BPD.<sup>47</sup> First evidence suggests that cardiac awareness training may improve performance in heartbeat perception tasks, increase HEP amplitudes,<sup>48</sup> and decrease symptom distress in patients with somatoform disorders.<sup>49</sup>

Taken together, there are indications of an association of emotional dysregulation with altered interoception in patients with BPD, but barely anything is known about the cortical representation of bodily signals and its neural correlates. Moreover, a group of patients with BPD in remission might allow for the identification of state-independent psychobiological characteristics. The present study aimed to investigate the cortical representation of bodily signals with HEPs, its association with emotional dysregulation, and its neural correlates in patients with BPD compared with healthy volunteers. We additionally explored HEPs in a group of patients with BPD in remission to gather information on the state dependence of alterations in interoceptive processes. We expected to find smaller HEP amplitudes in patients with BPD than in healthy volunteers, and we assumed that HEPs would correlate negatively with emotional dysregulation. We expected that HEP amplitudes as a measure of the cortical representation of bodily signals, and thus interoceptive signals, would be associated with anterior insular and dorsal anterior cingulate gray matter volume.

## Methods

### Participants

In our study, the participants were 34 medication-free patients who received a *DSM-IV* diagnosis of BPD (20 were female, and the mean [SD] age was 27.7 [6.8] years) and 31 healthy volunteers (16 were female, and the mean [SD] age was 27.9 [6.7] years) matched for age, sex, and intelligence. In addition, we collected EEG data from 17 medication-free female patients with BPD in remission. General exclusion criteria comprised neurological disorders; current alcohol or drug abuse or dependence in the last 12 months; lifetime diagnosis of schizophrenia, schizoaffective disorder, or bipolar disorder; severe medical illness, including cardiac dysregulations; and use of psychotropic medication (Table 1 and eAppendix in the Supplement). The study was part of the German Research Foundation Clinical Research Group 256<sup>50</sup> and was approved by the Ethics Committee of the Medical Faculty of the University of Heidelberg, Heidelberg, Germany, and all participants provided written informed consent.

### Measures

Qualified diagnosticians assessed borderline, comorbid antisocial, or avoidant personality (using the International Personality Disorder Examination<sup>51</sup>) and Axis I disorders (using the Structured Clinical Interview for *DSM-IV* for Axis I disorders<sup>52</sup>), emotional dysregulation (using the Difficulties in Emotion Regu-

Table 1. Demographic, Psychometric, and Self-reported Data and Current Comorbid Psychiatric Diagnoses of Patients

Data	Mean (SD)			$\chi^2$ or <i>F</i> Value	<i>P</i> Value <sup>a</sup>	<i>P</i> Value <sup>a</sup>		
	BPD (n = 34)	BPDR (n = 17)	HV (n = 31)			BPD vs BPDR	BPD vs HV	BPDR vs HV
Sex, No.				$\chi^2_2 = 12.1$	.002	.002	.56	.001
Female	20	17	16					
Male	14	00	15					
Age, y	27.7 (6.8)	30.2 (6.8)	27.9 (6.7)	$F_{2,79} = .96$	.39	.19	.87	.24
IQ	50.9 (8.3)	54.0 (5.6)	53.7 (4.4)	$F_{2,79} = 2.0$	.15	.11	.09	.87
BMI	25.7 (6.4)	27.9 (6.0)	28.7 (27.3)	$F_{2,79} = .248$	.78	.67	.50	.89
Heart rate, beats/min	65.5 (9.8)	68.3 (11.3)	62.5 (13.2)	$F_{2,79} = 1.5$	.24	.41	.30	.10
BPD criteria, IPDE score	6.2 (1.4)	1.0 (1.3)	0.0 (0.0)	$F_{2,79} = 342.8$	≤.001	<.001	<.001	.003
Borderline symptoms, ZAN-BPD score	11.9 (5.0)	3.2 (2.5)	0.4 (0.7)	$F_{2,79} = 95.1$	≤.001	<.001	<.001	.03
Borderline symptoms, BSL score	2.0 (0.6)	0.6 (0.4)	0.1 (0.2)	$F_{2,79} = 171.1$	≤.001	<.001	<.001	<.001
Depressiveness, BDI II score	26.7 (7.7)	9.8 (7.8)	3.0 (2.5)	$F_{2,79} = 121.8$	≤.001	<.001	<.001	<.001
Depersonalization, FDS score	19.7 (11.1)	7.8 (4.8)	2.5 (1.7)	$F_{2,79} = 43.5$	≤.001	<.001	<.001	<.001
Traumatization, CTQ score	57.6 (17.4)	56.1 (19.5)	7.6 (33.1)	$F_{2,75} = 23.2$	≤.001	.74	<.001	<.001
DERS score								
Emotional dysregulation	117.2 (14.8)	84.4 (29.9)	67.0 (19.8)	$F_{2,79} = 54.3$	≤.001	<.001	<.001	.01
Nonacceptance	19.3 (6.8)	13.9 (7.7)	10.0 (3.4)	$F_{2,79} = 19.7$	≤.001	.01	<.001	.06
Goals	18.7 (4.2)	14.0 (3.6)	10.7 (5.1)	$F_{2,79} = 29.9$	≤.001	<.001	<.001	.02
Impulsivity	16.2 (4.3)	11.4 (5.2)	9.2 (3.4)	$F_{2,79} = 23.8$	≤.001	.001	<.001	.09
Aware	19.7 (5.0)	18.3 (4.8)	16.19 (4.5)	$F_{2,79} = 7.6$	≤.001	.33	<.001	.03
Strategies	26.1 (5.5)	15.9 (6.9)	12.7 (6.2)	$F_{2,79} = 42.3$	≤.001	<.001	<.001	.11
Clarity	17.2 (3.6)	11.0 (3.9)	9.1 (3.7)	$F_{2,79} = 40.7$	≤.001	<.001	<.001	.10
Diagnoses, No. (%)								
Affective disorders	10 (29.4)	0 (0)	0 (0)					
Anxiety disorders	13 (38.2)	1 (5.9)	0 (0)					
Eating disorders	9 (26.5)	1 (5.9)	0 (0)					
PTSD	8 (23.5)	0 (0)	0 (0)					
Personality disorders								
Avoidant	6 (17.6)	2 (11.8)	0 (0)					
Antisocial	2 (5.9)	0 (0)	0 (0)					

Abbreviations: BDI II, Beck Depression Inventory II; BMI, body mass index (calculated as weight in kilograms divided by height in meters squared); BPD, patients with borderline personality disorder; BPDR, patients with BPD in remission; BSL, Borderline Symptom List; CTQ, Childhood Traumatization Questionnaire; DERS, Difficulties in Emotion Regulation Scale; FDS, Fewtrell

Depersonalization Scale; HV, healthy volunteers; IPDE, International Personality Disorder Examination; PTSD, posttraumatic stress disorder; ZAN-BPD, Zanarini Rating Scale for Borderline Personality Disorder.

<sup>a</sup> Uncorrected for multiple testing.

lation Scale<sup>53</sup>), depersonalization (using the Fewtrell Depersonalisation Scale<sup>54</sup>), borderline symptom severity (using the Zanarini Rating Scale for Borderline Personality Disorder<sup>55</sup> and the Borderline Symptom List<sup>56</sup>), depressiveness (using the Beck Depression Inventory II<sup>57</sup>), childhood traumatization (using the Childhood Traumatization Questionnaire<sup>58</sup>), and intelligence (using Raven's progressive matrices<sup>59</sup>). Demographic information was assessed in a standardized questionnaire; height and weight were measured on the day of assessment (Table 1).

### Electroencephalographic and Electrocardiographic Data Acquisition

A 5-minute resting-state EEG (eyes closed) and electrocardiogram (ECG; Einthoven II) were recorded in a sound-attenuated, dimly lit room (60 Ag/AgCl electrodes; EasyCap; an equidistant reference system with an average reference and all impedances <10 kΩ) amplified with a 72-channel Quick-

Amp amplifier (Brain Products; with a pass band of 0.01-200 Hz and a sampling rate of 1000 Hz).

### Data Processing

Data preprocessing (BrainVision Analyzer 2.0; Brain Products) included down sampling (250 Hz), digital filtering (band-pass of 0.1-35 Hz and 24-dB/octave roll-off), correction of eye-movement artifacts,<sup>17,38,41,43,60</sup> semiautomatic rejection of trials with nonphysiological artifacts, and segmentation (~200 to 1000 milliseconds of ECG R wave). Separate averages were computed for each electrode and participant. Statistical analyses were conducted on the mean voltage from 455 to 595 milliseconds after the ECG R wave because the impact of the electrocardiac-field artifact in this time window is less than 1%.<sup>17,39,41,43,61,62</sup> For mean HEP amplitudes, we calculated the average across all 60 head electrodes. ARTiiFACT<sup>63</sup> was used to calculate the heart rate.

### Statistical Analyses

Data were subjected to analysis of covariance (ANCOVA), controlling for age and sex, to analyze group differences in HEPs. First, mean HEP amplitudes were compared between patients with BPD and healthy volunteers; heart rate, body mass index, and current affective and anxiety disorders were added as control factors.<sup>64-67</sup> Next, a repeated-measures ANCOVA was performed that included electrode position (Fz, Cz, Pz, and Oz<sup>40,42,43</sup>; using the Huynh-Feldt procedure<sup>68</sup> [to correct for violations of sphericity] and the Dunn multiple comparison test as post hoc tests). Another ANCOVA of the 3 groups of participants (healthy volunteers, patients with BPD, and patients with BPD in remission) was also performed. Partial correlations (controlling for group) were used to explore associations between mean HEP amplitudes and questionnaire data.

### Structural Magnetic Resonance Imaging Data

#### Data Acquisition

A 3-dimensional sagittal isotropic magnetization-prepared rapid acquisition gradient-echo sequence was obtained using a 3-T scanner (Siemens Tim Trio) with a 32-channel head coil (240 slices,  $1 \times 1 \times 1$ -mm voxel size, flip angle of  $9^\circ$ , repetition time of 2300 milliseconds, echo time of 2.98 milliseconds, 256-mm field of view, matrix size of  $256 \times 256$  pixels, and 1-mm slice thickness). Experienced neuroradiologists reviewed all scans to exclude clinically important abnormalities.

#### Data Processing

Processing of structural magnetic resonance imaging (MRI) data was performed with Statistical Parametric Mapping, version 8 (<http://www.fil.ion.ucl.ac.uk/spm>), and the voxel-based morphometry (VBM) toolbox (VBM8; Department of Psychiatry, University of Jena) following the standard VBM8 workflow (eAppendix in the Supplement).

### Statistical Analyses

Using both unbiased VBM (uncorrected  $P < .001$  and an cluster-extension level of  $k > 79$  voxels) and region-of-interest (ROI)-based approaches (familywise error-corrected  $P < .05$ ), the relationship between mean HEP amplitudes and gray matter volumes was assessed with random-effects multiple regression analysis, which included mean HEP amplitudes, group, age, and sex as regressors. Hypothesis-driven ROI analyses were performed for the anterior cingulate and insular cortices (anatomical masks were derived from the Wake Forest University PickAtlas; <http://fmri.wfubmc.edu/software/pickatlas>). To assess the specificity of associations between HEP amplitudes and anterior cingulate and insular gray matter volumes, we performed further ROI analyses for the amygdalae and hippocampi<sup>2-6</sup> and ROI-based group comparisons of anterior cingulate and insular gray matter volumes.

### Source Analysis

#### Data Processing

To emphasize the functional relevance of the regions extracted from the MRI data for the HEP amplitudes, the averaged EEG data were subjected to an adapted form of spatiotemporal source analysis<sup>69-71</sup> implemented in Brain

Electric Source Analysis, version 5.1, with regional sources in the left anterior insular and right anterior cingulate cortices and bilateral opercula (spatial coordinates set to relevant peak voxels from the MRI analysis), as well as a principal component to control for ECG R-wave activity (eAppendix in the Supplement).

### Statistical Analysis

Mean dipole amplitudes in the time window from 455 to 595 milliseconds were subjected to ANCOVA with the factors of group and dipole and with the control factors of age, sex, and residual ECG R-wave activity.

## Results

### HEPs and BPD Symptomatology


The mean HEP amplitudes of patients with BPD were significantly reduced compared with healthy volunteers ( $F_{1,61} = 11.32$ ,  $P = .001$ ) (Figure 1), and the effect remained significant after including control variables ( $P < .05$ ). The group-by-electrode ANCOVA revealed the most pronounced reductions in amplitude in patients with BPD at parietal and occipital electrodes (group by electrode:  $F_{3,165} = 4.24$ ,  $P = .01$ ; at electrodes Pz and Oz,  $P < .01$ ), whereas the mean HEP amplitudes were larger in patients with BPD than in healthy volunteers at the frontal electrodes (ie, at Fz,  $P < .05$ ; eTable in the Supplement). Groups did not differ with regard to heart rate, body mass index, or age ( $P > .05$ ; Table 1).

The exploratory analysis of the mean HEP amplitudes of patients with BPD in remission compared with those of patients with BPD and healthy volunteers confirmed the significant main effect of group ( $F_{2,77} = 6.24$ ,  $P = .003$ ) (Figure 1), with post hoc tests revealing significantly reduced mean HEP amplitudes in patients with BPD compared with healthy volunteers ( $P < .01$ ), whereas the mean HEP amplitudes of patients with BPD in remission lie somewhere in between the patients with BPD and the healthy volunteers and did not significantly differ from the mean HEP amplitudes of these 2 groups ( $P > .05$ ; Figure 1). The mean HEP amplitudes were negatively associated with emotional dysregulation (using the Difficulties in Emotion Regulation Scale:  $R = -0.30$ ,  $P = .01$ ), BPD symptom severity (using the Zanarini Rating Scale for Borderline Personality Disorder:  $R = -0.21$ ,  $P = .06$ ; using the Borderline Symptom List:  $R = -0.28$ ,  $P = .01$ ), depressiveness (using the Beck Depression Inventory II:  $R = -0.25$ ,  $P = .03$ ), and emotional abuse (using the Childhood Traumatization Questionnaire:  $R = -0.21$ ,  $P = .06$ ).

### Structural MRI Analysis

The mean HEP amplitudes were significantly related to gray matter volumes in the dorsal anterior cingulate cortices ( $R = 0.47$ ; Figure 2) and the left anterior insula ( $R = 0.53$ , Figure 2). The VBM analysis also revealed an association between HEP amplitudes and bilateral frontal opercula (Table 2). No significant associations were found between HEP amplitudes and amygdala or hippocampal gray matter volumes, and group comparisons revealed no significant differences in

Figure 1. Mean Heartbeat Evoked Potential (HEP) Amplitudes



BPD indicates borderline personality disorder. Error bars indicate SEM.

insular or anterior cingulate cortex gray matter volumes between patients with BPD and healthy volunteers.

### Source Analysis

Our model with 4 regional sources (the left anterior insula, the right anterior cingulum, the left frontal operculum, and the right frontal operculum) and a principal component for ECG R-wave activity explained 74.4% of the mean dipole amplitude between 455 and 595 milliseconds. The mean dipole amplitudes of patients with BPD were significantly smaller (455-595 milliseconds) than those of healthy volunteers (main effect of group:  $F_{1,60} = 8.55$ ,  $P = .01$ ; no significant effects of dipole, dipole by group, or covariates:  $F_{1,60} \leq 1.74$ ,  $P \geq .19$ ) (Figure 3).

### Discussion

The present study revealed reduced HEP amplitudes, an indicator of cortical processing of bodily signals, in patients with BPD compared with healthy volunteers. In addition, HEP amplitudes correlated negatively with emotional dysregulation and with self-reported emotional childhood abuse. Morphometric analyses revealed specific, positive associations between HEP amplitudes and gray matter volumes of the left anterior insular cortex and the right anterior cingulate cortex. The


analysis of patients with BPD in remission indicated improved cortical representation of bodily signals with symptom remission.

In line with prominent theories such as Damasio's somatic marker hypothesis<sup>72</sup> or the James-Lange theory,<sup>9,10</sup> our results indicate the fundamental role played by afferent bodily signals in the perception and regulation of emotions. Similar to this, a number of studies have revealed positive correlations between interoceptive indicators and the ability to recognize and regulate emotions<sup>19,20</sup> and altered interoceptive signal processing in patients with major depression<sup>15,16</sup> or depersonalization disorder.<sup>17,18</sup> The HEP amplitudes of the patients with BPD were not only significantly reduced compared with healthy volunteers but negatively related to emotional dysregulation. The HEP amplitudes were also positively related to the gray matter volumes of the left anterior insular cortex and the bilateral dorsal anterior cingulate cortex, 2 structures that have been identified as core regions for interoception.<sup>33-35,73</sup> Together with a dipole analysis,<sup>45</sup> this suggests a common neural source of HEP amplitudes and IA measured in conventional heartbeat perception tasks.

Although previous studies highlighted the role played by the right anterior insula in interoceptive awareness<sup>33</sup> and conscious evaluation of bodily signals,<sup>74</sup> we found a strong correlation between HEP amplitudes and the gray matter vol-



**Figure 2. Correlations Between Mean Heartbeat Evoked Potential (HEP) Amplitudes and Gray Matter (GM) Volumes of the Bilateral Dorsal Anterior Cingulate Cortices (dACCs) and the Left Anterior Insula**



BPD indicates borderline personality disorder; L, left; and R, right.

**Table 2. Correlations Between Mean HEP Amplitudes and GM Volume in the Voxel-Based Morphometry and ROI Analyses**

Anatomical Location <sup>a</sup>	Peak MNI Coordinates			Cluster Size, <i>k</i>	<i>t</i> Value
	<i>x</i>	<i>y</i>	<i>z</i>		
Whole-brain analysis <sup>b</sup>					
L ant cingulate cortex/R supp motor area	−15	6	43	941	4.81
R roland operculum/R inf front operculum	52	3	16	223	4.64
R lingual	10	−87	−12	179	4.30
R ant cingulate cortex	20	38	19	1156	4.34
L ant insula/L inf orbitofrontal/L sup temp pole	−34	5	32	901	4.14
L inf front operculum	−48	12	19	305	3.93
Cerebellum/brainstem	14	−30	21	87	3.64
ROI analysis <sup>c</sup>					
R ant cingulate cortex	14	32	16	398	4.20
L ant insula	−26	17	−20	101	4.12

Abbreviations: GM, gray matter; HEP, heartbeat evoked potential; L, left; MNI, Montreal Neurological Institute; R, right; ROI, region-of-interest.

<sup>a</sup> Labels are based on the Automated Anatomic Labeling atlas

<sup>b</sup> Uncorrected  $P < .001$  (unbiased voxel-based morphometry approach) and an cluster-extension level of  $k > 79$  voxels.


<sup>c</sup> Familywise error-corrected  $P < .05$ .

ume of the left anterior insula. This is in line with the hypothesis that the left anterior insula is associated with the registration of interoceptive experiences and other low-level cortical processing of interoceptive states, whereas the right anterior insula seems to be of particular relevance for the further evaluation and subjective feeling of bodily signals.<sup>75,76</sup> The finding of reduced cortical representation of bodily signals in patients with BPD may increase our understanding of some of the most characteristic symptoms of this disorder. Parental invalidation, insecurity, and abuse in childhood may result in an individual with BPD focusing on external cues instead of perceiving, integrating, and interpreting his or her own bodily signals and emotions.<sup>7,8</sup> Consistent with this, we found a nega-

tive relationship between self-reported emotional abuse and HEP amplitudes, indicating early trauma-related deficits in the cortical representation of bodily signals that may underlie core BPD symptoms, such as emotional dysregulation or reduced pain sensitivity.<sup>25,26</sup> Despite this, results of 2 previous heartbeat tracking studies<sup>27,28</sup> that have investigated IA in samples that included individuals with BPD are contradictory. To overcome shortcomings of conventional methods, we measured HEP amplitudes in a large sample of patients with BPD, patients with BPD in remission, and healthy volunteers.

Schandry and Montoya<sup>77</sup> have hypothesized that HEP amplitudes at frontal electrodes may reflect a visceral-afferent signal transmission, whereas HEP activity at poste-

Figure 3. Mean Source Waveforms



Mean source waveforms of the source model with 4 regional sources (the left anterior insula, the right anterior cingulum, the left frontal operculum, and the right frontal operculum; spatial coordinates set to peak voxels from the magnetic resonance imaging analysis [Table 2]) and a principal component for electrocardiographic R-wave activity as derived by spatiotemporal source analysis. The principal component analysis (PCA) plot depicts the morphology of the R-wave-associated principal component.

rior electrodes may be related to somatosensory signal processing. In a dipole localization study,<sup>45</sup> HEP activity at frontal sites was associated with dipoles in the anterior cingulate cortex, whereas a dipole originating from the insula headed toward central sites. We observed reduced HEP amplitudes in patients with BPD at parietal and occipital electrodes, whereas the HEP amplitudes did not differ at central sites and were even larger at frontal electrodes compared with healthy volunteers.

In line with previous observations of patients with major depression,<sup>15</sup> not only the amplitude but also the spatial distribution of HEPs differed in patients with BPD, who also had smaller mean dipole amplitudes compared with healthy volunteers. One possible explanation for this could be that patients with BPD do not differ from healthy volunteers in the raw visceral-afferent neurotraffic and lower-brain relaying of this information, but show deficits in the integration of afferent bodily signals from different (eg, visceral and somatosensory) sources into neural self-representation. Alternatively, the different spatial distribution of HEP amplitudes in patients with BPD may reflect a disrupted cortical representation of visceral-afferent signals, which is compensated by afferent bodily signals from other sources, such as somatosensory or auditory sensations.<sup>78,79</sup> Because no differences in heart rate were found among the 3 groups of participants, the differences in HEP amplitudes may be attributed to alterations in central representation, not to peripheral differences.

The question remains as to why we have observed reduced mean HEP and dipole amplitudes in patients with BPD and positive correlations between HEP amplitudes and gray matter volumes in the insular and anterior cingulate cortices and frontal opercula, but no group differences in the gray matter volume of these regions. A possible explanation could be that HEP amplitudes not only reflect volume but also activity (as suggested by the dipole analysis) and the orchestral interplay or functional connectivity of these regions.

An additional analysis revealed that the HEP amplitudes of a smaller group of patient with BPD in remission lie somewhere in between those of patients with BPD and healthy volunteers, indicating an amelioration of the cortical representation of bodily signals with symptom remission. Although we do not have specific information on individual treatment history, it is well known that the most successful and widespread psychotherapies for BPD include elements of emotion recognition and regulation, the perception and integration of bodily signals, and mindfulness.<sup>8,80,81</sup> Owing to the cross-sectional nature of our study, the current finding can only be regarded as a first indicator of a possible improvement of interoceptive signal processing with symptom remission. The mechanisms of this relationship remain to be elucidated in longitudinal studies. Together with previous studies, the current results indicate that treatments focusing on body awareness not only may be a promising instrument to improve emotional dysregulation but might also decrease the enhanced risk for cardiovascular disease among patients with BPD.<sup>82</sup> Because HEP amplitudes might represent a state-dependent illness rather than a trait-dependent disorder marker, they could be used as an indicator of treatment response and symptom reduction in intervention studies.

To our knowledge, this is the first study to examine the cortical representation of bodily signals by use of HEP amplitudes and its neural correlates in BPD. However, several limitations of the present study should be noted. First, diagnostic interviews revealed a number of comorbidities. Although this reflects a typical pattern of comorbid psychiatric disorders in BPD<sup>83</sup> and effects remained significant after controlling for affective and anxiety disorders, we cannot rule out that the neurophysiological characteristics of these disorders may have affected our results. Future studies should include clinical control groups and should assess psychotherapeutic treatment experiences. Second, only exploratory

comparisons were possible with the smaller group of female patients with BPD in remission, and the exploratory correlation analyses were not corrected for multiple testing. Third, we decided not to include conventional heartbeat perception tasks, which have a limited face validity for the clinical phenomenology of BPD to separate the genuine cortical representation of visceral-afferent signals from possible attentional or evaluative processes. It remains an interesting question whether patients with BPD have specific deficits in the automatic, nonconscious cortical representation of bodily signals, which they are able to compensate for by motivation<sup>84</sup> or focused attention<sup>36</sup> in conscious tasks. Further studies, which should include different measures for the assessment of interoception and of emotion processing and regulation in patients with BPD, are needed.

## Conclusions

Taken together, the present study revealed reduced HEP amplitudes, an indicator of the cortical representation of bodily signals, in patients with BPD, a severe mental disorder that is characterized by emotional dysregulation, compared with healthy volunteers. In line with previous studies, morphometric analysis suggests that the anterior insular and the dorsal anterior cingulate cortices are neural correlates of interoceptive signal processing. Together with the additional, exploratory analysis of patients with BPD in remission, these results indicate deficits in interoceptive processes in patients with BPD, which may be a promising target for psychotherapeutic interventions.

## ARTICLE INFORMATION

**Submitted for Publication:** March 1, 2015; final revision received June 10, 2015; accepted June 10, 2015.

**Published Online:** September 16, 2015.  
doi:10.1001/jamapsychiatry.2015.1252.

**Author Contributions:** Ms Müller and Dr Bertsch had full access to all of the data in the study and take responsibility for the integrity of the data and the accuracy of the data analysis.  
*Study concept and design:* Schulz, Herpertz, Bertsch.

*Acquisition, analysis, or interpretation of data:* Müller, Schulz, Andermann, Gäbel, Gescher, Spohn, Bertsch.

*Drafting of the manuscript:* Müller, Schulz, Andermann, Bertsch.

*Critical revision of the manuscript for important intellectual content:* Schulz, Andermann, Gäbel, Gescher, Spohn, Herpertz, Bertsch.

*Statistical analysis:* Müller, Schulz, Andermann, Bertsch.

*Obtained funding:* Müller, Herpertz, Bertsch.

*Administrative, technical, or material support:* Gäbel, Gescher, Spohn, Herpertz, Bertsch.

*Study supervision:* Andermann, Herpertz, Bertsch.

**Conflict of Interest Disclosures:** None reported.

**Funding/Support:** The study was supported by grants from the German Research Foundation awarded to Dr Herpertz within the Clinical Research Unit 256 (grants He 2660/12-1 and He 2660/7-2).

**Role of the Funder/Sponsor:** The German Research Foundation had no role in the design and conduct of the study; collection, management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; and decision to submit the manuscript for publication.

**Additional Contributions:** We thank K. Herwig for technically supporting the EEG measurement and analysis, A. Rupp, MD, for feedback on the source analysis, L. Kramer, V. Sebold, and F. Mancke, MD, for their help with data collection, and the team of the Clinical Research Unit on Mechanisms of Disturbed Emotion Processing in Borderline Personality Disorder (Klinische Forschergruppe 256) for participant recruitment and organization. Dr F. Mancke, L. Kramer, and V. Sebold supported the study as members of the Clinical Research Unit, which was financially supported by the German Research Foundation.

## REFERENCES

- Lieb K, Zanarini MC, Schmahl C, Linehan MM, Bohus M. Borderline personality disorder. *Lancet*. 2004;364(9432):453-461.
- Hazlett EA, Zhang J, New AS, et al. Potentiated amygdala response to repeated emotional pictures in borderline personality disorder. *Biol Psychiatry*. 2012;72(6):448-456.
- Herpertz SC, Dietrich TM, Wenning B, et al. Evidence of abnormal amygdala functioning in borderline personality disorder: a functional MRI study. *Biol Psychiatry*. 2001;50(4):292-298.
- Schulz L, Domes G, Krüger A, et al. Neuronal correlates of cognitive reappraisal in borderline patients with affective instability. *Biol Psychiatry*. 2011;69(6):564-573.
- Minzenberg MJ, Fan J, New AS, Tang CY, Siever LJ. Fronto-limbic dysfunction in response to facial emotion in borderline personality disorder: an event-related fMRI study. *Psychiatry Res*. 2007;155(3):231-243.
- Silbersweig D, Clarkin JF, Goldstein M, et al. Failure of frontolimbic inhibitory function in the context of negative emotion in borderline personality disorder. *Am J Psychiatry*. 2007;164(12):1832-1841.
- Bateman A, Fonagy P. *Psychotherapy for Borderline Personality Disorder: Mentalization-Based Treatment*. Oxford, England: Oxford University Press; 2004.
- Linehan MM. *Cognitive-Behavioral Treatment of Borderline Personality Disorder*. New York, NY: Guilford Press; 1993.
- James W. What is an emotion? *Mind*. 1884;9(34):188-205. <http://www.jstor.org/stable/2246769>. Accessed August 12, 2015.
- Lange CG. The mechanisms of the emotions. In: Rand B, ed. *The Classical Psychologist: Selections Illustrating Psychology from Anaxagoras to Wundt*. Boston, MA: Houghton Mifflin; 1885:672-684.
- Schachter S, Singer JE. Cognitive, social, and physiological determinants of emotional state. *Psychol Rev*. 1962;69:379-399.
- Herbert BM, Herbert C, Pollatos O. On the relationship between interoceptive awareness and alexithymia: is interoceptive awareness related to emotional awareness? *J Pers*. 2011;79(5):1149-1175.
- Pollatos O, Kirsch W, Schandry R. On the relationship between interoceptive awareness, emotional experience, and brain processes. *Brain Res Cogn Brain Res*. 2005;25(3):948-962.
- Dunn BD, Galton HC, Morgan R, et al. Listening to your heart: how interoception shapes emotion experience and intuitive decision making. *Psychol Sci*. 2010;21(12):1835-1844.
- Terhaar J, Viola FC, Bär KJ, Debener S. Heartbeat evoked potentials mirror altered body perception in depressed patients. *Clin Neurophysiol*. 2012;123(10):1950-1957.
- Paulus MP, Stein MB. Interoception in anxiety and depression. *Brain Struct Funct*. 2010;214(5-6):451-463.
- Schulz A, Köster S, Beutel ME, et al. Altered patterns of heartbeat-evoked potentials in depersonalization/derealization disorder: neurophysiological evidence for impaired cortical representation of bodily signals. *Psychosom Med*. 2015;77(5):506-516.
- Michal M, Reuchlein B, Adler J, et al. Striking discrepancy of anomalous body experiences with normal interoceptive accuracy in depersonalization-derealization disorder. *PLoS One*. 2014;9(2):e89823.
- Weiss S, Sack M, Henningsen P, Pollatos O. On the interaction of self-regulation, interoception and pain perception. *Psychopathology*. 2014;47(6):377-382.
- Herbert BM, Ulbrich P, Schandry R. Interoceptive sensitivity and physical effort: implications for the self-control of physical load in everyday life. *Psychophysiology*. 2007;44(2):194-202.
- Domes G, Grabe HJ, Czeschnek D, Heinrichs M, Herpertz SC. Alexithymic traits and facial emotion recognition in borderline personality disorder. *Psychother Psychosom*. 2011;80(6):383-385.
- Vermetten E, Spiegel D. Trauma and dissociation: implications for borderline personality disorder. *Curr Psychiatry Rep*. 2014;16(2):434-444.
- Dyer A, Borgmann E, Feldmann RE Jr, et al. Body image disturbance in patients with borderline personality disorder: impact of eating disorders and perceived childhood sexual abuse. *Body Image*. 2013;10(2):220-225.
- Sansone RA, Chu JW, Wiederman MW. Body image and borderline personality disorder among



- psychiatric inpatients. *Compr Psychiatry*. 2010;51(6):579-584.
25. Ludäscher P, von Kalckreuth C, Parzer P, et al. Pain perception in female adolescents with borderline personality disorder. *Eur Child Adolesc Psychiatry*. 2015;24(3):351-357.
  26. Schmahl C, Greffrath W, Baumgärtner U, et al. Differential nociceptive deficits in patients with borderline personality disorder and self-injurious behavior: laser-evoked potentials, spatial discrimination of noxious stimuli, and pain ratings. *Pain*. 2004;110(1-2):470-479.
  27. Mussgay L, Klinkenberg N, Rüdell H. Heart beat perception in patients with depressive, somatoform, and personality disorders. *J Psychophysiol*. 1999;13(1):27-36. doi:10.1027//0269-8803.13.1.27.
  28. Hart N, McGowan J, Minati L, Critchley HD. Emotional regulation and bodily sensation: interoceptive awareness is intact in borderline personality disorder. *J Pers Disord*. 2013;27(4):506-518.
  29. Schandry R. Heart beat perception and emotional experience. *Psychophysiology*. 1981;18(4):483-488.
  30. Whitehead WE, Dreschner VM, Heiman P, Blackwell B. Relation of heart-rate control to heartbeat perception. *Biofeedback Self Regul*. 1977;2(4):371-392. doi:10.1007/BF00998623.
  31. Van der Does AJW, Van Dyck R, Spinhoven P. Accurate heartbeat perception in panic disorder: fact and artefact. *J Affect Disord*. 1997;43(2):121-130.
  32. Phillips GC, Jones GE, Rieger EJ, Snell JB. Effects of the presentation of false heart-rate feedback on the performance of two common heartbeat-detection tasks. *Psychophysiology*. 1999;36(4):504-510.
  33. Critchley HD, Wiens S, Rotshtein P, Ohman A, Dolan RJ. Neural systems supporting interoceptive awareness. *Nat Neurosci*. 2004;7(2):189-195.
  34. Pollatos O, Gramann K, Schandry R. Neural systems connecting interoceptive awareness and feelings. *Hum Brain Mapp*. 2007;28(1):9-18.
  35. Pollatos O, Schandry R, Auer DP, Kaufmann C. Brain structures mediating cardiovascular arousal and interoceptive awareness. *Brain Res*. 2007;1141:178-187.
  36. Montoya P, Schandry R, Müller A. Heartbeat evoked potentials (HEP): topography and influence of cardiac awareness and focus of attention. *Electroencephalogr Clin Neurophysiol*. 1993;88(3):163-172.
  37. Yuan H, Yan HM, Xu XG, Han F, Yan Q. Effect of heartbeat perception on heartbeat evoked potential waves. *Neurosci Bull*. 2007;23(6):357-362.
  38. Pollatos O, Schandry R. Accuracy of heartbeat perception is reflected in the amplitude of the heartbeat-evoked brain potential. *Psychophysiology*. 2004;41(3):476-482.
  39. Gray MA, Taggart P, Sutton PM, et al. A cortical potential reflecting cardiac function. *Proc Natl Acad Sci U S A*. 2007;104(16):6818-6823.
  40. Schandry R, Sparrer B, Weitkunat R. From the heart to the brain: a study of heartbeat contingent scalp potentials. *Int J Neurosci*. 1986;30(4):261-275.
  41. Schulz A, Ferreira de Sá DS, Dierolf AM, et al. Short-term food deprivation increases amplitudes of heartbeat-evoked potentials. *Psychophysiology*. 2015;52(5):695-703.
  42. Shao S, Shen K, Wilder-Smith EP, Li X. Effect of pain perception on the heartbeat evoked potential. *Clin Neurophysiol*. 2011;122(9):1838-1845.
  43. Schulz A, Strelzyk F, Ferreira de Sá DS, Naumann E, Vögele C, Schächinger H. Cortisol rapidly affects amplitudes of heartbeat-evoked brain potentials—implications for the contribution of stress to an altered perception of physical sensations? *Psychoneuroendocrinology*. 2013;38(11):2686-2693.
  44. Baskin-Sommers AR, Vitale JE, Maccoon D, Newman JP. Assessing emotion sensitivity in female offenders with borderline personality symptoms: results from a fear-potentiated startle paradigm. *J Abnorm Psychol*. 2012;121(2):477-483.
  45. Pollatos O, Kirsch W, Schandry R. Brain structures involved in interoceptive awareness and cardioafferent signal processing: a dipole source localization study. *Hum Brain Mapp*. 2005;26(1):54-64.
  46. Kabat-Zinn J. *Full Catastrophe Living: Using the Wisdom of Your Body and Mind to Face Stress, Pain, and Illness*. New York, NY: Delacorte; 1991.
  47. Krause-Utz A, Winter D, Niedtfeld I, Schmahl C. The latest neuroimaging findings in borderline personality disorder. *Curr Psychiatry Rep*. 2014;16(3):438.
  48. Schandry R, Weitkunat R. Enhancement of heartbeat-related brain potentials through cardiac awareness training. *Int J Neurosci*. 1990;53(2-4):243-253.
  49. Schaefer M, Egloff B, Gerlach AL, Witthöft M. Improving heartbeat perception in patients with medically unexplained symptoms reduces symptom distress. *Biol Psychol*. 2014;101:69-76.
  50. Schmahl C, Herpertz SC, Bertsch K, et al. Mechanisms of disturbed emotion processing and social interaction in borderline personality disorder: state of knowledge and research agenda of the German Clinical Research Unit [published online September 9, 2014]. *Borderline Pers Disord Emotion Dysregulation*. doi:10.1186/2051-6673-1-12.
  51. Loranger AW, Sartorius N, Andreoli A, et al. The International Personality Disorder Examination: the World Health Organization/Alcohol, Drug Abuse, and Mental Health Administration international pilot study of personality disorders. *Arch Gen Psychiatry*. 1994;51(3):215-224.
  52. First MB, Spitzer RL, Gibbon M, Williams JBW. *Structured Clinical Interview for DSM-IV (SCID-I)*. New York: New York Biometrics Research Dept; 1995.
  53. Gratz KL, Roemer L. Multidimensional assessment of emotion regulation and dysregulation: development, factor structure, and initial validation of the difficulties in emotion regulation scale. *J Psychopathol Behav*. 2004;26(1):41-54. doi:10.1023/B:JOBA.0000007455.08539.94.
  54. Fawcett WD. *Fewtrell Depersonalisation Scale (FDS)*. Leicester, England: APT Press; 2000.
  55. Zanarini MC, Vujanovic AA, Parachini EA, Boulanger JL, Frankenburg FR, Hennen J. Zanarini Rating Scale for Borderline Personality Disorder (ZAN-BPD): a continuous measure of DSM-IV borderline psychopathology. *J Pers Disord*. 2003;17(3):233-242.
  56. Bohus M, Limberger MF, Frank U, Chapman AL, Köhler T, Stieglitz RD. Psychometric properties of the Borderline Symptom List (BSL). *Psychopathology*. 2007;40(2):126-132.
  57. Beck A, Steer R, Brwon G. *Beck Depression Inventory-II Manual*. San Antonio, TX: The Psychological Corporation; 1996.
  58. Bernstein DP, Fink L. *Childhood Trauma Questionnaire: A Retrospective Self-report Manual*. San Antonio, TX: The Psychological Corporation; 1998.
  59. Heller KA, Kratzmeier H, Lengfelder A. *Matrizen-Test-Manual: Ein Handbuch zu den Advanced Progressive Matrizen von Raven*. Göttingen, Germany: Beltz-Testgesellschaft; 1998.
  60. Gratton G, Coles MG, Donchin E. A new method for off-line removal of ocular artifact. *Electroencephalogr Clin Neurophysiol*. 1983;55(4):468-484.
  61. Dirlich G, Vogl L, Plaschke M, Strian F. Cardiac field effects on the EEG. *Electroencephalogr Clin Neurophysiol*. 1997;102(4):307-315.
  62. Schulz A, Köster S, Beutel ME, et al. Altered patterns of heartbeat-evoked potentials in depersonalization/derealization disorder: neurophysiological evidence for impaired cortical representation of bodily signals. *Psychosom Med*. 2015;77(5):506-516.
  63. Kaufmann T, Sütterlin S, Schulz SM, Vögele C. ARTiiFACT: a tool for heart rate artifact processing and heart rate variability analysis. *Behav Res Methods*. 2011;43(4):1161-1170.
  64. Knapp-Kline K, Kline JP. Heart rate, heart rate variability, and heartbeat detection with the method of constant stimuli: slow and steady wins the race. *Biol Psychol*. 2005;69(3):387-396.
  65. Herbert BM, Blechert J, Hautzinger M, Matthias E, Herbert C. Intuitive eating is associated with interoceptive sensitivity: effects on body mass index. *Appetite*. 2013;70:22-30.
  66. Domschke K, Stevens S, Pfleiderer B, Gerlach AL. Interoceptive sensitivity in anxiety and anxiety disorders: an overview and integration of neurobiological findings. *Clin Psychol Rev*. 2010;30(1):1-11.
  67. Dunn BD, Dalgleish T, Ogilvie AD, Lawrence AD. Heartbeat perception in depression. *Behav Res Ther*. 2007;45(8):1921-1930.
  68. Huynh H, Feldt LS. Estimation of the box correction for degrees of freedom from sample data in randomized block and split-plot designs. *J Educ Stat*. 1976;1(1):69-82. doi:10.2307/1164736.
  69. Scherg M, Vajsar J, Picton TW. A source analysis of the late human auditory evoked potentials. *J Cogn Neurosci*. 1989;1(4):336-355.
  70. Scherg M. Fundamentals of dipole source potential analysis. In: Grandori F, Hoke M, Romani GL, et al. *Auditory Evoked Magnetic Fields and Electric Potentials (Advances in Audiology)*. Vol 6. Basel, Switzerland: Karger; 1990:40-69.
  71. Scherg M, Picton TW. Separation and identification of event-related potential components by brain electric source analysis. *Electroencephalogr Clin Neurophysiol Suppl*. 1991;42:24-37.
  72. Damasio AR. Descartes' error and the future of human life. *Sci Am*. 1994;271(4):144.

73. Couto B, Salles A, Sedeño L, et al. The man who feels two hearts: the different pathways of interoception. *Soc Cogn Affect Neurosci*. 2014;9(9):1253-1260.
74. Strigo IA, Matthews SC, Simmons AN, et al. Altered insula activation during pain anticipation in individuals recovered from anorexia nervosa: evidence of interoceptive dysregulation. *Int J Eat Disord*. 2013;46(1):23-33.
75. Craig AD. How do you feel? interoception: the sense of the physiological condition of the body. *Nat Rev Neurosci*. 2002;3(8):655-666.
76. Rilling JK, Young LJ. The biology of mammalian parenting and its effect on offspring social development. *Science*. 2014;345(6198):771-776.
77. Schandry R, Montoya P. Event-related brain potentials and the processing of cardiac activity. *Biol Psychol*. 1996;42(1-2):75-85.
78. Craig AD. How do you feel—now? the anterior insula and human awareness. *Nat Rev Neurosci*. 2009;10(1):59-70.
79. Wiens S. Interoception in emotional experience. *Curr Opin Neurol*. 2005;18(4):442-447.
80. Feliu-Soler A, Pascual JC, Borràs X, et al. Effects of dialectical behaviour therapy-mindfulness training on emotional reactivity in borderline personality disorder: preliminary results. *Clin Psychol Psychother*. 2014;21(4):363-370.
81. Perroud N, Nicastro R, Jermann F, Huguelet P. Mindfulness skills in borderline personality disorder patients during dialectical behavior therapy: preliminary results. *Int J Psychiatry Clin Pract*. 2012;16(3):189-196.
82. El-Gabalawy R, Katz LY, Sareen J. Comorbidity and associated severity of borderline personality disorder and physical health conditions in a nationally representative sample. *Psychosom Med*. 2010;72(7):641-647.
83. Zanarini MC, Frankenburg FR, Hennen J, Reich DB, Silk KR. Axis I comorbidity in patients with borderline personality disorder: 6-year follow-up and prediction of time to remission. *Am J Psychiatry*. 2004;161(11):2108-2114.
84. Weiskunat R, Schandry R. Motivation and heartbeat evoked potentials. *J Psychophysiol*. 1990;4(1):33-40.