

# **Q Notes**

By Peter Montgomery

December 16, 2020

# Contents

<b>1</b>	<b>Complex Arithmetic</b>	<b>3</b>
1.1	Complex Inner Product . . . . .	3
1.2	Complex Polar Form . . . . .	4
1.3	Roots of Unity . . . . .	5
1.4	Summing Roots of Unity . . . . .	6
1.5	Kronecker Delta . . . . .	9

# Chapter 1

## Complex Arithmetic

Complex numbers are written in the form

$$\mathbb{C} = \{a + bi \mid a \in \mathbb{R}, b \in \mathbb{R} \text{ and } i^2 = -1\}$$

$a$  is the called the real part

$b$  is the imaginary part

$a + bi$  together are called complex

### 1.1 Complex Inner Product

#### Real Inner Product

The inner product definition for real vectors states that

$$\vec{a} \cdot \vec{b} = \sum_i \alpha_i \beta_i = \alpha_T \beta; \alpha, \beta \in \mathbb{R}$$

Which naturally leads to

$$|\vec{a}| = \sqrt{\vec{a} \cdot \vec{a}} \geq 0, \in \mathbb{R}$$

$$\text{Ex. } \vec{v} = \begin{pmatrix} 3 \\ 2 \end{pmatrix}, |\vec{v}|^2 = \vec{v} \cdot \vec{v} = \begin{pmatrix} 3 \\ 2 \end{pmatrix} \cdot \begin{pmatrix} 3 \\ 2 \end{pmatrix} = 9 + 4 = 13 \Rightarrow |\vec{v}| = \sqrt{13}$$

However, for complex vectors,  $\vec{a} \cdot \vec{a}$  could be negative and leads to  $|\vec{a}| \in \mathbb{C}$ .

So, using the above inner product definition, for  $\vec{v} = \begin{pmatrix} 3i \\ -2 \end{pmatrix}$ ,

$$|\vec{v}|^2 = \vec{v} \cdot \vec{v} = 9(i^2) + 4 = -9 + 4 = -5 \therefore |\vec{v}| = \sqrt{-5}$$

However, the magnitude of a vector can't be a complex value!

#### Complex Conjugate

$$\text{For } \vec{\gamma} = \begin{pmatrix} a + ib \\ c + id \end{pmatrix}, \vec{\gamma}^* = \begin{pmatrix} a - ib \\ c - id \end{pmatrix}$$

Conjugation is distributive across sums and products.

$$(\alpha + \beta)^* = \alpha^* + \beta^*$$

$$(\alpha \times \beta)^* = \alpha^* \times \beta^*$$

### Complex Inner Product

Now, the inner product definition changes:

$$\vec{\alpha} \cdot \vec{\beta} = (\vec{\alpha}_T)^* \cdot \vec{\beta}$$

This implies

$$\vec{\alpha} \cdot \vec{\alpha} = \vec{\alpha}_T^* \vec{\alpha} \in \mathbb{R}^+$$

For the above problem,

$$|\vec{v}|^2 = \vec{v} \cdot \vec{v} = (\vec{v}_T)^* \vec{v} = \begin{pmatrix} -3i & -2 \end{pmatrix} \begin{pmatrix} 3i \\ -2 \end{pmatrix} = -9(i^2) + 4 = 13 \Rightarrow |\vec{v}| = \sqrt{13} \in \mathbb{R}^+$$

This definition gives non-negative real magnitude to all complex vectors because

$$(a + bi)(a - bi) = a^2 + b^2, \forall a, b \in \mathbb{R}$$

$\alpha_T^*$  is also called the Hermitian conjugate  $\equiv \alpha^\dagger$  (alpha-dagger).

**Exercise:** Proof that all n-dimensional complex vectors have real magnitude from the complex inner product definition

## 1.2 Complex Polar Form

Any complex number can be transformed from the form  $z = a + bi$  to the form  $z = r(\cos \theta + i \sin \theta)$ , where  $r = \sqrt{a^2 + b^2}$  &  $\theta = \tan^{-1} \left( \frac{b}{a} \right)$ .

From Euler's formula, this becomes

$$z = r e^{i\theta}$$

For example,

$$1 + i = \sqrt{2} \left( \cos \frac{\pi}{4} + i \sin \frac{\pi}{4} \right) = \sqrt{2} e^{i\frac{\pi}{4}}$$

### Some Results of Euler's Formula

For  $z = e^{i\theta}$  &  $w = e^{i\phi}$ ,

Multiplication:  $zw = rs e^{i\theta+\phi}$

Division:  $\frac{z}{w} = \frac{r}{s} e^{i\theta-\phi}$

Conjugation:  $z^* = re^{-i\theta}$

## 1.3 Roots of Unity

For  $z \in \mathbb{R}$  &  $n \in \mathbb{Z}$ , there are two solutions to the equation  $z^n = 1$  ( $z = \pm 1$ )

The number of solutions depends on the power  $n$ : 2 for even  $n$ , 1 for odd  $n$ .

However, when the domain of  $z$  is extended to complex numbers, there are  $n$  unique complex solutions for every  $n$ .

In the complex polar form,

$$z^n = e^{i\theta n} = 1$$

$$e^{i(n\theta)} = \cos(n\theta) + i \sin(n\theta) = 1 + 0i$$

We can solve either equations to get  $\theta$

$$\cos(n\theta) = 1 \text{ or } \sin(n\theta) = 0$$

Both leads to

$$\theta = \frac{2\pi k}{n}, k \in \mathbb{Z}_{\geq 0}$$

Notice that (**Excercise: Proof this**)

$$e^{i\frac{2\pi k}{n}} = e^{i\frac{2\pi(k+an)}{n}}$$

We can define arbitrary set  $\mathcal{R}_n$  to be the set of all  $k$ , where

$$f(k) \equiv e^{i\frac{2\pi}{n}k} \text{ is unique}$$

From above, we can deduce a periodicity in the range of this function, where

$$f(\mathcal{R}_n) = f(\mathcal{R}_n + an)$$

In fact,  $\mathcal{R}_n$  is a least residue system modulo  $n$ . This just means that we can use the properties of modular arithmetic on  $\mathcal{R}_n$ .

$$f(\mathcal{R}_n) = f(\mathcal{R}_n \bmod n)$$

Going back the roots of unity,

$$z = e^{i\frac{2\pi}{n}(k \bmod n)}, k \bmod n \in \{0, \dots, n-1\}$$

Thus,

$$z^n = 1$$

In some context,  $z$  is written as  $\omega_n$ , so in summary

$$(\omega_n)^k = \sqrt[n]{1} = \cos\left(\frac{2\pi k}{n}\right) + i \sin\left(\frac{2\pi k}{n}\right) = e^{i\frac{2\pi}{n}k}, k \in \{0, \dots, n-1\}$$

For example,  $\sqrt[5]{1} = 0.309 + 0.951i$  &  $-0.809 - 0.5878i$  & 3 other numbers

**Exercise:** Proof this


## 1.4 Summing Roots of Unity

The sum of all root  $n$ th root of unity is 0

$$\sum_{k=0}^{n-1} (\omega_n)^k = 0$$

### Geometric Proof

- To prove this geometrically, draw all the  $(\omega_n)^k$  as vectors with origin at 0 in the complex plane.
- Connect the end of the vectors together to end up with an  $n$ -sided regular.
- Notice that each vector is perpendicular to the opposite side of the polygon, so each can be seen as a scaled up and rotated version the polygon side.
- Because the differences of the angle between two adjacent vectors is  $\frac{2\pi}{n}$  (same as the polygon), when put end to end, the vectors will form a close loop that is another  $n$ -sided regular polygon.
- This means that the sum of the vector ends up at the origin and thus is 0



An  $n = 5$  example from SunilK

**Algebraic Proof**

Let

$$\sum_{k=0}^{n-1} (\omega_n)^k = A$$

Going back to set  $\mathcal{R}_n$  from above, define a function  $S$  for which

$$S(\mathcal{R}_n) = \sum_{k=0}^{n-1} \omega^k \quad (\omega \equiv \omega_n)$$

To repeat,  $\mathcal{R}_n$  is a cyclic group of order  $n$  with integer and modular addition. In other words,  $\mathcal{R}_n$  is the set integers modulo  $n$ . A property of  $\mathcal{R}_n$  is

$$\mathcal{R}_n = \mathcal{R}_n + \alpha \pmod{n}, \quad \alpha \in \mathbb{Z}$$

For example, let  $\mathcal{R}_n = \{0, 1, 2, 3\}$ , and  $\alpha = 2$

$$\mathcal{R}_n + \alpha = \{2, 3, 4, 5\} \pmod{4} = \{0, 1, 2, 3\}$$

This implies that

$$S(\mathcal{R}_n) = (\mathcal{R}_n + \alpha)$$

Thus

$$\sum_{k=0}^{n-1} \omega^k = \sum_{k=0}^{n-1} \omega^{k+\alpha} = \omega^\alpha \sum_{k=0}^{n-1} \omega^k$$

Substituting from above

$$\omega^\alpha \times A = A$$

The trivial solution is  $\omega^\alpha = 1$ . However, we're trying to solve  $A$

$$(\omega^\alpha - 1)A = 0 \quad \therefore A = 0$$

In conclusion this result agrees with the geometric proof,

$$\sum_{k=0}^{n-1} \omega^k = 0$$

**A More General Sum**

A broader definition for the function  $S$  from above is  $S_a$ ,

$$S_a(\mathcal{R}_n) = S(a \cdot \mathcal{R}_n \pmod n) = \sum_{k=0}^{n-1} (\omega^k)^a, \quad a \in \mathbb{Z}_{\geq 0, < n}$$

This is when every element in the  $\mathcal{R}_n$  is mod- $n$  multiplied by a constant  $a$ . The sum of roots of unity above,  $S(\mathcal{R}_n)$ , is the special case where  $a = 1$ .

Right off the bat, for the case  $a = 0$ , the sum evaluates to  $n$ .


Evaluating the sum for  $a > 1$  requires making some assumptions.

### Conjecture 1

The operation  $a \cdot \mathcal{R}_n \pmod n$  collapses the cyclic group  $\mathcal{R}_n$  uniformly into a smaller cyclic group that is a factor of  $n$ .

$$a \cdot \mathcal{R}_n \pmod n \rightarrow \mathcal{R}_\phi, \quad \phi = \frac{n}{\gcd(n, a)}$$

The digram shows the geometric representation of the  $n = 6, a = 2$  example, where the hexagon is collapsed into a triangle.



### An $n = 6, a = 2$ example

The left figure shows the mapping of points by  $S_a$ .

$$2 \cdot \{0, 1, 2, 3, 4, 5, 6\} \pmod n = \{0, 2, 4, 0, 2, 4\} = 2 \text{ sets of } \{0, 2, 4\}$$

The right figure shows the end-result of the mapping. The number on top of the vertex is the number of points mapped into that coordinate.

Running a Monte Carlo simulation of 100,000 shots on this conjecture on Python 3.6 & 3.7 confirms this result 100% for  $n < 100,000$ .

Assuming Conjecture 1 is true, this implies that



$a \cdot \mathcal{R}_n \pmod n \rightarrow \left(\frac{n}{\phi}\right)$  repetitions of  $\mathcal{R}_\phi \rightarrow \left(\frac{n}{\phi}\right)$  identical sets of  $\mathcal{R}_\phi$

This means that

$$S_a(\mathcal{R}_n) = S(a \cdot \mathcal{R}_n \pmod n) = S\left(\frac{n}{\phi} \cdot \mathcal{R}_\phi\right)$$

Because  $S_a$  is a sum (linear operator), we can bring the  $\frac{n}{\phi}$  outside of the sum.

$$S\left(\frac{n}{\phi} \cdot \mathcal{R}_\phi\right) = \frac{n}{\phi} \cdot S(\mathcal{R}_\phi) = \frac{n}{\phi} \cdot \sum_{k=0}^{\phi-1} \omega^k$$

We have already prove that the sum of roots of unity equals 0. This means that our sum also equals 0. In summary,

$$S_a(\mathcal{R}_n) = \sum_{k=0}^{n-1} (\omega^k)^a = \begin{cases} n, & a = 0 \\ 0, & 0 < a < n \end{cases}$$

## 1.5 Kronecker Delta

The Kronecker delta is a useful notation to expresses a result to be 0 for all index, except index m, where the the result is 1.

$$\delta_{jm} = \begin{cases} 1, & \text{if } j = m \\ 0, & \text{otherwise} \end{cases}$$

Expressing the general sum above using the Kronecker delta,

$$\sum_{k=0}^{n-1} (\omega^k)^a = \sum_{k=0}^{n-1} \omega^{(j-m)k} = n \cdot \delta_{jm}$$