H2 Math Summary

Notes

Maximus Pung

Contents

1 Systems of Linear Equations 2

1.1 Forming the Equations . . . 2
1.2 Special Case: Number of Equations > Number of Variables . . . 2
1.3 Special Case: Number of Equations < Number of Variables . . . 3

2 Inequalities 4

2.1 Solving Inequalities Graphically . . . 4
2.2 Sign Test Technique . . . 5
2.3 Common Substitutions . . . 6

3 Graphing Techniques 7

3.1 Key Features of a Curve . . . 7
3.2 Graphs of Rational Functions . . . 10
3.3 Symmetry . . . 11
3.4 Basic Graphs . . . 11
3.5 Parametric Equations . . . 12
3.5.1 Finding the Cartesian Equation . . . 12
3.5.2 Asymptotes (Parametric Equations) . . . 13
3.5.3 Intersections . . . 13
3.6 Conics . . . 14

4 Graphing Transformations 16

4.1 Translations and Scalings . . . 16
4.2 Reflections . . . 17
4.3 Combining Transformations . . . 17
4.4 Modulus Graphs . . . 18
4.5 Reciprocal Graphs $y = \frac{1}{\mathrm{f}(x)}$ . . . 20
4.6 Derivative Graphs $y = \mathrm{f}'(x)$ . . . 20
4.7 Reverse Transformations . . . 22
4.8 Transformations of Conics . . . 22

5 Functions 23

5.1 Functions, Domain and Range . . . 23
5.2 Inverse Functions . . . 23
5.2.1 Removing the Modulus . . . 24
5.3 Composite Functions . . . 25
5.3.1 Composite with a Piecewise Function . . . 25
5.4 Functional Powers and Self-Inverse Functions . . . 25
5.5 Piecewise and Periodic Functions . . . 26
5.5.1 Inverse of a Piecewise Function . . . 27

1 Systems of Linear Equations

1.1 Forming the Equations

We read the context of the question and form equations depending on the context of the question.

Example 1.1

$$\left.\begin{aligned} a+b+c&=176 && —(1)\\ 0.85a+0.90b+0.95c&=161.80 && —(2)\\ 0.93a+0.95b+0.98c&=169.56 && —(3)\end{aligned}\right\} \Rightarrow a=32,\ b=44,\ c=100$$

Example 1.2

The "variables" may be expressions. To find x, y and z, use the given information to form the equations:

$$\left.\begin{aligned} 2x^3+3\ln y-\mathrm{e}^z&=18 && —(1)\\ x^3-\ln y+2\mathrm{e}^z&=9 && —(2)\\ x^3+2\ln y+\mathrm{e}^z&=11 && —(3)\end{aligned}\right\} \Rightarrow x^3=8,\ \ln y=1,\ \mathrm{e}^z=1$$

It is technically still 'linear' because we can think of $A=x^3$, $B=\ln y$, $C=\mathrm{e}^z$, which turns it into a linear system in A, B and C. A truly nonlinear equation could look like $x^2+x+y^2+2y-3z^3-xz+xy=1$; it cannot be made linear this way, because x appears as both x^2 and x, and there are product terms such as xz and xy.

1.2 Special Case: Number of Equations > Number of Variables

Sometimes the number of equations is more than the number of variables, and the solution must satisfy all of the equations.

Example 1.3

$$\left.\begin{aligned} \lambda+\mu&=1 && —(1)\\ 2\lambda-\mu&=5 && —(2)\end{aligned}\right\} \Rightarrow \lambda=2,\ \mu=-1$$

$$\lambda+3\mu=-1 \qquad —(3)$$

Sometimes there may be no solution, e.g. skew lines in vectors. In this case, solve for λ and μ using equations (1) and (2). Then sub into (3).

> **Note**
>
> Check (3): if LHS = RHS, the equations are consistent; if LHS ≠ RHS, there is no solution.

1.3 Special Case: Number of Equations < Number of Variables

Sometimes the number of equations is less than the number of variables, so there are infinitely many solutions.

Example 1.4

$$\left.\begin{aligned} x+y+z=6 \quad &—(1) \\ x-y+2z=5 \quad &—(2) \end{aligned}\right\} \Rightarrow x=\tfrac{11}{2}-\tfrac{3}{2}z,\ y=\tfrac{1}{2}+\tfrac{1}{2}z,\ z=z$$

> **Note**
>
> Useful trick: in contextual questions, the variables are usually non-negative integers, $z \in \mathbb{Z}_0^+$, so use this to find the solution from the general solution.

When $z=z$, it means that there are infinitely many solutions, and z can be any real value. So just sub that into x and y to get a correct solution.

$z=-1: (7,0,-1)$ $z=0: (\tfrac{11}{2},\tfrac{1}{2},0)$ $z=1: (4,1,1)$ $z=2: (\tfrac{5}{2},\tfrac{3}{2},2)$

$z=3: (1,2,3)$ $z=4: (-\tfrac{1}{2},\tfrac{5}{2},4)$ $z=5: (-2,3,5)$ $z=6: (-\tfrac{7}{2},\tfrac{7}{2},6)$

We can use the GC table to find all these solutions.

2 Inequalities


2.1 Solving Inequalities Graphically

Some inequalities cannot be solved algebraically, so we sketch the graphs and read off the solution.

- Rearrange to $f(x) > 0$, or compare $y = f(x)$ with $y = g(x)$.
- Sketch on the GC; mark the intersections and the asymptotes.
- Read off the x-values where the required graph is above or below.

Example 2.1

$$|x^2 - 4| \leqslant \ln(x+3)$$



Using GC, intersections at $x = 1.5746$, $x = 2.3840$

$$\therefore x = -2 \text{ or } 1.57 \leqslant x \leqslant 2.38$$


2.2 Sign Test Technique

We solve rational inequalities by finding where the expression changes sign.

- Make the right-hand side 0.
- Do **not** cross multiply unless the expression or bracket is always positive.
- Expressions that are always positive can be divided across, e.g. a quadratic with discriminant <0 and positive leading coefficient: $x^2+2x+5=(x+1)^2+4>0$.

Example 2.2

$$\frac{(x-1)(x-2)^2}{(x-3)(x-4)} \leqslant 0$$



$x=2$: numerator $=0$ (included)

$x=3,\ x=4$: denominator $=0$ (excluded)

$\therefore x \leqslant 1$ or $x=2$ or $3<x<4$

You can sketch the graph to check the answer.

> **Note**
>
> Crossing a factor with an odd power changes the sign; crossing a factor with an even power keeps the sign the same.

2.3 Common Substitutions

Once an inequality in x is solved, we can replace x with an expression to deduce the solution of a related inequality.


Replace x by	**Original inequality**		
	$x>2$	$x<-1$	$-1<x<2$
e^x	$x>\ln 2$	no solution	$x<\ln 2$
$\ln x$	$x>e^2$	$0<x<e^{-1}$	$e^{-1}<x<e^2$
$-x$	$x<-2$	$x>1$	$-2<x<1$
$x+1$	$x>1$	$x<-2$	$-2<x<1$
$\lvert x\rvert$	$x<-2$ or $x>2$	no solution	$-2<x<2$
x^2	$x<-\sqrt{2}$ or $x>\sqrt{2}$	no solution	$-\sqrt{2}<x<\sqrt{2}$
$\frac{1}{x}$	$\left(0<x<\frac{1}{2}\right)$	$(-1<x<0)$	$\left(x<-1 \text{ or } x>\frac{1}{2}\right)$
$\cos x,\ 0\leqslant x\leqslant 2\pi$	(no solution)	(no solution)	$(0\leqslant x<\pi \text{ or } \pi<x\leqslant 2\pi)$

Regardless of whatever substitution you use, using a graph will always work, but some can be solved without a graph.

For $\frac{1}{x}$, $\sin x$ and $\cos x$, draw the graph to solve.


Example 2.3

$$\frac{1}{x}>2, \quad \frac{1}{x}<-1, \quad -1<\frac{1}{x}<2$$



Example 2.4

$$\cos x>\tfrac{1}{2}, \quad \cos x<-\tfrac{1}{2},$$
$$-\tfrac{1}{2}<\cos x<\tfrac{1}{2}, \quad 0\leqslant x\leqslant 2\pi$$




3 Graphing Techniques


3.1 Key Features of a Curve

Key features of a curve:

(a) **x-intercepts:** let $y=0$.
e.g. $y=x^2-1$: $x^2-1=0 \Rightarrow x=\pm 1$.




y-intercepts: let $x=0$.
e.g. $y=x+2$: $y=2$.




(b) **Stationary points**, when $\frac{dy}{dx}=0$ or $\frac{dy}{dx}$ is undefined. Label upon request.

Horizontal turning point: $\frac{dy}{dx}=0$ (tangent horizontal).
e.g. $y=-(x-1)^2+2$, maximum point $(1,2)$.






Vertical turning point: $\frac{dy}{dx}$ undefined (tangent vertical).
e.g. $x=(y-1)^2+1$, turning point $(1,1)$.



(c) Asymptotes


(i) **Vertical asymptotes.** If as $x \to k$, $y \to \pm\infty$, then $x = k$ is a vertical asymptote.

Common Occurrence	**Condition**	**Example and asymptote**	**Graph**
Fractions	denominator $= 0$	$y = \dfrac{1}{x-1}$ $x - 1 = 0 \Rightarrow x = 1$ $\therefore x = 1$ is a vertical asymptote	
Logarithmic graphs	inside ln $= 0$	$y = \ln(x-2)$ $x - 2 = 0 \Rightarrow x = 2$ $\therefore x = 2$ is a vertical asymptote	
Tangent graphs	inside tan $= \dfrac{\pi}{2} + k\pi$, $k \in \mathbb{Z}$	$y = \tan x$ $x = \pm\dfrac{\pi}{2}, \pm\dfrac{3\pi}{2}, \ldots$	


(ii) **Horizontal asymptotes.** As $x \to \pm\infty$, if $y \to k$, then $y = k$ is a horizontal asymptote.

- To find horizontal asymptotes, one should sub in $x \to \pm\infty$ into the equation. This is the only foolproof way to do it.
- A curve **CAN** intersect its horizontal asymptote.

Note: there are more cases of horizontal asymptotes, but these are the common cases.






e.g. $y = \dfrac{\sin x}{x}$: curve crosses $y = 0$



e.g. $y = 3 + e^{-x}$


Useful trick to find horizontal asymptotes

For rational functions $\frac{P(x)}{Q(x)}$:


Case	Horizontal asymptote	Example	Graph
numerator power > denominator power	no horizontal asymptote	$y=\frac{x^2}{x-1}$	
numerator power < denominator power	$y=0$	$y=\frac{1}{x^2+1}$	
numerator power = denominator power	ratio of coefficients of highest power	$y=\frac{2x+1}{x-1}$ $y=2$	

(iii) **Oblique asymptotes.** An asymptote of the form $y=mx+c$, $m\neq 0$.

Graphs with oblique asymptotes come in the form $y=mx+c+\mathrm{g}(x)$, where $\mathrm{g}(x)\to 0$ as $x\to\pm\infty$; the oblique asymptote is $y=mx+c$.



e.g. $y=2x+1+\frac{3}{x-2}$









e.g. $y=3x-5+\mathrm{e}^{-x}$

Note

For parametric curves, oblique asymptotes (if they exist) can only be found via the cartesian equation.

3.2 Graphs of Rational Functions


Two common shapes for each type.

Type and form	Shape 1	Shape 2
Linear / linear $y=\dfrac{Ax+B}{Cx+D}$ Horizontal asymptote: $y=\dfrac{A}{C}$ Vertical asymptote: $x=-\dfrac{D}{C}$	 $y=\dfrac{x+1}{x-1}$	 $y=\dfrac{x-1}{x+1}$
Quadratic / linear $y=\dfrac{ax^2+bx+c}{dx+e}$ $=Ax+B+\dfrac{C}{dx+e}$ Oblique asymptote: $y=Ax+B$ Vertical asymptote: $x=-\dfrac{e}{d}$	 2 turning points $y=x+1+\dfrac{1}{x-1}\quad\left(\dfrac{Cd}{A}>0\right)$	 No turning points $y=x+1-\dfrac{1}{x-1}\quad\left(\dfrac{Cd}{A}<0\right)$
Linear / quadratic $y=\dfrac{ax+b}{cx^2+dx+e}$ Horizontal asymptote: $y=0$ Vertical asymptotes: solve $cx^2+dx+e=0$ (possibly no roots)	 $y=\dfrac{x}{(x-1)(x+1)}$	 $y=\dfrac{x-2}{(x-1)(x+1)}$


3.3 Symmetry

Even functions: $f(-x)=f(x)$.

The graph is symmetric about the y-axis (line of symmetry $x=0$, dotted).



e.g. $y=x^2$




e.g. $y=\cos x$

Odd functions: $f(-x)=-f(x)$.


The graph is symmetric about the origin.

e.g. $y=x^3-x$















Symmetry about $y=x$: swapping x and y gives the same equation.

e.g. $y=\dfrac{1}{x}$ $(xy=1)$



3.4 Basic Graphs

 Linear $y=x+1$	 Quadratic $y=x^2-1$	 Cubic $y=x^3-x$
 Exponential $y=e^x$	 Logarithm $y=\ln x$	 Reciprocal $y=\frac{1}{x}$
 $y=\sin x$	 $y=\cos x$	 $y=\tan x$
 $y=\sin^{-1}x$	 $y=\cos^{-1}x$	 $y=\tan^{-1}x$

3.5 Parametric Equations

A curve C can be described by giving both coordinates in terms of a parameter t:

$$x=x(t), \quad y=y(t), \quad t\in T$$


Example.

$x=2\cos t+1, \quad y=3\sin t-2, \quad 0\leqslant t\leqslant \pi$

$t=0$: $(3,-2)$ $\quad t=\frac{\pi}{2}$: $(1,1)$ $\quad t=\pi$: $(-1,-2)$

$x=0$: $\left(0, \frac{3\sqrt{3}}{2}-2\right)$

$y=0$: $\left(1\pm\frac{2\sqrt{5}}{3}, 0\right)$



CAUTION: sketching from parametric equations

ALWAYS take note of the range of t when sketching graphs. The restriction on t decides which part of the curve exists: above, $0\leqslant t\leqslant\pi$ gives only the upper half of the ellipse, with endpoints $(3,-2)$ and $(-1,-2)$, not the whole ellipse.

3.5.1 Finding the Cartesian Equation

The cartesian equation of C is an equation without the parameter t. Common techniques:

① **Make t the subject.** e.g. $x=2t-1, y=3t^2+e^t, t\in\mathbb{R}$

$$t=\frac{x+1}{2}$$

$$y=3\left(\frac{x+1}{2}\right)^2+e^{\frac{x+1}{2}}$$

② **Trigo identity formulas.** e.g. $x=2\cos t+1, y=3\sin t-2, 0\leqslant t\leqslant\pi$

$$\cos t=\frac{x-1}{2}, \quad \sin t=\frac{y+2}{3}$$

$$\cos^2 t+\sin^2 t=1$$

$$\frac{(x-1)^2}{4}+\frac{(y+2)^2}{9}=1, \quad y\geqslant -2$$

③ **Others.** e.g. $x=t-\frac{1}{t}, y=t+\frac{1}{t}, t\neq 0$

$$y+x=2t, \quad y-x=\frac{2}{t}$$

$$(y+x)(y-x)=4$$

$$y^2-x^2=4$$

Note: Avoid finding the cartesian eqn if the question did not ask you to.

3.5.2 Asymptotes (Parametric Equations)

- Find (if possible) $t \to t_0$ such that $x \to x_0, y \to \pm\infty$ (vertical asymptote $x = x_0$), or $y \to y_0, x \to \pm\infty$ (horizontal asymptote $y = y_0$).
- Oblique asymptotes (if they exist) can only be found via the cartesian equation.


Example. $x = \dfrac{1}{t-1},\ y = 2t,\ t \neq 1$

As $t \to \pm\infty,\ x \to 0,\ y \to \pm\infty$

$\therefore x = 0$ is a vertical asymptote

As $t \to 1,\ x \to \pm\infty,\ y \to 2$

$\therefore y = 2$ is a horizontal asymptote




Oblique asymptotes. Cannot be found from the parametric form alone; find the cartesian equation first.

e.g. $x = 2\sec\theta,\ y = 3\tan\theta$

$$\sec^2\theta - \tan^2\theta = 1$$

$$\frac{x^2}{4} - \frac{y^2}{9} = 1$$

Oblique asymptotes: $y = \pm\dfrac{3}{2}x$



3.5.3 Intersections

Intersection between a parametric curve and another curve: sometimes both equations are needed.

e.g. $x = 2t + \ln t,\ y = t^2 - 1,\ t > 0$ and the circle $x^2 + y^2 = 25$:

$$(2t + \ln t)^2 + (t^2 - 1)^2 = 25$$

Using GC, $t = 0.00735$ or $t = 1.86$ (both > 0)

$\therefore$ intersections: $(-4.90,\ -1.00)$ and $(4.35,\ 2.47)$

This equation in t cannot be solved algebraically, so the GC is needed.


3.6 Conics

Circle

$$(x-h)^2+(y-k)^2=r^2$$

Centre: (h,k)

Radius: r




Parabola

$$y-k=a(x-h)^2 \qquad x-h=b(y-k)^2$$

$y-k=a(x-h)^2$:
turning point (h,k), $\dfrac{dy}{dx}=0$

$x-h=b(y-k)^2$:
turning point (h,k), $\dfrac{dy}{dx}$ undefined



Ellipse


$$\frac{(x-h)^2}{a^2}+\frac{(y-k)^2}{b^2}=1$$

Centre: (h,k)

"Horizontal radius" $=a$, "vertical radius" $=b$

For $a>b$: major axis horizontal (length $2a$, semi-major a); minor axis vertical (length $2b$, semi-minor b).

⋆ RHS ⇒ numerator coefficients of x^2, y^2 must all be 1.




Hyperbola: Type I

$$\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$$

Centre: (h, k)

Distance from centre to vertices $= a$

Asymptotes (dashed): $y - k = \pm\frac{b}{a}(x-h)$



- RHS $\Rightarrow$ numerator coefficients of x^2, y^2 must all be 1.
- Equation of asymptotes: $y - k = \pm\frac{b}{a}(x-h)$


Hyperbola: Type II

$$\frac{(y-k)^2}{b^2} - \frac{(x-h)^2}{a^2} = 1$$

Centre: (h, k)

Distance from centre to vertices $= b$

Asymptotes (dashed): $y - k = \pm\frac{b}{a}(x-h)$



- RHS $\Rightarrow$ numerator coefficients of x^2, y^2 must all be 1.
- Equation of asymptotes: $y - k = \pm\frac{b}{a}(x-h)$





---

4 Graphing Transformations

Every transformation comes from **replacing** x **or** y in the equation $y=\mathrm{f}(x)$.

Throughout, blue is $y=\mathrm{f}(x)=x(x-2)$, with intercepts $(0,0)$, $(2,0)$ and minimum point $(1,-1)$; red is the transformed graph.



4.1 Translations and Scalings

Transformation	Replacement	Result	Graph
Translation of a units in the positive y direction	replace y with $y-a$: $y-a=\mathrm{f}(x)$	$y=\mathrm{f}(x)+a$ (graph: $a=2$)	
Translation of a units in the positive x direction	replace x with $x-a$	$y=\mathrm{f}(x-a)$ (graph: $a=2$)	
Scaling by factor a parallel to the y-axis	replace y with $\frac{y}{a}$: $\frac{y}{a}=\mathrm{f}(x)$	$y=a\mathrm{f}(x)$ (graph: $a=2$)	
Scaling by factor a parallel to the x-axis	replace x with $\frac{x}{a}$	$y=\mathrm{f}\left(\frac{x}{a}\right)$ (graph: $a=2$)	

CAUTION

Every replacement acts the "opposite" way to what it looks like: $x-2$ moves the graph in the **positive** x direction; $y=\mathrm{f}(2x)$ is a scaling by factor $\frac{1}{2}$ (not 2) parallel to the x-axis.
Replacing x does **not** mean replacing the input of f, and replacing y does not mean replacing the output $\mathrm{f}(x)$.

4.2 Reflections

Transformation	Replacement	Result	Graph
Reflection about the x-axis	replace y with $-y$: $-y=\mathrm{f}(x)$	$y=-\mathrm{f}(x)$	
Reflection about the y-axis	replace x with $-x$	$y=\mathrm{f}(-x)$	

4.3 Combining Transformations


Functions of the form $y=a\mathrm{f}(bx+c)+d,\ \ a,b,c,d\in\mathbb{R}$: replace x and y **one step at a time**.

e.g. $y=\mathrm{f}(x)\ \longrightarrow\ y=2\mathrm{f}(2x-1)+1$

Replacement	Transformation	Function
replace x with $x-1$	translation of 1 unit in the positive x direction	$y=\mathrm{f}(x-1)$
replace x with $2x$	scaling by factor $\frac{1}{2}$ parallel to the x-axis	$y=\mathrm{f}(2x-1)$
replace y with $\frac{y}{2}$	scaling by factor 2 parallel to the y-axis	$y=2\mathrm{f}(2x-1)$
replace y with $y-1$	translation of 1 unit in the positive y direction	$y=2\mathrm{f}(2x-1)+1$

For $f(x)=x(x-2)$:

$$(1,-1)\to(2,-1)\to(1,-1)$$
$$\to(1,-2)\to(1,-1)$$
$$(0,0)\to(1,0)\to(\tfrac{1}{2},0)$$
$$\to(\tfrac{1}{2},0)\to(\tfrac{1}{2},1)$$





CAUTION: order of transformations

Inside f: do the translation **before** the scaling/reflection in x. $y=f(2x-1)$ is "translate 1 unit, then scale by $\frac{1}{2}$", **not** "scale by $\frac{1}{2}$, then translate 1 unit" (that gives $y=f(2(x-1))=f(2x-2)$).
Outside f: do the scaling before the translation in y.

4.4 Modulus Graphs

Every modulus graph can be partitioned into a "+" part and a "−" part: $|x|=x$ for $x\geqslant 0$, $|x|=-x$ for $x<0$.

Graph	How to sketch	Graph
$y=\|f(x)\|$	Keep the part on or above the x-axis; reflect the part below the x-axis about the x-axis. Sharp points at the x-intercepts.	
$y=f(\|x\|)$	Remove the part on the left of the y-axis; reflect the part on the right of the y-axis about the y-axis.	

e.g. $y=f(x)\longrightarrow y=|f(-2x+1)|$

Replacement	Transformation	Function
replace x with $x+1$	translation of 1 unit in the negative x direction	$y=f(x+1)$
replace x with $-x$	reflection about the y-axis	$y=f(-x+1)$
replace x with $2x$	scaling by factor $\frac{1}{2}$ parallel to the x-axis	$y=f(-2x+1)$
$\|f\|$	reflect parts below the x-axis about the x-axis	$y=\|f(-2x+1)\|$

CAUTION

$y=\mathrm{f}(|x|+1)$ and $y=\mathrm{f}(|x+1|)$ are **different** curves.

$\mathrm{f}(|x|+1)$: translate first, then apply $|x|$. $\mathrm{f}(|x+1|)$: apply $|x|$ first, then translate.

4.5 Reciprocal Graphs $y=\frac{1}{\mathrm{f}(x)}$

Analyse the new curve at the critical points of $y=\mathrm{f}(x)$.


$y=\mathrm{f}(x)$	$y=\frac{1}{\mathrm{f}(x)}$
x-intercept $(a,0)$	vertical asymptote $x=a$
vertical asymptote $x=a$	$y\to 0$: point $(a,0)$ (excluded)
maximum point (p,q), $q\neq 0$	minimum point $\left(p,\frac{1}{q}\right)$
minimum point (p,q), $q\neq 0$	maximum point $\left(p,\frac{1}{q}\right)$
horizontal asymptote $y=k$, $k\neq 0$	horizontal asymptote $y=\frac{1}{k}$
$y\to\pm\infty$ (incl. oblique asymptote)	horizontal asymptote $y=0$
$y=\pm 1$	same point $(\frac{1}{\pm 1}=\pm 1)$
$\mathrm{f}(x)>0$ / $\mathrm{f}(x)<0$	$\frac{1}{\mathrm{f}(x)}>0$ / $\frac{1}{\mathrm{f}(x)}<0$ (same sign)

e.g. $\mathrm{f}(x)=x^2-4$:

$(-2,0),\ (2,0)\to x=\pm 2$

$(0,-4)\to\left(0,-\frac{1}{4}\right)$

$y\to\infty\to y=0$



4.6 Derivative Graphs $y=\mathrm{f}'(x)$

$\mathrm{f}'(x)=\frac{\mathrm{d}y}{\mathrm{d}x}$ is the gradient of $y=\mathrm{f}(x)$: read off the gradient on each interval.


$y=\mathrm{f}(x)$	$y=\mathrm{f}'(x)$
stationary point at $x=a$	x-intercept $(a,0)$
increasing / decreasing	above / below the x-axis
point of inflexion (max. or min. gradient)	turning point
vertical asymptote $x=a$	vertical asymptote $x=a$
horizontal asymptote $y=k$	horizontal asymptote $y=0$
oblique asymptote $y=mx+c$	horizontal asymptote $y=m$

e.g. $\mathrm{f}(x) = x^3 - 3x$:

$$\mathrm{f}'(x) = 3x^2 - 3$$

SP at $x = \pm 1 \rightarrow (\pm 1, 0)$

inflexion at $x = 0 \rightarrow$ min. $(0, -3)$



CAUTION

Mark x-coordinates only where they come from $y = \mathrm{f}(x)$: the y-values of $y = \mathrm{f}'(x)$ are generally **not** known from the graph of $y = \mathrm{f}(x)$.

4.7 Reverse Transformations

If $y=\mathrm{f}(x)$ undergoes a sequence of transformations to give a known $y=\mathrm{g}(x)$, undo the transformations **in reverse order**, each with its opposite replacement.

e.g. translate 2 units in the negative x direction, reflect about the y-axis, scale by factor 3 parallel to the y-axis; result $y=6\mathrm{e}^{-x+3}$.

Opposite transformation	**Replacement**	**Function**
scale by factor $\frac{1}{3}$ parallel to the y-axis	replace y with $3y$	$y=2\mathrm{e}^{-x+3}$
reflect about the y-axis	replace x with $-x$	$y=2\mathrm{e}^{x+3}$
translate 2 units in the positive x direction	replace x with $x-2$	$y=2\mathrm{e}^{x+1}$

$$\therefore \mathrm{f}(x)=2\mathrm{e}^{x+1}$$

4.8 Transformations of Conics


Apply exactly the same replacements of x and y.

e.g. $x^2+y^2=1$: scale by factor 2 parallel to the x-axis, translate 2 units in the positive x direction, translate 1 unit in the positive y direction.

Transformation	**Replacement**	**Equation**
scale by factor 2 parallel to the x-axis	replace x with $\frac{x}{2}$	$\frac{x^2}{4}+y^2=1$
translate 2 units in the positive x direction	replace x with $x-2$	$\frac{(x-2)^2}{4}+y^2=1$
translate 1 unit in the positive y direction	replace y with $y-1$	$\frac{(x-2)^2}{4}+(y-1)^2=1$

$$\therefore \frac{(x-2)^2}{4}+(y-1)^2=1$$

Centre $(2,1)$.



CAUTION

Replace x by $(x-2)$ **inside** the bracket: $\frac{(x-2)^2}{4}$, not $\frac{x^2}{4}-2$ or $\frac{x^2-2}{4}$.

5 Functions

5.1 Functions, Domain and Range

$$\mathrm{f}: x \mapsto \underbrace{x^2+3x+2}_{\text{rule f}(x)}, \quad x \in \mathbb{R}, \quad \underbrace{x \geqslant 0}_{\text{domain } D_\mathrm{f}}$$

A function is a relation that gives **exactly one** value of y for each $x \in D_\mathrm{f}$.

- **Is a function:** every vertical line $x=a$, $a \in D_\mathrm{f}$, cuts $y=\mathrm{f}(x)$ **exactly once**.
- **Not a function:** find **one** vertical line $x=a$, $a \in D_\mathrm{f}$, that cuts the graph more than once, or not at all.
- Two functions are equal only if they have the same rule **and** the same domain.

Inequality	Set notation	Interval
$a<x<b$	$\{x \in \mathbb{R} : a<x<b\}$	(a, b)
$a \leqslant x \leqslant b$	$\{x \in \mathbb{R} : a \leqslant x \leqslant b\}$	$[a, b]$
$a \leqslant x<b$	$\{x \in \mathbb{R} : a \leqslant x<b\}$	$[a, b)$
$x>a$	$\{x \in \mathbb{R} : x>a\}$	(a, ∞)
$x \in \mathbb{R},\ x \neq 0$	$\mathbb{R} \setminus \{0\}$	$(-\infty, 0) \cup (0, \infty)$

CAUTION


To find the range of a function, **ALWAYS** sketch its graph $y=\mathrm{f}(x)$, $x \in D_\mathrm{f}$, and read the range off the graph. **Never** just sub in the endpoints of the domain.

5.2 Inverse Functions

	f^{-1}	**Working**
Exists	f is one-to-one: every line $y=k$, $k \in \mathbb{R}$, cuts $y=\mathrm{f}(x)$ **at most once**	Does not exist: one line $y=l$ cuts more than once, or $\mathrm{f}(x_1)=\mathrm{f}(x_2)$ with $x_1 \neq x_2$
Rule	make x the subject of $y=\mathrm{f}(x)$	then swap: $\mathrm{f}^{-1}(x)=\ldots$
Domain	$D_{\mathrm{f}^{-1}}=R_\mathrm{f}$	sketch $y=\mathrm{f}(x)$ for R_f
Range	$R_{\mathrm{f}^{-1}}=D_\mathrm{f}$	
Graph	reflection of $y=\mathrm{f}(x)$ about $y=x$	all coordinates swapped

e.g. $\mathrm{f}: x \mapsto (x-3)^2 - 7,\ x \geqslant 3$

$$y = (x-3)^2 - 7$$
$$x - 3 = \sqrt{y+7} \quad (x \geqslant 3)$$
$$x = 3 + \sqrt{y+7}$$
$$R_\mathrm{f} = [-7, \infty)$$
$$\therefore \mathrm{f}^{-1}: x \mapsto 3 + \sqrt{x+7},\ x \in \mathbb{R},\ x \geqslant -7$$



CAUTION


- When drawing $y = \mathrm{f}(x)$ and $y = \mathrm{f}^{-1}(x)$ together, use the **same scale** on both axes.
- $\mathrm{ff}^{-1}(x) = \mathrm{f}^{-1}\mathrm{f}(x) = x$, but $D_{\mathrm{ff}^{-1}} = D_{\mathrm{f}^{-1}} = R_\mathrm{f}$ while $D_{\mathrm{f}^{-1}\mathrm{f}} = D_\mathrm{f}$.
- Solving $\mathrm{f}(x) = \mathrm{f}^{-1}(x)$: it is **not always** $\mathrm{f}(x) = x$ (only when the curves meet on $y = x$, e.g. f increasing).

5.2.1 Removing the Modulus

On a restricted domain, the sign inside the modulus is fixed, so the modulus can be removed: $|x-a| = x-a$ for $x \geqslant a$, $|x-a| = -(x-a)$ for $x < a$.

e.g. $\mathrm{f}: x \mapsto \dfrac{3}{|x-2|+1},\ x < 2$

$$x < 2: |x-2| = 2 - x$$
$$\mathrm{f}(x) = \frac{3}{3-x}$$
$$R_\mathrm{f} = (0, 3)$$
$$y = \frac{3}{3-x} \Rightarrow x = 3 - \frac{3}{y}$$
$$\therefore \mathrm{f}^{-1}: x \mapsto 3 - \frac{3}{x},\ x \in \mathbb{R},\ 0 < x < 3$$



Linear/linear functions with a modulus: remove the modulus using the domain first, then make x the subject. The domain of f^{-1} comes from the graph of $y = \mathrm{f}(x)$, not from the rule.

5.3 Composite Functions

	fg	**Working**
Exists	$R_g \subseteq D_f$	Does not exist: $x_0 \in R_g, x_0 \notin D_f$ $\therefore R_g \not\subseteq D_f$
Rule	sub $g(x)$ into $f(x)$: $f[g(x)]$	simplify
Domain	$D_{fg} = D_g$	rightmost function's domain (e.g. $D_{fgh} = D_h$)
Range	$D_g \xrightarrow{g} R_g \xrightarrow{f} R_{fg}$	sketch $y = f(x)$ on the domain R_g

e.g. $f: x \mapsto e^x, x \in \mathbb{R}$; $g: x \mapsto x^2+2, x > 0$

$$R_g = (2, \infty) \subseteq \mathbb{R} = D_f \therefore \text{fg exists}$$

$$fg(x) = e^{x^2+2}, \quad D_{fg} = (0, \infty)$$

$$(0, \infty) \xrightarrow{g} (2, \infty) \xrightarrow{f} (e^2, \infty) \therefore R_{fg} = (e^2, \infty)$$

5.3.1 Composite with a Piecewise Function

Split according to which piece of f the value $g(x)$ lands in.


e.g. $f(x) = \begin{cases} x^2, & 0 \leqslant x < 1 \\ 2-x, & 1 \leqslant x \leqslant 2 \end{cases}$ $\quad g(x) = \frac{x}{2}, \ 0 \leqslant x \leqslant 4$

$$R_g = [0, 2] \subseteq D_f \therefore \text{fg exists}$$

$$0 \leqslant \tfrac{x}{2} < 1 \iff 0 \leqslant x < 2$$

$$1 \leqslant \tfrac{x}{2} \leqslant 2 \iff 2 \leqslant x \leqslant 4$$

$$fg(x) = \begin{cases} \frac{x^2}{4}, & 0 \leqslant x < 2 \\ 2 - \frac{x}{2}, & 2 \leqslant x \leqslant 4 \end{cases} \qquad R_{fg} = [0, 1]$$



5.4 Functional Powers and Self-Inverse Functions


$f^2(x) = f[f(x)]$ exists if $R_f \subseteq D_f$; similarly f^3, f^4, … Note $f^2(x) \neq [f(x)]^2$.

Self-inverse: $f = f^{-1} \iff f^2(x) = x$ for all $x \in D_f$ (graph symmetric about $y = x$).

e.g. $\mathrm{g}: x \mapsto \dfrac{x+1}{x-1},\ x \neq 1$

$$\mathrm{g}^2(x) = \frac{\frac{x+1}{x-1}+1}{\frac{x+1}{x-1}-1} = \frac{2x}{2} = x$$

$$\therefore \mathrm{g}^{-1}(x) = \mathrm{g}(x)$$



Periodic composition pattern. If $\mathrm{f}^2(x) = x$:

$$\mathrm{f}^n(x) = \begin{cases} \mathrm{f}(x), & n \text{ odd} \\ x, & n \text{ even} \end{cases}$$

$$\mathrm{g}^{2027}(x) = \mathrm{g}(x) = \frac{x+1}{x-1}$$

$$\mathrm{g}^{2026}(2x) = 2x$$

$$\mathrm{g}^{-1}(2x) = \mathrm{g}(2x) = \frac{2x+1}{2x-1}$$

Similarly, if $\mathrm{f}^3(x) = x$, the pattern repeats every 3: $\mathrm{f}^n = \mathrm{f}^{n \bmod 3}$, and $\mathrm{f}^{-1} = \mathrm{f}^2$.

5.5 Piecewise and Periodic Functions


$$\mathrm{f}: x \mapsto \begin{cases} \mathrm{f}_1(x), & x \in D_1 \\ \mathrm{f}_2(x), & x \in D_2 \end{cases}, \qquad \mathrm{f}(x) = \mathrm{f}(x+p)$$

$\mathrm{f}(x) = \mathrm{f}(x+p)$: the graph repeats every p units (period p).

→ Always sketch the **original copy** first, then copy & paste.

→ Never sub any x value into f that is outside the domain of the original copy: shift by multiples of p first.

e.g. $\mathrm{f}(x) = \begin{cases} 1-x^2, & 0 \leqslant x < 1 \\ x-1, & 1 \leqslant x < 2 \end{cases},\ \mathrm{f}(x) = \mathrm{f}(x+2),\ -3 \leqslant x \leqslant 4.5$



$$\mathrm{f}(4.5) = \mathrm{f}(4.5-4) = \mathrm{f}(0.5) = 1 - 0.5^2 = 0.75$$

Area under a periodic function: area over n full periods $= n\times$ area over one period.

$$\int_0^2 \mathrm{f}(x)\,\mathrm{d}x = \int_0^1 (1-x^2)\,\mathrm{d}x + \int_1^2 (x-1)\,\mathrm{d}x = \frac{2}{3}+\frac{1}{2}=\frac{7}{6}$$

$$\int_{-2}^4 \mathrm{f}(x)\,\mathrm{d}x = 3\times\frac{7}{6}=\frac{7}{2}$$

5.5.1 Inverse of a Piecewise Function

Check one-to-one using the ranges of the pieces (they must not overlap and each piece is one-to-one); then invert each piece, with domain = range of that piece.

e.g. $\mathrm{h}(x)=\begin{cases} x^2, & 0\leqslant x\leqslant 1 \\ 2x-1, & 1<x\leqslant 3 \end{cases}$

ranges: $[0,1]$ and $(1,5]$ (no overlap)

$$\mathrm{h}^{-1}(x)=\begin{cases} \sqrt{x}, & 0\leqslant x\leqslant 1 \\ \dfrac{x+1}{2}, & 1<x\leqslant 5 \end{cases}$$

