H2 Math Summary

Notes

Maximus Pung

Contents

1 Systems of Linear Equations 2

1.1 Forming the Equations . . . 2
1.2 Special Case: Number of Equations > Number of Variables . . . 2
1.3 Special Case: Number of Equations < Number of Variables . . . 3

2 Inequalities 4

2.1 Solving Inequalities Graphically . . . 4
2.2 Sign Test Technique . . . 5
2.3 Common Substitutions . . . 6

3 Graphing Techniques 7

3.1 Key Features of a Curve . . . 7
3.2 Asymptotes . . . 7
3.2.1 Vertical Asymptotes . . . 7
3.2.2 Horizontal Asymptotes . . . 8
3.2.3 Oblique Asymptotes . . . 9
3.3 Graphs of Rational Functions . . . 10
3.4 Symmetry . . . 11
3.5 Basic Graphs . . . 12
3.6 Parametric Equations . . . 13
3.6.1 Finding the Cartesian Equation . . . 13
3.6.2 Asymptotes . . . 14
3.6.3 Intersections . . . 14
3.7 Conics . . . 15

1 Systems of Linear Equations

1.1 Forming the Equations

We read the context of the question and form equations depending on the context of the question.

Example 1.1

$$\left.\begin{aligned} a+b+c&=176 && —(1)\\ 0.85a+0.90b+0.95c&=161.80 && —(2)\\ 0.93a+0.95b+0.98c&=169.56 && —(3)\end{aligned}\right\} \Rightarrow a=32,\ b=44,\ c=100$$

Example 1.2

The "variables" may be expressions. To find x, y and z, use the given information to form the equations:

$$\left.\begin{aligned} 2x^3+3\ln y-\mathrm{e}^z&=18 && —(1)\\ x^3-\ln y+2\mathrm{e}^z&=9 && —(2)\\ x^3+2\ln y+\mathrm{e}^z&=11 && —(3)\end{aligned}\right\} \Rightarrow x^3=8,\ \ln y=1,\ \mathrm{e}^z=1$$

It is technically still 'linear' because we can think of $A=x^3$, $B=\ln y$, $C=\mathrm{e}^z$, which turns it into a linear system in A, B and C. A truly nonlinear equation could look like $x^2+x+y^2+2y-3z^3-xz+xy=1$; it cannot be made linear this way, because x appears as both x^2 and x, and there are product terms such as xz and xy.

1.2 Special Case: Number of Equations > Number of Variables

Sometimes the number of equations is more than the number of variables, and the solution must satisfy all of the equations.

Example 1.3

$$\left.\begin{aligned} \lambda+\mu&=1 && —(1)\\ 2\lambda-\mu&=5 && —(2)\end{aligned}\right\} \Rightarrow \lambda=2,\ \mu=-1$$

$$\lambda+3\mu=-1 \qquad —(3)$$

Sometimes there may be no solution, e.g. skew lines in vectors. In this case, solve for λ and μ using equations (1) and (2). Then sub into (3).

> **Note**
>
> Check (3): if LHS = RHS, the equations are consistent; if LHS ≠ RHS, there is no solution.

1.3 Special Case: Number of Equations < Number of Variables

Sometimes the number of equations is less than the number of variables, so there are infinitely many solutions.

Example 1.4

$$\left.\begin{aligned} x+y+z=6 \quad &—(1) \\ x-y+2z=5 \quad &—(2) \end{aligned}\right\} \Rightarrow x=\tfrac{11}{2}-\tfrac{3}{2}z,\ y=\tfrac{1}{2}+\tfrac{1}{2}z,\ z=z$$

> **Note**
>
> Useful trick: in contextual questions, the variables are usually non-negative integers, $z \in \mathbb{Z}_0^+$, so use this to find the solution from the general solution.

When $z=z$, it means that there are infinitely many solutions, and z can be any real value. So just sub that into x and y to get a correct solution.

$z=-1: (7,0,-1)$ $z=0: (\tfrac{11}{2},\tfrac{1}{2},0)$ $z=1: (4,1,1)$ $z=2: (\tfrac{5}{2},\tfrac{3}{2},2)$

$z=3: (1,2,3)$ $z=4: (-\tfrac{1}{2},\tfrac{5}{2},4)$ $z=5: (-2,3,5)$ $z=6: (-\tfrac{7}{2},\tfrac{7}{2},6)$

We can use the GC table to find all these solutions.

2 Inequalities


2.1 Solving Inequalities Graphically

Some inequalities cannot be solved algebraically, so we sketch the graphs and read off the solution.

- Rearrange to $f(x) > 0$, or compare $y = f(x)$ with $y = g(x)$.
- Sketch on the GC; mark the intersections and the asymptotes.
- Read off the x-values where the required graph is above or below.

Example 2.1

$$|x^2 - 4| \leqslant \ln(x+3)$$



Using GC, intersections at $x = 1.5746$, $x = 2.3840$

$$\therefore x = -2 \text{ or } 1.57 \leqslant x \leqslant 2.38$$


2.2 Sign Test Technique

We solve rational inequalities by finding where the expression changes sign.

- Make the right-hand side 0.
- Do **not** cross multiply unless the expression or bracket is always positive.
- Expressions that are always positive can be divided across, e.g. a quadratic with discriminant <0 and positive leading coefficient: $x^2+2x+5=(x+1)^2+4>0$.

Example 2.2

$$\frac{(x-1)(x-2)^2}{(x-3)(x-4)} \leqslant 0$$



$x=2$: numerator $=0$ (included)

$x=3,\ x=4$: denominator $=0$ (excluded)

$\therefore x \leqslant 1$ or $x=2$ or $3<x<4$

You can sketch the graph to check the answer.

> **Note**
>
> Crossing a factor with an odd power changes the sign; crossing a factor with an even power keeps the sign the same.

2.3 Common Substitutions

Once an inequality in x is solved, we can replace x with an expression to deduce the solution of a related inequality.


Replace x by	**Original inequality**		
	$x>2$	$x<-1$	$-1<x<2$
e^x	$x>\ln 2$	no solution	$x<\ln 2$
$\ln x$	$x>e^2$	$0<x<e^{-1}$	$e^{-1}<x<e^2$
$-x$	$x<-2$	$x>1$	$-2<x<1$
$x+1$	$x>1$	$x<-2$	$-2<x<1$
$\lvert x\rvert$	$x<-2$ or $x>2$	no solution	$-2<x<2$
x^2	$x<-\sqrt{2}$ or $x>\sqrt{2}$	no solution	$-\sqrt{2}<x<\sqrt{2}$
$\frac{1}{x}$	$\left(0<x<\frac{1}{2}\right)$	$(-1<x<0)$	$\left(x<-1 \text{ or } x>\frac{1}{2}\right)$
$\cos x,\ 0\leqslant x\leqslant 2\pi$	(no solution)	(no solution)	$(0\leqslant x<\pi \text{ or } \pi<x\leqslant 2\pi)$

Regardless of whatever substitution you use, using a graph will always work, but some can be solved without a graph.

For $\frac{1}{x}$, $\sin x$ and $\cos x$, draw the graph to solve.


Example 2.3

$$\frac{1}{x}>2, \quad \frac{1}{x}<-1, \quad -1<\frac{1}{x}<2$$



Example 2.4

$$\cos x>\tfrac{1}{2}, \quad \cos x<-\tfrac{1}{2},$$
$$-\tfrac{1}{2}<\cos x<\tfrac{1}{2}, \quad 0\leqslant x\leqslant 2\pi$$




3 Graphing Techniques


3.1 Key Features of a Curve

Key features of a curve:

(a) **x-intercepts:** let $y=0$.
e.g. $y=x^2-1$: $x^2-1=0 \Rightarrow x=\pm 1$.




y-intercepts: let $x=0$.
e.g. $y=x+2$: $y=2$.




(b) **Stationary points**, when $\frac{dy}{dx}=0$ or $\frac{dy}{dx}$ is undefined. Label upon request.

Horizontal turning point: $\frac{dy}{dx}=0$ (tangent horizontal).
e.g. $y=-(x-1)^2+2$, maximum point $(1,2)$.



Vertical turning point: $\frac{dy}{dx}$ undefined (tangent vertical).
e.g. $x=(y-1)^2+1$, turning point $(1,1)$.



(c) **Asymptotes**


3.2 Asymptotes

3.2.1 Vertical Asymptotes

If as $x \to k$, $y \to \pm\infty$, then $x=k$ is a vertical asymptote. Common occurrences:


1. Fractions. Denominator $=0$.

e.g. $y=\dfrac{1}{x-1}$: vertical asymptote $x=1$.




2. Logarithmic graphs. Inside $\ln=0$.

e.g. $y=\ln(x-2)$: vertical asymptote $x=2$.



3. Tan graphs. Inside $\tan=\dfrac{\pi}{2}+k\pi,\ k\in\mathbb{Z}$.

e.g. $y=\tan x$: vertical asymptotes $x=\pm\dfrac{\pi}{2},\ \pm\dfrac{3\pi}{2},\ldots$



3.2.2 Horizontal Asymptotes

As $x\to\pm\infty$, if $y\to k$, then $y=k$ is a horizontal asymptote.

- To find horizontal asymptotes, one should sub in $x\to\pm\infty$ into the equation. This is the only foolproof way to do it.
- A curve **CAN** intersect its horizontal asymptote.


Useful trick to find horizontal asymptotes

For rational fns $\frac{P(x)}{Q(x)}$.

numerator power > denom power

⇒ no horizontal asymptote


e.g. $y=\frac{x^2}{x-1}$



numerator power < denom power

⇒ horiz asymptote: $y=0$


e.g. $y=\frac{1}{x^2+1}$



numerator power = denom power

⇒ horiz asymptote: ratio of coeff of highest power.

e.g. $y=\frac{2x+1}{x-1}$, $y=2$



3.2.3 Oblique Asymptotes


An asymptote of the form $y=mx+c$, $m\neq 0$.

e.g. $y=2x+1+\frac{3}{x-2}$, $y=3x-5+e^{-x}$

$$y=\frac{ax^2+bx+c}{dx+e}=Ax+B+\frac{C}{dx+e}$$

Oblique asymptote: $y=Ax+B$. Vertical asymptote: $x=-\frac{e}{d}$.


2 turning points



e.g. $y=x+1+\frac{1}{x-1}$

$\left(\frac{Cd}{A}>0\right)$

No turning points.



e.g. $y=x+1-\frac{1}{x-1}$

$\left(\frac{Cd}{A}<0\right)$


Note

For parametric curves, oblique asymptotes (if they exist) can only be found via the cartesian equation.


3.3 Graphs of Rational Functions

Two common shapes for each type.

Linear / linear: $y=\dfrac{Ax+B}{Cx+D}$ Horizontal asymptote: $y=\dfrac{A}{C}$. Vertical asymptote: $x=-\dfrac{D}{C}$.




e.g. $y=\dfrac{x+1}{x-1}$




e.g. $y=\dfrac{x-1}{x+1}$

Quadratic / linear: $y=\dfrac{Ax^2+Bx+C}{Dx+E}$ Oblique asymptote; vertical asymptote $x=-\dfrac{E}{D}$.




e.g. $y=x+\dfrac{1}{x-1}$




e.g. $y=x-\dfrac{1}{x-1}$

Linear / quadratic: $y=\dfrac{ax+b}{cx^2+dx+e}$ Horizontal asymptote: $y=0$. Vertical asymptotes: Solve $cx^2+dx+e=0$ (possibly no roots).



e.g. $y=\dfrac{x}{(x-1)(x+1)}$



e.g. $y=\dfrac{x-2}{(x-1)(x+1)}$


Quadratic curves: $y=a(x-h)^2+k$, $y=\alpha(x-m)(x-n)$.

3.4 Symmetry

Even functions: $\mathrm{f}(-x)=\mathrm{f}(x)$.


The graph is symmetric about the y-axis.

e.g. $y=x^2$



Symmetry about the x-axis: replacing y by $-y$ gives the same equation.


e.g. $y^2=x$



Odd functions: $\mathrm{f}(-x)=-\mathrm{f}(x)$.


The graph is symmetric about the origin.

e.g. $y=x^3-x$




3.5 Basic Graphs


- Polynomial (linear, quadratic, cubic).
- Exponential. $y=ae^x+b$.
- Logarithm. $y=\ln(ax+b)$.
- Trigo. $y=\sin x$, $\cos x$, $\tan x$.
- Inverse Trigo. $y=\sin^{-1}x$, $\cos^{-1}x$, $\tan^{-1}x$.




Linear $y=x+1$




Quadratic $y=x^2-1$




Cubic $y=x^3-x$




Exponential $y=e^x$




Logarithm $y=\ln x$




Reciprocal $y=\frac{1}{x}$




$y=\sin x$




$y=\cos x$




$y=\tan x$



$y=\sin^{-1}x$



$y=\cos^{-1}x$



$y=\tan^{-1}x$

3.6 Parametric Equations

A curve C can be described by giving both coordinates in terms of a parameter t:

$$x=x(t), \quad y=y(t), \quad t\in T$$


Example.

$x=2\cos t+1, \quad y=3\sin t-2, \quad 0\leqslant t\leqslant \pi$

$t=0$: $(3,-2)$ $\quad t=\frac{\pi}{2}$: $(1,1)$ $\quad t=\pi$: $(-1,-2)$

$x=0$: $\left(0, \frac{3\sqrt{3}}{2}-2\right)$

$y=0$: $\left(1\pm\frac{2\sqrt{5}}{3}, 0\right)$



CAUTION: sketching from parametric equations

ALWAYS take note of the range of t when sketching graphs. The restriction on t decides which part of the curve exists: above, $0\leqslant t\leqslant\pi$ gives only the upper half of the ellipse, with endpoints $(3,-2)$ and $(-1,-2)$, not the whole ellipse.

3.6.1 Finding the Cartesian Equation

The cartesian equation of C is an equation without the parameter t. Common techniques:

① **Make t the subject.** e.g. $x=2t-1, y=3t^2+e^t, t\in\mathbb{R}$

$$t=\frac{x+1}{2}$$

$$y=3\left(\frac{x+1}{2}\right)^2+e^{\frac{x+1}{2}}$$

② **Trigo identity formulas.** e.g. $x=2\cos t+1, y=3\sin t-2, 0\leqslant t\leqslant\pi$

$$\cos t=\frac{x-1}{2}, \quad \sin t=\frac{y+2}{3}$$

$$\cos^2 t+\sin^2 t=1$$

$$\frac{(x-1)^2}{4}+\frac{(y+2)^2}{9}=1, \quad y\geqslant -2$$

③ **Others.** e.g. $x=t-\frac{1}{t}, y=t+\frac{1}{t}, t\neq 0$

$$y+x=2t, \quad y-x=\frac{2}{t}$$

$$(y+x)(y-x)=4$$

$$y^2-x^2=4$$

Note: Avoid finding the cartesian eqn if the question did not ask you to.

3.6.2 Asymptotes

- Find (if possible) $t \to t_0$ such that $x \to x_0, y \to \pm\infty$ (vertical asymptote $x = x_0$), or $y \to y_0, x \to \pm\infty$ (horizontal asymptote $y = y_0$).
- Oblique asymptotes (if they exist) can only be found via the cartesian equation.


3.6.3 Intersections

Intersection between a parametric curve and another curve: sometimes both equations are needed.

3.7 Conics

Circle: $(x-h)^2+(y-k)^2=r^2$

Centre: (h,k) radius: r




Ellipse: $\dfrac{(x-h)^2}{a^2}+\dfrac{(y-k)^2}{b^2}=1$

Centre: (h,k), "horizontal radius" $=a$, "vertical radius" $=b$

(for $a>b$: major axis horizontal, length $2a$, semi-major a; minor axis vertical, length $2b$, semi-minor b)

minor axis $2b$ (vertical)




major axis $2a$ (horizontal)

⋆ RHS ⇒ numerator coefficients of x^2, y^2 must all be 1.

Hyperbola: Type I

$\dfrac{(x-h)^2}{a^2}-\dfrac{(y-k)^2}{b^2}=1$


Centre: (h,k), Distance from centre to vertices $=a$



Type II

$\dfrac{(y-k)^2}{b^2}-\dfrac{(x-h)^2}{a^2}=1$

Centre: (h,k), Distance from centre to vertices $=b$




⋆ RHS ⇒ numerator coefficients of x^2, y^2 must all be 1.

⋆ Equation of asymptotes: $y-k=\pm\dfrac{b}{a}(x-h)$ (dashed in the sketches).

Parabola:

· $y-k=a(x-h)^2$ Turning point $\left(\dfrac{dy}{dx}=0\right)$: (h,k)

· $x-h=b(y-k)^2$ Turning point $\left(\dfrac{dy}{dx}\text{ undefined}\right)$: (h,k)



Common question types for curve sketching

→ Find the range/set of values that y can/cannot take. [VJC 2025]

→ Find the range of values of a such that C has no turning points [SAJC 2025]

→ Line rotation about pivot question. [SAJC 2025]

→ Conic superimposed onto graph. [DHS 2025]