Syllabus 9758 Statistics

Example Questions

Maximus Pung

Contents

1 Permutations and Combinations 3

Question 1.1 Choosing with "at least" conditions; circular arrangements . . . 3
Question 1.2 Probabilities from a table; access codes from BOOKKEEPER; "at least one from each group" . . 5
Question 1.3 Choosing teams (football) . . . 8
Question 1.4 Arranging in rows ($2 \times n$); round table . . . 10

2 Probability 12

Question 2.1 Tree diagram . . . 12
Question 2.2 Sampling; Venn diagram: range of values of a probability . . . 14
Question 2.3 Sum to infinity of a GP; distribution of a score; fair game . . . 17
Question 2.4 Contextual probability . . . 19

3 Discrete Random Variables 21

Question 3.1 Unknowns in the table; solving simultaneously . . . 21
Question 3.2 Fair game . . . 22
Question 3.3 Total of two games (not CLT) . . . 24
Question 3.4 Expectation of a new variable (new table) . . . 26
Question 3.5 Central Limit Theorem . . . 28

4 Binomial Distribution 30

Question 4.1 Double binomial . . . 30
Question 4.2 The kth trial is the mth success . . . 32
Question 4.3 Bug turning left and right . . . 33
Question 4.4 Range of p given the mode . . . 34
Question 4.5 Central Limit Theorem . . . 35

5 Normal Distribution 37

Question 5.1 Combining independent normal variables . . . 37
Question 5.2 Sketching a normal curve . . . 39
Question 5.3 invNorm simultaneous equations; by symmetry . . . 41
Question 5.4 Binomial count of a normal event . . . 43

6 Hypothesis Testing 47

Question 6.1 Basic test; reverse: smallest significance level, range of variance . . . 47
Question 6.2 Data as $(x-k)$ and $(x-k)^2$ . . . 50
Question 6.3 Data in a frequency table (GC steps) . . . 53
Question 6.4 Sample variance given ($\frac{n}{n-1} \times$ sample variance) . . . 56
Question 6.5 Data as x and $(x-k)^2$ . . . 59

7 Correlation and Linear Regression 62

Question 7.1 Comparing models (scatter diagram and r); estimate and reliability 62
Question 7.2 Missing y value from the LSRL via $(\bar{x}, \bar{y})$ 64
Question 7.3 Sum of squares of residuals 67
Question 7.4 Abi and Bhani: residuals and least squares 71

Question 6.2 Data as $\Sigma(x-k)$ and $\Sigma(x-k)^2$ 50
Question 6.3 Data in a frequency table (GC steps) 53
Question 6.4 Sample variance given ($\frac{n}{n-1}\times$ sample variance) 56
Question 6.5 Data as Σx and $\Sigma(x-k)^2$ 59

7 Correlation and Linear Regression 62

Question 7.1 Comparing models (scatter diagram and r); estimate and reliability 62
Question 7.2 Missing y value from the LSRL via $(\bar{x},\bar{y})$ 64
Question 7.3 Sum of squares of residuals 67
Question 7.4 Abi and Bhani: residuals and least squares 71

1 Permutations and Combinations

1.1 A treasure hunt game is to be played in a 3-storey building. There are 12 specific locations where 5 treasures are to be hidden. Among the 5 treasures, 4 of them are identical. The locations are distributed across the storeys as follows:

- 3 locations on the 1st storey,
- 3 locations on the 2nd storey, and
- 6 locations on the 3rd storey.

Each location can hold at most one treasure.

(a) Find the number of ways to distribute the 5 treasures among the 12 locations for each of the following separate cases:

(i) The 5 treasures can be hidden in any of the 12 locations. [1]


(ii) Exactly 2 treasures must be hidden on the 3rd storey. [2]

(iii) There must be at least one treasure on each of the 1st and 2nd storeys, and at least 2 treasures on the 3rd storey. [3]

Five teams of 2 players each take part in the game. The 10 players and 2 game masters are to be seated around a round table for a briefing.


(b) Find the number of different seating arrangements such that players from the same team must be seated next to each other but the two game masters must not be seated next to each other. [3]

[TJC 2025 Prelim P2 Q6]

Solution		Comments
(a)(i) **[1]**	Required number of ways $= \binom{12}{5}\binom{5}{1} = 3960$  choose 5 of the 12 locations: $\binom{12}{5}$; the distinct treasure ($\star$) takes 1 of these 5: $\binom{5}{1}$	

(a)(ii) [2]

Required number of ways $= \binom{6}{2}\binom{6}{3}\binom{5}{1} = 1500$



then the distinct treasure goes to 1 of the 5 chosen locations: $\binom{5}{1}$

Alternative method:

Case 1: 2 not identical at 3rd storey $\binom{6}{2} \times 2! \binom{6}{3} = 600$

Case 2: 2 identical at 3rd storey $\binom{6}{2}\binom{6}{3}\frac{3!}{2!} = 900$

(a)(iii) [3]

Case 1: 2 in 1st storey, 1 in 2nd storey, 2 in 3rd storey

Number of ways $= \binom{3}{2}\binom{3}{1}\binom{6}{2}\binom{5}{1} = 675$

Case 2: 1 in 1st storey, 2 in 2nd storey, 2 in 3rd storey

Number of ways $= \binom{3}{1}\binom{3}{2}\binom{6}{2}\binom{5}{1} = 675$

Case 3: 1 in 1st storey, 1 in 2nd storey, 3 in 3rd storey

Number of ways $= \binom{3}{1}\binom{3}{1}\binom{6}{3}\binom{5}{1} = 900$

Total number of ways $= 2250$

Case	1st (3)	2nd (3)	3rd (6)	ways
1	2	1	2	675
2	1	2	2	675
3	1	1	3	900


each row: $\binom{3}{\cdot}\binom{3}{\cdot}\binom{6}{\cdot} \times \binom{5}{1}$ (distinct treasure); 5 treasures in total

(b) [3]

Number of ways to arrange the 10 participants $= (5-1)!(2!)^5 = 768$

Number of ways to slot in the 2 game masters $= \binom{5}{2} 2! = 20$

Required number $= 768 \times 20 = 15360$



$(5-1)!$ ways round the table,
$(2!)^5$ within the 5 teams;
game masters in 2 of the 5 gaps ∧:
$\binom{5}{2}$ 2! (never next to each other)

1.2 **(a)** In a survey conducted by a library, a group of 100 members were asked the number of days in a month they visit the library. The results are shown below:

	1 or less	2 to 4	5 or more
Male	15	25	22
Female	20	$18-n$	n

A member is selected at random from the group. The events are defined as follows:

A: The member visits the library on 4 days or less in a month.

B: The member visits the library on 2 days or more in a month.

F: The member is a female.

Find the following probabilities in terms of n.

(i) $\mathrm{P}(A \cup B')$ [1]

(ii) $\mathrm{P}(F \mid A')$ [2]

(iii) Given that $\mathrm{P}(A \cap B) = \frac{3}{10}$, find the value of n. Hence determine if A and B are independent, justifying your answer. [3]

(b) A librarian is organising a large community reading event. To manage registration, he needs to create 4-letter access codes for the participants using the letters from the word BOOKKEEPER.

(i) How many access codes can be formed if all 4 letters are distinct from one another? [1]


(ii) Find the total number of access codes that can be created given that there is no restriction. [3]

To run an event, the librarian plans to select a team of 12 volunteers from a group of 18. The volunteers comprise three age groups: 5 youths, 6 young adults, and 7 seniors.

(iii) In how many ways can the team be formed if at least one volunteer must be chosen from each age group? [3]

[ASRJC 2025 Prelim P2 Q11]

Solution		Comments
(a)(i) **[1]**	$\mathrm{P}(A \cup B') = \mathrm{P}(A)$ $= \dfrac{15+25+20+(18-n)}{100}$ or $\dfrac{100-22-n}{100}$ $= \dfrac{78-n}{100}$	
(a)(ii) **[2]**	$\mathrm{P}(F \mid A') = \dfrac{\mathrm{P}(F \cap A')}{1-P(A)} = \dfrac{\frac{n}{100}}{1-\frac{78-n}{100}} = \dfrac{n}{22+n}$	
(a)(iii) **[3]**	Given $\mathrm{P}(A \cap B) = \dfrac{3}{10}$ $\mathrm{P}(A \cap B) = \dfrac{25+18-n}{100} = \dfrac{43-n}{100}$ $\dfrac{43-n}{100} = \dfrac{3}{10} \Rightarrow n = 13$ $\mathrm{P}(A)\mathrm{P}(B) = \dfrac{65}{100} \times \dfrac{65}{100} = \dfrac{169}{400} \neq \dfrac{3}{10}$ Since $\mathrm{P}(A)\mathrm{P}(B) \neq \mathrm{P}(A \cap B)$, A and B are **NOT** independent **Alternatively**, comparing $\mathrm{P}(A \mid B)$ with $\mathrm{P}(A)$ $\mathrm{P}(A \mid B) = \dfrac{\mathrm{P}(A \cap B)}{\mathrm{P}(B)} = \dfrac{\frac{3}{10}}{\frac{65}{100}} = \dfrac{6}{13}$ Since $\mathrm{P}(A) = \dfrac{13}{20}$ and $\mathrm{P}(A \mid B) \neq \mathrm{P}(A)$, A and B are **NOT** independent	
(b)(i) **[1]**	3Es, 2Os, 2Ks, 1P, 1R, 1B Number of 4-letter code words $= {}^6C_4 \times 4! = 360$ or ${}^6P_4 = 360$	
(b)(ii) **[3]**	Case 1: All distinct (from b(i)) Number of 4-letter code words $= {}^6C_4 \times 4! = 360$ Case 2: One pair of repeated letters. (Eg XXYZ) Number of 4-letter code words $= {}^3C_1 \times {}^5C_2 \times \dfrac{4!}{2!} = 360$ Case 3: Two pairs of repeated letters. (Eg XXYY) Number of 4-letter code words $= {}^3C_2 \times \dfrac{4!}{2!2!} = 18$ Case 4: One set of 3 repeated letters (Eg EEEX) Number of 4-letter code words $= {}^5C_1 \times \dfrac{4!}{3!} = 20$ $\therefore$ Total number of 4-letter code word $= 360+360+18+20 = 758$ E E E O O K K P R B — 3 E, 2 O, 2 K (repeatable); P, R, B X Y Z W — Case 1: all distinct X X Y Z — Case 2: one pair X X Y Y — Case 3: two pairs X X X Y — Case 4: three alike (only E)	

(b)(iii) **[3]**	Total number of ways $= {}^{18}C_{12} = 18564$ No youth $= {}^{13}C_{12} = 13$ No young adult $= {}^{12}C_{12} = 1$ Total possible ways $= 18564 - 13 - 1 = 18550$  all teams – (no youth: 12 from 13) – (no young adult: 12 from 12); no senior is impossible (only 11 others)	

1.3 A group of 12 players is available for a street football match. The group consists of 5 attackers, 4 defenders and 3 goalkeepers. A team of 5 players is selected at random from the group.

(a) Find the

(i) number of different teams containing exactly 2 attackers, 2 defenders and 1 goalkeeper, [2]


(ii) probability that the team contains exactly 1 goalkeeper and at most 2 attackers. [3]

The coach has already decided which goalkeeper will play in a match. To help him select from the 5 attackers and 4 defenders, he conducts a passing drill involving these players. For each pass, 1 player is chosen at random to receive the ball. After each pass, the ball is returned to the coach before the next player is chosen. A player may receive the ball more than once. The coach makes 4 passes in this manner. Find the probability that

(b) exactly 3 passes are made to attackers, [2]

(c) either the first 3 passes or the last 3 passes are made to attackers (or both). [3]

[JPJC 2026 Prelim P2 Q8]

Solution		**Comments**
(a)(i) **[2]**	Number of teams $= {}^5C_2 \times {}^4C_2 \times {}^3C_1 = 180$ 	

(a)(ii) **[3]**

Case 1: 1 goalkeeper, 2 attackers, 2 defenders $^3C_1 \times {}^5C_2 \times {}^4C_2 = 180$

Case 2: 1 goalkeeper, 1 attacker, 3 defenders $^3C_1 \times {}^5C_1 \times {}^4C_3 = 60$

Case 3: 1 goalkeeper, 0 attacker, 4 defenders $^3C_1 \times {}^5C_0 \times {}^4C_4 = 3$

Total number of ways $= {}^{12}C_5 = 792$

Hence probability that team has at most 2 attackers

$= \frac{180+60+3}{792} = \frac{243}{792}$ or 0.307

Case	G (3)	A (5)	D (4)	ways
1	1	2	2	180
2	1	1	3	60
3	1	0	4	3

team of 5: exactly 1 goalkeeper, at most 2 attackers

(b) **[2]**

Req probability

$= \mathrm{P}(\mathrm{AAAD}) + \mathrm{P}(\mathrm{AADA}) + \mathrm{P}(\mathrm{ADAA}) + \mathrm{P}(\mathrm{DAAA})$

$= \left(\frac{5}{9}\right)^3 \left(\frac{4}{9}\right) \times 4 = \frac{2000}{6561}$ or 0.305

pass 1	pass 2	pass 3	pass 4	
D	A	A	A	$\left(\frac{5}{9}\right)^3 \left(\frac{4}{9}\right)$
A	D	A	A	$\left(\frac{5}{9}\right)^3 \left(\frac{4}{9}\right)$
A	A	D	A	$\left(\frac{5}{9}\right)^3 \left(\frac{4}{9}\right)$
A	A	A	D	$\left(\frac{5}{9}\right)^3 \left(\frac{4}{9}\right)$

×4 positions for D


(c) **[3]**

Method 1:

Req probability $= \mathrm{P}(\mathrm{AAA_}) + \mathrm{P}(\mathrm{_AAA}) - \mathrm{P}(\mathrm{AAAA})$

$= \left(\frac{5}{9}\right)^3 + \left(\frac{5}{9}\right)^3 - \left(\frac{5}{9}\right)^4$

$\frac{1625}{6561}$ or 0.248



Method 1: add both, subtract overlap

Method 2: AAAD + DAAA + AAAA

Method 2:

Req prob $= \mathrm{P}(\mathrm{AAAD}) + \mathrm{P}(\mathrm{DAAA}) + \mathrm{P}(\mathrm{AAAA})$

$= \left(\frac{5}{9}\right)^3 \left(\frac{4}{9}\right) + \left(\frac{4}{9}\right) \left(\frac{5}{9}\right)^3 + \left(\frac{5}{9}\right)^4$

$\frac{1625}{6561}$ or 0.248

1.4 An office team of 10 people includes 7 men and 3 women named Anne, Beth, and Cathie. For an upcoming fire drill exercise, 5 individuals will be chosen, each assigned a unique role, to carry out the drill. Determine the number of possible ways to select 5 people from this group of 10





(i) to conduct the fire drill, [1]

(ii) such that at most 1 woman is selected to conduct the fire drill. [2]

(iii) After the fire drill exercise, the 10 people are to hold a discussion at a round table with 10 identical seats. Determine the number of ways in which Beth is seated between Anne and Cathie. [1]

(iv) A group photo of the 10 people, arranged in two rows of five, was taken after the discussion. Determine the number of ways in which Beth is not standing beside Anne or Cathie. [4]

[ASRJC 2024 Prelim P2 Q8]

Solution		Comments
(i) **[1]**	Number of ways $= \binom{10}{5} \times 5! = 30240$  choose 5 of 10: $\binom{10}{5}$, then assign the 5 distinct roles: 5!	
(ii) **[2]**	Number of ways $= \left[\binom{7}{5}\binom{3}{0} + \binom{7}{4}\binom{3}{1}\right] 5!$ $= (21 \times 1 + 35 \times 3)5! = 15120$	
(iii) **[1]**	Number of ways $= (8-1)!2! = 10080$  $(8-1)!$ round the table, $2!$ for A B C or C B A	
(iv) **[4]**	Case 1 – Beth in 1 row while Anne and Cathie are in another row $\binom{7}{4} \times 5! \times 5! \times 2 = 1008000$ Case 2 – Beth and one of them in 1 row $\binom{7}{3} \times 3! \times \binom{2}{1} \times \binom{4}{2} \times 2! \times 5! \times 2 = 1209600$ Case 3 – Anne, Beth and Cathie are in the same row $\underbrace{\binom{7}{2} \times 2! \times \binom{3}{2} \times 2! \times 2! \times 5! \times 2}_{\text{A and C together}} + \underbrace{\binom{7}{2} \times 2! \times 3! \times 5! \times 2}_{\text{A and C separated}} = 181440$ Thus number of ways $= 1008000 + 1209600 + 181440 = 2399040$  Case 1: B in one row, A and C in the other. Case 2: B and one of A, C in the same row, not side by side.  Case 3: A, B, C in the same row, B beside neither A nor C.	

2 Probability

2.1 An online game consists of three stages. All players start with Stage 1, and no matter whether they win or lose, they proceed to Stage 2. Players are allowed to proceed to Stage 3 only if they have won at least one of the first two stages.

Ming's probability of winning Stage 1 is p.


If Ming wins a stage, his probability of winning the next stage is halved. If he loses a stage, his probability of winning remains unchanged in the next stage.

(a) Draw a probability tree diagram to represent all possible outcomes for Ming over the three stages. [3]

(b) Find the range of possible values of p if the probability that Ming wins all three stages is more than 0.1. [2]

(c) It is now known that $p = 0.9$. Given that Ming proceeds to Stage 3, find the probability that he wins exactly two stages out of the three. [4]

[EJC 2025 Prelim P2 Q6]

Solution		Comments
(a) **[3]**		
(b) **[2]**	$p(0.5p)\ (0.5^2p) > 0.1$ $p^3 > 0.8$ $p > 0.928$	
(c) **[4]**	$\mathrm{P}(\text{wins exactly 2 stages out of 3} \mid \text{proceeds to Stage 3})$ $= \dfrac{\mathrm{P}(\text{wins exactly 2 stages out of 3} \cap \text{proceeds to Stage 3})}{\mathrm{P}(\text{proceeds to Stage 3})}$ $= \dfrac{0.9(0.45)(1-0.225)+0.9(0.55)(0.45)+0.1(0.9)(0.45)}{1-(0.1)(0.1)}$ $= \dfrac{0.577125}{0.99} = 0.583$	

2.2 (a) In a certain university, there are 110 Co-Curricular Activities (CCAs) clustered into 4 categories. There are 22 Arts and Culture CCAs, 16 Community Service CCAs, 38 Physical Sports CCAs and 34 Special Interest CCAs.

(i) Albert wishes to find out about approaches to training of the Physical Sports CCAs, so he sends a questionnaire to 22 Physical Sports CCAs. Explain whether these 22 Physical Sports CCAs form a sample or a population. [1]

(ii) Benedict wishes to investigate the level of student engagement in CCAs, but does not want to obtain the detailed information necessary from all 110 CCAs. Explain how he should carry out his investigation, and why he should do the investigation in this way. [2]

(iii) Find the number of different possible samples of 16 CCAs, with 4 CCAs chosen from each category. [2]

(b) The probability of events A, B and C occurring are given by $\mathrm{P}(A)=\frac{8}{25}$, $\mathrm{P}(B)=\frac{61}{100}$ and $\mathrm{P}(C)=\frac{7}{20}$ respectively. It is also known that $\mathrm{P}(A\cap C)=\frac{11}{100}$, $\mathrm{P}(B\mid C)=\frac{13}{35}$ and $\mathrm{P}(A\cap B\cap C)=\frac{7}{100}$.

(i) Find $\mathrm{P}(B\cap C)$. [2]

(ii) If events A and B are independent, find $\mathrm{P}(A\cap B\cap C')$. [2]


(iii) If events A and B are **not** known to be independent, find the greatest and least possible values of $\mathrm{P}(A\cap B\cap C')$. [4]

[CJC 2024 Prelim P2 Q10]

Solution		Comments
(a)(i) **[1]**	These 22 CCAs form a sample since this number is less than the population size of Physical Sports CCAs which is 38.	
(a)(ii) **[2]**	Benedict should select a random sample from the 110 CCAs: every CCA has an equal probability to be selected to be in the sample and the CCAS are selected independently. (If define without stating "random", need 2 full points.) A random sample helps to avoid bias.	
(a)(iii) **[2]**	Number of samples $={}^{22}C_4\times{}^{16}C_4\times{}^{38}C_4\times{}^{34}C_4$ $=4.56\times10^{16}$ (3 sf)	

(b)(i) **[2]**	$\mathrm{P}(B \mid C) = \frac{13}{35}$ $\frac{\mathrm{P}(B \cap C)}{\mathrm{P}(C)} = \frac{13}{35}$ $\frac{\mathrm{P}(B \cap C)}{\frac{7}{20}} = \frac{13}{35}$ $\mathrm{P}(B \cap C) = \frac{13}{100}$	
(b)(ii) **[2]**	$\mathrm{P}(A \cap B) = \mathrm{P}(A) \times \mathrm{P}(B)$ $= \frac{8}{25} \times \frac{61}{100}$ $= \frac{122}{625}$ $\mathrm{P}(A \cap B \cap C')$ $= \frac{122}{625} - \frac{7}{100}$ $= \frac{313}{2500}$	

(b)(iii)
[4]



Let $\mathrm{P}(A\cap B\cap C')=x$ where $0\leqslant x\leqslant 1$ ... (1)

Then

$\mathrm{P}(A\cap B'\cap C')=\frac{8}{25}-\frac{11}{100}-x=\frac{21}{100}-x$

$\mathrm{P}(A'\cap B\cap C')=\frac{61}{100}-\frac{13}{100}-x=\frac{12}{25}-x$

$\mathrm{P}(A\cup B\cup C)$

$=\mathrm{P}(C)+\mathrm{P}(A\cap B'\cap C')+\mathrm{P}(A'\cap B\cap C')+\mathrm{P}(A\cap B\cap C')$

$=\frac{7}{20}+\left(\frac{21}{100}-x\right)+\left(\frac{12}{25}-x\right)+x$

$=\frac{26}{25}-x$

Consider the probabilities $\mathrm{P}(A\cap B'\cap C')$, $\mathrm{P}(A'\cap B\cap C')$ and $\mathrm{P}(A\cup B\cup C)$,

$0\leqslant \frac{21}{100}-x\leqslant 1$ and $0\leqslant \frac{12}{25}-x\leqslant 1$ and $0\leqslant \frac{26}{25}-x\leqslant 1$

$-\frac{79}{100}\leqslant x\leqslant \frac{21}{100}$ ... (2) $-\frac{13}{25}\leqslant x\leqslant \frac{12}{25}$ ... (3)

$\frac{1}{25}\leqslant x\leqslant \frac{26}{25}$ ... (4)

Taking intersection of inequalities (1) to (4), $\frac{1}{25}\leqslant x\leqslant \frac{21}{100}$

$\min \mathrm{P}(A\cap B\cap C')=\frac{1}{25}$ and $\max \mathrm{P}(A\cap B\cap C')=\frac{21}{100}$

2.3 A bag contains one green disc, r red discs and $2r$ blue discs, where $r > 1$.

(a) Tim and Sim play a game by taking turns to draw a disc from the bag at random. The drawn disc is replaced after each turn. The first person to draw the green disc wins. If Sim started the game first, show that the probability of Sim winning the game is $\frac{3r+1}{6r+1}$. [2]

(b) In another game, a player draws 2 discs from the bag at random. If he draws a red disc, he scores 5 points, if he draws a blue disc, he scores 2 points and if he draws a green disc he scores 0. The player's score, denoted by T, is found by taking the difference in the number of points he scores for the two discs.

(i) Find $\mathrm{P}(T=t)$ for all possible values of t, leaving your answers in terms of r. [3]

(ii) Show that the expectation of the player's score is $\frac{2(2r+3)}{3r+1}$. [1]

(c) In a third game, a player draws a disc from the bag at random. If he draws a green disc, he wins \$0.15, if he draws a blue disc, he wins \$0.10 and if he draws a red disc, he loses \$0.25. May and Jun each draws a disc from the bag once, with the disc replaced in the bag after the first player drew. The probability that May ends up with more money than Jun is $\frac{27}{112}$. Find the value of r. [2]

Is the game fair for a player? Explain. [1]

[ACJC 2023 Prelim P2 Q9]

Solution		Comments
(a) **[2]**	$\frac{1}{3r+1}+\left(\frac{3r}{3r+1}\right)^2\times\frac{1}{3r+1}+\left(\frac{3r}{3r+1}\right)^4\times\frac{1}{3r+1}+\ldots$ $=\frac{\frac{1}{3r+1}}{1-\left(\frac{3r}{3r+1}\right)^2}=\frac{3r+1}{6r+1}$	
(b)(i) **[3]**	$\mathrm{P}(T=0)=\frac{r}{3r+1}\times\frac{r-1}{3r}+\frac{2r}{3r+1}\times\frac{2r-1}{3r}=\frac{5r-3}{3(3r+1)}$ $\mathrm{P}(T=2)=\frac{2r}{3r+1}\times\frac{1}{3r}\times 2!=\frac{4}{3(3r+1)}$ $\mathrm{P}(T=3)=\frac{2r}{3r+1}\times\frac{r}{3r}\times 2!=\frac{4r}{3(3r+1)}$ $\mathrm{P}(T=5)=\frac{r}{3r+1}\times\frac{1}{3r}\times 2!=\frac{2}{3(3r+1)}$	
(b)(ii) **[1]**	$\mathrm{E}(T)=\sum t\mathrm{P}(T=t)$ $=0\times\frac{10r-4}{3(3r+1)}+2\times\frac{4}{3(3r+1)}+3\times\frac{4r}{3(3r+1)}+5\times\frac{2}{3(3r+1)}$ $=\frac{8+12r+10}{3(3r+1)}=\frac{2(2r+3)}{3r+1}$	
(c) **[1]**	$\mathrm{P}(X_1>X_2)=\mathrm{P}(X_1=0.15,X_2=-0.25)$ $+\mathrm{P}(X_1=0.15,X_2=0.10)$ $+\mathrm{P}(X_1=0.10,X_2=-0.25)$ $\frac{2r^2+3r}{(3r+1)^2}=\frac{27}{112}$ $19r^2-174r+27=0$ $r=\frac{3}{19}\approx 0.15789$ (rejected) $\quad\therefore r=9$ $\mathrm{E}(X)=\sum x\mathrm{P}(X=x)$ $=-0.25\times\frac{9}{28}+0.10\times\frac{18}{28}+0.15\times\frac{1}{28}$ $=-0.010714\approx -0.0107$ Since $\mathrm{E}(x)<0$, the game is <u>not</u> fair for a player, as he/she will be expected to lose 1.07 cents every time he/she plays.	

x	-0.25	0.10	0.15
$\mathrm{P}(X=x)$	$\frac{r}{3r+1}$	$\frac{2r}{3r+1}$	$\frac{1}{3r+1}$

t	0	2	3	5
$\mathrm{P}(T=t)$	$\frac{5r-3}{3(3r+1)}$	$\frac{4}{3(3r+1)}$	$\frac{4r}{3(3r+1)}$	$\frac{2}{3(3r+1)}$

2.4 A basketball free throw game involves a team of two players, Ben and John. The game consists of at most three throws and the moment a player makes a successful shot, the team wins, and the game will end.

The game uses the following rules.

- Only one player is selected for each game and the probability that Ben is selected in a game is 0.7.
- The probabilities that Ben and John make a successful shot in any single attempt are 0.1 and 0.07 respectively.
- The shots are independent of each other.

(a) Find the probability that the team wins the game. [3]

(b) The team did not win the game. Find the probability that John was the one who was selected to shoot. [3]

(c) The team attempts the game repeatedly until the first game is won. Find the least number of attempts required such that the probability of winning within n games is at least 0.95. [2]

[HCI 2025 Prelim P2 Q6]

Solution		Comments
(a) **[3]**	$\mathrm{P}(\text{wins})$ $=\mathrm{P}(\text{wins and Ben selected})+\mathrm{P}(\text{wins and John selected})$ $=\mathrm{P}(\text{wins}\mid\text{Ben selected})\mathrm{P}(\text{Ben selected})$ $+\mathrm{P}(\text{wins}\mid\text{John selected})\mathrm{P}(\text{John selected})$ $=(0.1+0.9\times0.1+0.9^2\times0.1)\ (0.7)$ $+(0.07+0.93\times0.07+0.93^2\times0.07)\ (0.3)$ $=0.2483929$ OR $\mathrm{P}(\text{wins})$ $=\mathrm{P}(\text{wins and Ben selected})+\mathrm{P}(\text{wins and John selected})$ $=\mathrm{P}(\text{wins}\mid\text{Ben selected})\mathrm{P}(\text{Ben selected})+$ $\mathrm{P}(\text{wins}\mid\text{John selected})\mathrm{P}(\text{John selected})$ $=(1-0.9^3)\ (0.7)+(1-0.93^3)\ (0.3)$ $=0.2483929$	

(b) **[3]**

$$\text{P(John selected} \mid \text{did not win)}$$
$$= \frac{\text{P(John selected and did not win)}}{\text{P(did not win)}}$$
$$= \frac{0.3 \times 0.93^3}{1 - 0.2483929}$$
$$= 0.321 \text{ (3sf)}$$

(c) **[2]**

Required probability $= 1 -$ P(did not win in all n games)

$$= 1 - (1 - 0.2483929)^n$$

OR

Required probability

= P(win in 1st game) + P(win in 2nd game)

+ P(win in 3rd game) + … + P(win in n^{th} game)

$$= 0.2483929 + (1 - 0.2483929)(0.2483929)$$
$$+ (1 - 0.2483929)^2 (0.2483929) + \ldots$$
$$+ (1 - 0.2483929)^{n-1} (0.2483929)$$
$$= \frac{0.2483929\,(1 - (1 - 0.2483929)^n)}{1 - (1 - 0.2483929)}$$
$$= 1 - (1 - 0.2483929)^n$$

$$1 - (1 - 0.2483929)^n \geqslant 0.95$$

$$(1 - 0.2483929)^n \leqslant 0.05$$

n	$(1 - 0.2483929)^n$
10	$0.0575 > 0.05$
11	$0.0432 < 0.05$

Smallest n is 11.

See also Example 3 (b), (c) for another contextual probability question.

3 Discrete Random Variables

3.1 The probability distribution of a discrete random variable X is given as follows:

x	-2	-1	0	1	3
$\mathrm{P}(X=x)$	0.1	a	0.3	b	c

Given that $\mathrm{E}(X)=0.9$ and $\mathrm{Var}(X)=2.99$, find the values of a, b and c. [5]

[ACJC 2025 Prelim P2 Q8]

Solution		**Comments**
[5]	$\mathrm{E}(X)=(-2)(\frac{1}{10})+(-1)(a)+(0)(\frac{3}{10})+(1)(b)+(3)(c)$ $\frac{9}{10}=-\frac{2}{10}-a+b+3c$ $-a+b+3c=\frac{11}{10}$ ..... (1) $\mathrm{E}(X^2)=(-2)^2(\frac{1}{10})+(-1)^2(a)+(0)^2(\frac{3}{10})+(1)^2(b)+(3)^2(c)=$ $\frac{4}{10}+a+b+9c$ $\mathrm{Var}(X)=\mathrm{E}(X^2)-[\mathrm{E}(X)]^2$ $\frac{299}{100}=\left(\frac{4}{10}+a+b+9c\right)-\left(\frac{9}{10}\right)^2$ $a+b+9c=\frac{34}{10}$ ..... (2) $\sum \mathrm{P}(X=x)=1$ $\frac{1}{10}+a+\frac{3}{10}+b+c=1$ $a+b+c=\frac{6}{10}$ ..... (3) Solving (1), (2) and (3), we have $a=\frac{1}{10}$ or 0.1, $b=\frac{3}{20}$ or 0.15, $c=\frac{7}{20}$ or 0.35	

3.2 Anthony plays a game using a four-sided fair die with the numbers 1, 2, 3 and 4 printed on its sides. Anthony rolls the die and should the outcome be a prime number, he wins an amount equivalent to that number in dollars. Conversely, if the outcome is a non-prime number, Anthony will lose an amount equivalent to that number in dollars. The random variable X represents the amount of money Anthony wins in a roll of the die.

(a) Tabulate the probability distribution of X and find the value of $\mathrm{E}(X)$. [2]

(b) Find the value of the variance of X. [1]

(c) The random variables X_1 and X_2 are two independent observations of X. By drawing a table of outcomes for $|X_1+X_2|$ or otherwise, find $\mathrm{E}(|X_1+X_2|)$. [2]

Assume now that a biased six-sided die with numbers 2, 3, 4, 5, 6 and 7 printed on its sides, is used instead for the same game. It is known that for this biased die, the probability of obtaining a "3" is p, the probability of obtaining a "5" is $2p$ and the remaining four numbers each have an equal probability of occurring.

(d) Find the exact value of p for the game to be fair using the biased die. [3]

[ASRJC 2025 Prelim P2 Q7]

Solution		Comments
(a) **[2]**	$\begin{array}{\|c\|c\|c\|c\|c\|}\hline x & -4 & -1 & 2 & 3 \\ \hline \mathrm{P}(X=x) & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} & \frac{1}{4} \\ \hline \end{array}$ $\mathrm{E}(X)=\frac{-4}{4}+\frac{-1}{4}+\frac{2}{4}+\frac{3}{4}=0$	
(b) **[1]**	$\mathrm{Var}(X)=\left[\frac{(-4)^2}{4}+\frac{(-1)^2}{4}+\frac{2^2}{4}+\frac{3^2}{4}\right]-0^2=\frac{15}{2}$	
(c) **[2]**	Let $Y=\lvert X_1+X_2\rvert$	
	$\begin{array}{\|c\|c\|c\|c\|c\|}\hline X_2 \backslash X_1 & -4 & -1 & 2 & 3 \\ \hline -4 & 8 & 5 & 2 & 1 \\ \hline -1 & 5 & 2 & 1 & 2 \\ \hline 2 & 2 & 1 & 4 & 5 \\ \hline 3 & 1 & 2 & 5 & 6 \\ \hline \end{array}$	
	$\begin{array}{\|c\|c\|c\|c\|c\|c\|c\|}\hline y & 1 & 2 & 4 & 5 & 6 & 8 \\ \hline \mathrm{P}(Y=y) & \frac{4}{16} & \frac{5}{16} & \frac{1}{16} & \frac{4}{16} & \frac{1}{16} & \frac{1}{16} \\ \hline \end{array}$	
	$\mathrm{E}(Y)=\frac{4}{16}+2\left(\frac{5}{16}\right)+4\left(\frac{1}{16}\right)+5\left(\frac{4}{16}\right)+6\left(\frac{1}{16}\right)+8\left(\frac{1}{16}\right)=\frac{13}{4}$	
(d) **[3]**	The new probability distribution for X is as tabulated below:	
	$\begin{array}{\|c\|c\|c\|c\|c\|c\|c\|}\hline x & -6 & -4 & 2 & 3 & 5 & 7 \\ \hline \mathrm{P}(X=x) & \frac{1-3p}{4} & \frac{1-3p}{4} & \frac{1-3p}{4} & p & 2p & \frac{1-3p}{4} \\ \hline \end{array}$	
	$\mathrm{E}(X)=-6\left(\frac{1-3p}{4}\right)-4\left(\frac{1-3p}{4}\right)+2\left(\frac{1-3p}{4}\right)+3p+10p+$ $7\left(\frac{1-3p}{4}\right)=-\left(\frac{1-3p}{4}\right)+13p=-\frac{1}{4}+\frac{55}{4}p$	
	For the game to be fair, $-\frac{1}{4}+\frac{55}{4}p=0$	
	Hence $p=\frac{1}{55}$	

3.3 Andrea participates in a game show. In each round of the game, Andrea responds to a series of questions until she answers a question wrongly or she has answered five questions correctly. She is awarded one point for every correct answer and is awarded an additional 3 bonus points if she answers all five questions correctly. It is given that she answers each question independently and the probability she answers each question correctly is $\frac{4}{5}$.

(a) Show that the probability that Andrea obtains 2 points in one round of the game is $\frac{16}{125}$. [1]

(b) Find the probability distribution of the points that Andrea obtains in one round of the game. [2]

(c) Find the expectation and variance of the points that Andrea obtains in one round of the game. [2]

(d) Find the probability that Andrea obtains less than 3 points after two rounds of the game. [3]

[CJC 2024 Prelim P2 Q5]

Solution		Comments
(a) **[1]**	Probability that Andrea obtains 2 points in a round of the game $=\left(\frac{4}{5}\right)\left(\frac{4}{5}\right)\left(\frac{1}{5}\right)=\frac{16}{125}$	
(b) **[2]**	Let X be the random variable denoting the number of points obtained in a round.	
(c) **[2]**	**Method 1**: Using G.C., Expectation $=3.67232$ (exact); Variance $=3.22334^2=10.4$ (3 s.f.)	
	Method 2: $\mathrm{E}(X)=0\left(\frac{1}{5}\right)+1\left(\frac{4}{25}\right)+2\left(\frac{16}{125}\right)+3\left(\frac{64}{625}\right)+4\left(\frac{256}{3125}\right)+8\left(\frac{1024}{3125}\right)=3.67232$ $\mathrm{E}(X^2)=0^2\left(\frac{1}{5}\right)+1^2\left(\frac{4}{25}\right)+2^2\left(\frac{16}{125}\right)+3^2\left(\frac{64}{625}\right)+4^2\left(\frac{256}{3125}\right)+8^2\left(\frac{1024}{3125}\right)=23.87584$ $\mathrm{Var}(X)=\mathrm{E}(X^2)-[\mathrm{E}(X)]^2=23.87584-(3.67232)^2=10.4$ (to 3 s.f.)	
(d) **[3]**	Probability that Andrea obtains less than 3 points after two rounds $=\mathrm{P}(X_1+X_2<3)=\mathrm{P}(X_1+X_2=0)+\mathrm{P}(X_1+X_2=1)+\mathrm{P}(X_1+X_2=2)$ $=\mathrm{P}(X_1=0\cap X_2=0)+2\mathrm{P}(X_1=0\cap X_2=1)+2\mathrm{P}(X_1=0\cap X_2=2)+\mathrm{P}(X_1=1\cap X_2=1)$ $=\left(\frac{1}{5}\right)^2+2\left(\frac{1}{5}\right)\left(\frac{4}{25}\right)+2\left(\frac{1}{5}\right)\left(\frac{16}{125}\right)+\left(\frac{4}{25}\right)^2=\frac{113}{625}$ or 0.1808 (exact)	

x	0	1	2	3	4	8
$\mathrm{P}(X=x)$	$\frac{1}{5}$	$\frac{4}{25}$	$\frac{16}{125}$	$\frac{64}{625}$	$\frac{256}{3125}$	$\frac{1024}{3125}$

3.4 The number of muffins, X, bought by a randomly chosen customer at a muffin shop has probability distribution as shown in the table below:

x	1	2	3	4	5	6	>6
$\mathrm{P}(X=x)$	$\frac{1}{12}$	$\frac{2}{12}$	$\frac{3}{12}$	a	$\frac{2}{12}$	$\frac{1}{12}$	0

(i) Find the value of a.

(ii) State the expected value of X.

(iii) Evaluate the variance of X.

Initially, the shopkeeper charges \$2.50 for each muffin, which cost 50 cents to produce.

(iv) Using the answer to part (ii), or otherwise, find the expected profit made per customer.

The shopkeeper decides to promote the sale of muffins by giving a free muffin to all customers who buy 4 or more muffins. As a result, the probability distribution of the number of muffins, Y, taken away by a customer is given in the following table.

y	1	2	3	4	5	6	>6
$\mathrm{P}(Y=y)$	$\frac{3}{36}$	$\frac{4}{36}$	$\frac{3}{36}$	0	$\frac{20}{36}$	$\frac{6}{36}$	0

(v) Find the expected profit made per customer.

(vi) If the number of customers increases by 8%, is the promotion more profitable for the shopkeeper? Give a reason for your answer.

[SAJC DRV HW2 Review Q8]

Solution		Comments
(i)	$\frac{1}{4}$	
(ii)	3.5	
(iii)	$1\frac{11}{12}$	
(iv)	*No official answer is printed for this part.*	
(v)	\$6.86	
(vi)	*No official answer is printed for this part.*	

Only the answer line printed with the question (SAJC DRV HW2 Review Q8) is available; no official worked solution was found.

3.5 In a game, a player selects one ball at random from a bag which contains four balls numbered '1', '1', '2' and '3'. The player notes the number on the selected ball, and then tosses that number of fair coins. The number of heads obtained is denoted by X and has a probability distribution as follows:


x	0	1	2	3
$\mathrm{P}(X=x)$	a	$\frac{15}{32}$	b	$\frac{1}{32}$

(i) Find the exact values of a and b. [3]

(ii) Find $\mathrm{E}(X)$ and show that $\mathrm{Var}(X)=\frac{39}{64}$. [2]

(iii) The player plays 50 games. Find the probability that the average number of heads obtained per game exceed 1. [2]

[JPJC 2025 Prelim P2 Q7]

Solution		Comments
(i) **[3]**	$a = \mathrm{P}(X=0) = \left(\frac{1}{2}\times\frac{1}{2}\right) + \left(\frac{1}{4}\times\left(\frac{1}{2}\right)^2\right) + \left(\frac{1}{4}\times\left(\frac{1}{2}\right)^3\right) = \frac{11}{32}$ $b = \mathrm{P}(X=2) = \left(\frac{1}{4}\times\left(\frac{1}{2}\right)^2\right) + \left(\frac{1}{4}\times{}^3C_2\times\left(\frac{1}{2}\right)^3\right) = \frac{5}{32}$ Or $b = \mathrm{P}(X=2) = 1 - \frac{11}{32} - \frac{15}{32} - \frac{1}{32} = \frac{5}{32}$ $a = \frac{11}{32},\ b = \frac{5}{32}$ 	
(ii) **[2]**	Probability distribution of X: x: 0, 1, 2, 3 $\mathrm{P}(X=x)$: $\frac{11}{32}$, $\frac{15}{32}$, $\frac{5}{32}$, $\frac{1}{32}$ $\mathrm{E}(X) = \sum x\mathrm{P}(X=x) = (0)\left(\frac{11}{32}\right) + (1)\left(\frac{15}{32}\right) + (2)\left(\frac{5}{32}\right) + (3)\left(\frac{1}{32}\right) = \frac{7}{8}$ $\mathrm{E}(X^2) = \sum x^2\mathrm{P}(X=x) = (0)^2\left(\frac{11}{32}\right) + (1)^2\left(\frac{15}{32}\right) + (2)^2\left(\frac{5}{32}\right) + (3)^2\left(\frac{1}{32}\right) = \frac{11}{8}$ $\mathrm{Var}(X) = \mathrm{E}(X^2) - [\mathrm{E}(X)]^2 = \frac{11}{8} - \left(\frac{7}{8}\right)^2 = \frac{39}{64}$ (Shown)	
(iii) **[2]**	Let X = number of heads obtained in one game; $\overline{X}$ = mean number of heads obtained per game Since $n = 50$ is large, by the Central Limit Theorem, $\overline{X} \sim \mathrm{N}\left(\frac{7}{8}, \frac{39/64}{50}\right) = \mathrm{N}\left(\frac{7}{8}, \frac{39}{3200}\right)$ approximately $\mathrm{P}(\overline{X} > 1) = 0.12876 \approx 0.129$	

4 Binomial Distribution

4.1 A fruit seller sources his oranges from a supplier with a large plantation where 1% of the oranges are rotten. The supplier picks the oranges regardless of whether the oranges are rotten or not, and packs 20 oranges in each box at random.

(a) The random variable X denotes the number of rotten oranges in a randomly chosen box. State in context two assumptions for X to follow a binomial distribution. [2]

Assume that X follows a binomial distribution for the rest of this question.

The fruit seller considers a box containing between 2 and 5 rotten oranges inclusive as substandard.

(b) Find the probability that a randomly chosen box is substandard. [1]

(c) Find the expected number of rotten oranges in a box. [1]

A truckload of 10 randomly chosen boxes of oranges is then sent to the fruit seller.

(d) Find the probability that not more than 2 boxes from the entire truckload are substandard. [2]

Upon receiving the oranges, the fruit seller inspects the boxes with these conditions:

- Accept the entire truckload if the first box opened has no rotten oranges.
- Reject the entire truckload if the first box opened has more than 1 rotten orange.
- If the first box has exactly 1 rotten orange, the fruit seller opens a second box.
- If the second box has at most 1 rotten orange, he will accept the entire truckload. Otherwise, he will reject the entire truckload.

(e) Find the probability that he will reject the entire truckload. [2]

[ACJC 2025 Prelim P2 Q9]

Solution		Comments
(a) **[2]**	The probability that an orange is rotten is a constant of 0.01. Whether an orange is rotten is independent of whether any other oranges are rotten.	
(b) **[1]**	Let the random variable X denote the number of rotten oranges in a box of 20 oranges. $X \sim \mathrm{B}(20, 0.01)$ $\mathrm{P}(2 \leqslant X \leqslant 5) = \mathrm{P}(X \leqslant 5) - \mathrm{P}(X \leqslant 1) = 0.0169$	
(c) **[1]**	$\mathrm{E}(X) = (20)(0.01) = 0.2$	
(d) **[2]**	Let the random variable Y denote the number of substandard boxes out of the entire truckload of 10 boxes. $Y \sim \mathrm{B}(10, p)$ where $p = 0.0169$ $\mathrm{P}(Y \leqslant 2) = 0.999$ (3 s.f.)	
(e) **[2]**	Let A = {he rejects the entire truckload} $\mathrm{P}(A) = \mathrm{P}(X \geqslant 2) + \mathrm{P}(X = 1)\mathrm{P}(X \geqslant 2) = [1 - \mathrm{P}(X \leqslant 1)] + \mathrm{P}(X = 1)[1 - \mathrm{P}(X \leqslant 1)] = 0.0196$	

4.2 A manufacturer produces a large number of tumblers of three colours: black, pink and white in the respective ratios $3:2:1$. The tumblers are packed into boxes of 20 each.

Assume that the number of white tumblers in each box follows a binomial distribution.

(a) Find the probability that the number of white tumblers in a randomly chosen box is between 4 and 9 inclusive. [2]


(b) Find the probability that the 19th tumbler chosen from the box is the fourth white tumbler. [2]

It is known that, on average, p% of the tumblers are defective. The number of defective tumblers in a box also follows a binomial distribution. The probability that a box contains at most one defective tumbler is 0.95.

(c) Write down an equation satisfied by p. Hence find the value of p. [3]

[YIJC 2024 Prelim P2 Q8]

Solution		Comments
(a) **[2]**	Let X be the number of white tumblers in a box of 20. $X \sim \mathrm{B}(20, \frac{1}{6})$ $\mathrm{P}(4 \leqslant X \leqslant 9) = \mathrm{P}(X \leqslant 9) - \mathrm{P}(X \leqslant 3) \approx 0.433$ Alternative: $\mathrm{P}(X=4) + \mathrm{P}(X=5) + \cdots + \mathrm{P}(X=9) \approx 0.433$	
(b) **[2]**	Let W be the number of white tumblers from 18 tumblers. $W \sim \mathrm{B}(18, \frac{1}{6})$ Required probability $= \mathrm{P}(W=3) \times \frac{1}{6} \approx 0.0409$	
(c) **[3]**	Let D be the number of defective tumblers in a box of 20. $D \sim \mathrm{B}(20, 0.01p)$ $\mathrm{P}(D \leqslant 1) = 0.95$ $\mathrm{P}(D=0) + \mathrm{P}(D=1) = 0.95$ $^{20}C_0 (0.01p)^0 (1-0.01p)^{20} + {}^{20}C_1 (0.01p)^1 (1-0.01p)^{19} = 0.95$ $(1-0.01p)^{20} + 0.2p(1-0.01p)^{19} = 0.95$ Using G.C., $p \approx 1.81$	



4.3

In a computer game, a bug moves from left to right through a network of connected paths. The bug starts at S and, at each junction, randomly takes the left fork with probability p or the right fork with probability q, where $q = 1 - p$. The forks taken at each junction are independent. The bug finishes its journey at one of the 9 endpoints labelled A–I (see diagram).

(i) Show that the probability that the bug finishes its journey at D is $56p^5q^3$. [2]

(ii) Given that the probability that the bug finishes its journey at D is greater than the probability that the bug finishes its journey at any one of the other endpoints, find exactly the possible range of values of p. [4]

In another version of the game, the probability that, at each junction, the bug takes the left fork is $0.9p$, the probability that the bug takes the right fork is $0.9q$ and the probability that the bug is swallowed up by a 'black hole' is 0.1.

(iii) Find the probability that, in this version of the game, the bug reaches one of the end-points A–I, without being swallowed up by a black hole. [1]

[N2018 P2 Q6]

Solution		**Comments**
(i) **[2]**	*No official answer is printed for this part.*	
(ii) **[4]**	$\frac{5}{9} < p < \frac{2}{3}$	
(iii) **[1]**	0.430	

No official SEAB solution is published; answers as printed with this question in RI 2025 Tutorial S2B Q10.

4.4 'Crunchers' are sweets that are sold in packets of 8. Each packet is made up of randomly chosen coloured sweets. On average 15% of Crunchers are yellow.

(a) Find the probability that a randomly chosen packet of Crunchers contains no more than one yellow sweet. [2]

(b) Kev buys 70 randomly chosen packets of Crunchers. Find the probability that at least 50 of these packets contain no more than one yellow sweet. [3]

On average the proportion of Crunchers that are red is p. It is known that the modal number of red sweets in a packet is 3.

(c) Use this information to find exactly the range of values that p can take. [4]

[9758 Specimen Paper (for 2025) P2 Q7]

Solution		Comments
(a) **[2]**	Use of B(8, 0.15); 0.657	
(b) **[3]**	Use of B(70, 0.657); Use of $1-\mathrm{P}(x \leqslant 49)$; 0.189	
(c) **[4]**	$\frac{8\times7\times6}{1\times2\times3}p^3(1-p)^5 > \frac{8\times7}{1\times2}p^2(1-p)^6; \quad p > \frac{1}{3}$ $\frac{8\times7\times6}{1\times2\times3}p^3(1-p)^5 > \frac{8\times7\times6\times5}{1\times2\times3\times4}p^4(1-p)^4; \quad p < \frac{4}{9}$	

4.5 A factory produces porcelain bowls. It is known that, on average, 8% of the bowls are faulty. The bowls are packed in boxes of 20. A box is considered imperfect if it contains more than 2 faulty bowls. Assume that the number of faulty bowls in a box follows a binomial distribution.

(a) Find the probability that a randomly chosen box is imperfect. [2]

These boxes are packed into cartons of 12 boxes each.

(b) Find the probability that, out of 3 randomly chosen cartons, there are 2 cartons that contains fewer than 15 faulty bowls each. [3]

Let Y be the number of imperfect boxes in a randomly chosen carton.

(c) State the values of $\mathrm{E}(Y)$ and $\mathrm{Var}(Y)$. [2]

(d) Hence using a suitable approximation, find the probability that the total number of imperfect boxes in a random sample of 35 cartons is at most 85. [2]

[TJC 2025 Prelim P2 Q7]

Solution		Comments
(a) **[2]**	Let X be the number of faulty bowls in a box of 20 bowls. $X \sim \mathrm{B}(20, 0.08)$ $\mathrm{P}(X > 2) = 1 - \mathrm{P}(X \leqslant 2) = 1 - 0.78795 = 0.21205$ (5 s.f.) $= 0.212$ (3 s.f.)	
(b) **[3]**	Let W be the number of faulty bowls in a carton (of 12 boxes). $W \sim \mathrm{B}(12 \times 20, 0.08)$, i.e. $W \sim \mathrm{B}(240, 0.08)$ $\mathrm{P}(W < 15) = \mathrm{P}(W \leqslant 14) = 0.12933$ (5 s.f.) Required Probability $= {}^3C_2 (0.12933)^2 (1 - 0.12933) = 0.0437$ (3 s.f.) **OR** Let A be the number of cartons out of 3 contains fewer than 15 faulty bowls each. $A \sim \mathrm{B}(3, 0.12933)$ Required Probability $= \mathrm{P}(A = 2) = 0.0437$ (3 s.f.)	
(c) **[2]**	$Y \sim \mathrm{B}(12, 0.21205)$ $\mathrm{E}(Y) = np = 12 \times 0.21205 = 2.5446 = 2.54$ (3 s.f.) $\mathrm{Var}(Y) = np(1-p) = 12 \times 0.21205 \times (1 - 0.21205) = 2.0050$ (5 s.f.) $= 2.01$ (3s.f.)	
(d) **[2]**	$Y \sim \mathrm{B}(12, 0.21205)$. Let $T = Y_1 + \cdots + Y_{35}$. Since sample size 35 is large, by Central Limit Theorem, $T \sim \mathrm{N}(35 \times 2.5446,\ 35 \times 2.0050)$ approximately i.e. $T \sim \mathrm{N}(89.061,\ 70.175)$ approximately Required probability $= \mathrm{P}(T \leqslant 85) = 0.314$ (3 s.f.)	

5 Normal Distribution

5.1 In this question, you should state clearly all the distributions that you use, together with the values of the appropriate parameters.

A toy designer creates weighted wooden blocks used in balance-based educational toys. There are two types of blocks: cylindrical blocks and cuboidal blocks. The mass, in grams, of each cylindrical block and cuboidal block follow a normal distribution with the following parameters:

	Mean	Standard Deviation
cylindrical block	280	6
cuboidal block	220	5

(a) Find the probability that the mass of a randomly selected cylindrical block exceeds the mass of a randomly selected cuboidal block by more than 65 grams. [2]

(b) The probability that the total of twice the mass of a cylindrical block and thrice the mass of a cuboidal block being less than M g is 0.355. Find the value of M, giving your answer to the nearest integer. [2]

To build a deluxe seesaw model, the designer uses:

- 2 cylindrical blocks,
- 3 cuboidal blocks,
- 7 screws,
- 1 wooden plank.

The blocks are drilled before assembly to fit screws:

- Each cylindrical block loses 5% of its mass after drilling.
- Each cuboidal block loses 3% of its mass after drilling.

The masses of the remaining components follow a normal distribution with the following parameters:

	Mean	Standard Deviation
screw	8	1.2
wooden plank	540	8

When assembling the deluxe seesaw model, all of its components are chosen at random.

(c) Find the probability that the total mass of all the modified components in a deluxe seesaw model is more than 1785 g. [4]

[ASRJC 2025 Prelim P2 Q9]

Solution		Comments
(a) **[2]**	Let X and Y be the mass of a randomly selected cylindrical and cuboidal block (in g) respectively. $X \sim \mathrm{N}(280,\ 6^2)$ and $Y \sim \mathrm{N}(220,\ 5^2)$ $X-Y \sim \mathrm{N}(280-220,\ 5^2+6^2)$ $X-Y \sim \mathrm{N}(60,\ 61)$ $\mathrm{P}(X-Y>65)=0.261$	
(b) **[2]**	Let $C=2X+3Y$ $C \sim \mathrm{N}(2\times280+3\times220,\ 2^2\times6^2+3^2\times5^2)$ $C \sim \mathrm{N}(1220,\ 369)$ Given $\mathrm{P}(C<M)=0.355$ From GC, $M=1212.8568$ $M=1213$	
(c) **[4]**	Let S and W be the mass of a randomly selected screw and wooden plank (in g) respectively. $S \sim \mathrm{N}(8,\ 1.2^2)$ and $W \sim \mathrm{N}(540,\ 8^2)$ Let $T=0.95(X_1+X_2)+0.97(Y_1+Y_2+Y_3)+S_1+\cdots+S_7+W$ $\mathrm{E}(T)=(0.95\times2\times280)+(0.97\times3\times220)+(7\times8)+540$ $=532+426.8+56+540$ $=1768.2$ $\mathrm{Var}(T)=(0.95^2\times2\times6^2)+(0.97^2\times3\times5^2)+(7\times1.2^2)+64$ $=64.98+70.5675+10.08+64$ $=209.6275$ $\mathrm{P}(T>1785)=0.123$	

5.2 In this question you should state the parameters of any distribution you use.

An electric power service company keeps records of the installation time of its electricity meters in the houses in a new estate. The time taken to install an electricity meter is normally distributed with mean 45 minutes and standard deviation 6 minutes.


(a) Sketch the distribution for the installation time of an electricity meter between 20 minutes and 70 minutes. [2]

(b) A house that has its electricity meter take more than 1 hour to be installed is considered 'inefficient'. The company randomly selects n houses in the estate for quality control in their services. Find the greatest value of n such that the probability that fewer than 3 of these n houses are 'inefficient' is at least 0.90. [3]

(c) Each month, the amount of electricity, in kilowatt-hours (kWh), used by a particular household in the estate has the distribution $\mathrm{N}(524, 27^2)$. The company charges households for electricity used at \$0.26 per kWh and each household is billed every two months. Find the probability that a randomly chosen bill for this household is more than \$270 given that it is between \$250 and \$280. State an assumption that is needed for your calculations to be valid. [4]

(d) The company also installs gas meters in the houses in the estate. The time taken in minutes to install a gas meter follows a normal distribution. 40% of the gas meters each has an installation time greater than 53 minutes and 15% of the gas meters each has an installation time less than 38 minutes. Find the mean and standard deviation of the installation times of the gas meters in the estate. [4]

[NJC 2024 Prelim P2 Q12]

Solution		**Comments**
(a) **[2]**	Let X denote the random variable 'time taken (min) to install an electricity meter.' $X \sim \mathrm{N}(45, 6^2)$  Note: $\mathrm{P}(20 < X < 70) = 0.99997$	

(b) **[3]**

Let E be the random variable 'number of 'inefficient' houses out of n houses.'

$E \sim \mathrm{B}(n,\ \mathrm{P}(X>60))$

$E \sim \mathrm{B}(n,\ 0.0062097)$

$\mathrm{P}(E<3) = \mathrm{P}(E \le 2) \ge 0.90$

Using GC,

n	$\mathrm{P}(E \le 2)$
177	$0.9012 \ge 0.9$
178	$0.8999 < 0.9$
179	0.8986

Largest $n = 177$

(c) **[4]**

Let W be the random variable 'amount of electricity (kWh) used in the household in a month.'

$W \sim \mathrm{N}(524,\ 27^2)$

Let $T = 0.26(W_1 + W_2)$.

$\mathrm{E}(T) = 0.26 \times 524 \times 2 = 272.48$

$\mathrm{Var}(T) = 0.26^2 \times 2 \times 27^2 = 98.5608$

$T \sim \mathrm{N}(272.48,\ 98.5608)$

$$\mathrm{P}(T>270 \mid 250<T<280) = \frac{\mathrm{P}(270<T<280)}{\mathrm{P}(250<T<280)}$$
$$= \frac{0.36425}{0.76383}$$
$$= 0.490 \ \text{(3 s.f.)}$$

We need to assume that the electricity used in each of these two months is independent for this particular household.

(d) **[4]**

Let G denote the random variable 'time taken (min) to install a gas meter

$G \sim \mathrm{N}(\mu, \sigma^2)$.

$\mathrm{P}(G<38) = 0.15, \qquad \mathrm{P}(G>53) = 0.4$

$$\mathrm{P}\left(Z < \frac{38-\mu}{\sigma}\right) = 0.15, \qquad \mathrm{P}\left(Z > \frac{53-\mu}{\sigma}\right) = 0.4$$

By G.C.,

$$\frac{38-\mu}{\sigma} = -1.0364, \qquad \frac{53-\mu}{\sigma} = 0.25335$$

$\mu - 1.0364\sigma = 38 \quad ---(1), \qquad \mu + 0.25335\sigma = 53 \quad ---(2)$

Solving (1) and (2):

$\mu = 50.0535 = 50.1$ (3 s.f.)



$\sigma = 11.6298 = 11.6$ (3 s.f.)

5.3 The waiting times for customers to get their pastries from a popular bakery is modelled by a normal distribution with mean μ minutes and standard deviation σ minutes. On average, 10% of customers wait for more than 25 minutes and the same percentage of customers wait for less than 5 minutes.

(i) Find the values of μ and σ, leaving your answers to the nearest minute. [3]

(ii) Explain if the normal distribution is a suitable model for the waiting times for customers to get pastries from this bakery. [1]

[MI 2025 Prelim P2 Q6]

Solution		**Comments**
(i) **[3]**	Let X be the waiting time to get pastries from the popular bakery of a randomly chosen customer. $X \sim \mathrm{N}(\mu,\ \sigma^2)$ Given $\mathrm{P}(X<5)=\mathrm{P}(X>25)=0.10$. **Method 1** By symmetry, $\mu=\dfrac{5+25}{2}=15$  $\mathrm{P}(X>25)=0.10$ $\mathrm{P}\left(Z>\dfrac{25-15}{\sigma}\right)=0.10$ $\dfrac{10}{\sigma}=1.28155$ $\sigma=7.80305$ --- ***Alternatively,*** $\mathrm{P}(X>25)=0.10$ Using GC,  $\sigma=7.80304$ --- $\therefore \mu=15,\ \sigma=8$ (to the nearest minute).	

	Method 2 $\mathrm{P}(X>25)=0.10$ $\quad$ $\mathrm{P}(X<5)=0.10$ $\mathrm{P}\left(Z>\frac{25-\mu}{\sigma}\right)=0.10$ $\quad$ $\mathrm{P}\left(Z<\frac{5-\mu}{\sigma}\right)=0.10$ $\frac{25-\mu}{\sigma}=1.28155$ $\quad$ $\frac{5-\mu}{\sigma}=-1.28155$ $\mu+1.28155\sigma=25$ … (1) $\quad$ $\mu-1.28155\sigma=5$ … (2) Solving (1) and (2) simultaneously, $\mu=15,\ \sigma=7.80305.$ $\therefore\mu=15,\ \sigma=8$ (to the nearest minute).	
(ii) **[1]**	Consider $X\sim\mathrm{N}(15,\ 8^2)$. $\mathrm{P}(X<0)=0.0303.$ (or 0.02728 if we use $\sigma=7.80305$) The waiting times cannot be negative but $\mathrm{P}(X<0)$ is about 3%, which is not insignificant. Thus, the normal distribution is not a suitable model. **Note:** For normal distribution: $\mathrm{P}(\mu-3\sigma<X<\mu-3\sigma)\approx 0.997$ i.e. 99.7%. Hence, whenever the probability of obtaining unreasonable values of x exceeds 0.003, we can conclude that the normal distribution is not suitable	

5.4 A factory produces metal discs and metal bars for constructing machinery parts. The mass of a randomly chosen metal disc follows a normal distribution with mean 18 grams and standard deviation 0.6 grams. The mass of a randomly chosen metal bar has an independent normal distribution with mean 20 grams and standard deviation σ grams.

(a) Sketch the distribution for the masses of the metal disc between 14 grams and 22 grams. [1]

(b) 2 metal discs are randomly chosen. Find the probability that the mass of one of the discs is less than 19 grams and the other metal disc has mass more than 19 grams. [2]

(c) Find the range of values of k such that less than 90% of the metal discs have masses within k grams of the mean mass of the metal discs. [2]

(d) It is known that 7.65% of the metal bars have masses more than 21 grams. Find the value of σ. [2]

It is given that $\sigma = 0.5$ for the rest of the question.


(e) A client places an order for a new kind of metal strips. The metal strips are modelled as similar solids to the metal bars with a 20% reduction in mass.

A production worker selects 8 of the metal strips for inspection. Find the probability that at least 5 of the metal strips have masses between 15.8 grams and 17.2 grams. [3]


(f) A protective case is designed for the safe transportation of the metal bar. Each metal bar is placed inside a protective case during transportation. The masses of the protective cases are modelled as 30% of the masses of the metal bars.

Let T denote the total mass of 4 of the metal bars and their protective cases. Find the variance of T. [2]

[RVHS 2025 Prelim P2 Q12]

Solution		Comments
(a) **[1]**		

(b) **[2]**	Let D be the random variable denoting the mass of a randomly chosen metal disc. $D \sim N(18,\ 0.6^2)$ Probability required $= \binom{2}{1} [P(D<19)]\,[P(D>19)] - - - (*)$ $= 0.0910128294$ ≈ 0.0910 (3 s.f.)	
(c) **[2]**	$P(18-k<D<18+k)<0.9$ Since $P(17.01308782<D<18.98691218)=0.9$, then $18+k<18.98691218$ $k<0.987$	
(d) **[2]**	7.65% of the metal bars have masses more than 21 grams. Let B be the random variable denoting the mass of a randomly chosen metal bar. $B \sim N(20, \sigma^2)$ $P(B>21)=0.0765$ $P\left(Z>\frac{21-20}{\sigma}\right)=0.0765$ $P\left(Z>\frac{1}{\sigma}\right)=0.0765$ $\frac{1}{\sigma}=1.429014729$ $\sigma=0.6997$ $\sigma=0.700$ (3.s.f)	

(e) **[3]**	Let S be the random variable denoting the mass of a randomly chosen metal strip. $S = 0.8B \sim N(0.8 \times 20,\ 0.8^2 \times 0.5^2)$ $S \sim N(16,\ 0.4^2)$ $P(15.8 < S < 17.2) = 0.690125006$ Let X be the random variable denoting the number of metal strips whose mass lies between 15.8 grams and 17.2 grams, out of 8 bars. $X \sim B(8,\ 0.690125006)$ $P(X \geq 5)$ $= 1 - P(X \leq 4)$ $= 0.7875986163$ ≈ 0.788 (3.s.f)  8 strips, each "success" if $15.8 < S < 17.2$ (prob. 0.690125006): $X \sim B(8,\ 0.690125006)$, need $X \geq 5$	

(f) **[2]**	Let C be the random variable denoting the mass of a randomly chosen protective case. $C \sim N(0.3\times 20,\ 0.3^2\times 0.5^2)$ $C \sim N(6,\ 0.15^2)$ $B \sim N(20,\ 0.5^2)$ Let $Y = B + C$ $Y \sim N(20+6,\ 0.5^2+0.15^2)$ $Y \sim N(26,\ 0.2725)$ Let $T = Y_1 + Y_2 + Y_3 + Y_4$ $\text{Var}(T) = 4\times 0.2725$ $= 1.09$ (3 s.f) **Common Mistake** Let C be the random variable denoting the mass of a randomly chosen protective case. $C = 0.3B$ $B \sim N(20,\ 0.5^2)$ Let $Y = B + C$ $Y = B + 0.3B = 1.3B$ $Y \sim N(1.3\times 20,\ 1.3^2\times 0.5^2)$ $Y \sim N(26,\ 0.4225)$ Let $T = Y_1 + Y_2 + Y_3 + Y_4$ $T \sim N(4\times 26,\ 4\times 0.4225)$ $T \sim N(104,\ 1.69)$	

6 Hypothesis Testing

6.1 Energy bars are a convenient source of nutrition for athletes. A typical energy bar weighs between 30-50 g and is likely to supply about 200-300 calories. An American company sells energy bars and a nutritionist working in the company claims that the mean number of calories in a product, Perfect Protein bars, is 350 calories.

A random sample of 30 bars is collected and the number of calories, x cal, in the 30 bars is summarised as follows.

$$\sum x = 10446, \qquad \sum x^2 = 3638000.$$

(i) Test at the 3% level of significance whether the mean number of calories in Perfect Protein bars is less than 350 calories, defining any symbols that you use. [7]


(ii) Explain why the test would be inadmissible if the nutritionist had taken a random sample of 15 energy bars and hence state an assumption needed for him to carry out the test. [1]


(iii) If the test was for whether the mean number of calories in Perfect Protein bars is 350 calories, find the smallest level of significance for which the nutritionist's claim will be rejected. [2]

The company launched a new product called Dynamic Protein bars which are advertised to be higher in calories than Perfect Protein bars. A random sample of 45 Dynamic Protein bars is collected to test this and the mean number of calories of the sample is found to be 352.2.

(iv) The result of this investigation shows that there is no reason to believe that the mean number of calories in the new product is more than 350 calories at the 4% level of significance. Find the range of possible values of the variance used in calculating the test statistic. [3]

[ACJC 2023 Prelim P2 Q11]

Solution		Comments
(i) **[7]**	$\bar{x} = \dfrac{\sum x}{n} = \dfrac{10446}{30} = 348.2$ (exact) $s^2 = \dfrac{1}{n-1}\left(\sum x^2 - \dfrac{\left(\sum x\right)^2}{n}\right) = \dfrac{1}{29}\left(3638000 - \dfrac{(10446)^2}{30}\right) =$ 24.23448 Let μ be the population mean number of calories in Perfect Protein bars. Null hypothesis $H_0 : \mu = 350$ Alternative hypothesis $H_1 : \mu < 350$ Perform 1-tail test at 3 % significance level. Under H_0, $\bar{X} \sim N\left(350, \dfrac{24.23448}{30}\right)$ by Central Limit Theorem since $n = 30$ is large enough. From GC, $z = \dfrac{\bar{x} - \mu_0}{\sqrt{\frac{s^2}{n}}} = \dfrac{348.2 - 350}{\sqrt{\frac{24.23448}{30}}} = -2.0027$ p-value $= P(Z < -2.0027) = 0.0226$  p-value < 0.03 ∴Reject H_0 There is sufficient evidence at 3% level of significance to conclude that the mean number of calories in Perfect Protein bars is less than 350 cal.	
(ii) **[1]**	The test would be inadmissible if the nutritionist had taken a random sample of 15 energy bars as the distribution of the population is unknown (and Central Limit Theorem cannot be applied). The nutritionist will need to assume that the number of calories in Perfect Protein bars follow a normal distribution.	
(iii) **[2]**	Null hypothesis $H_0 : \mu = 350$ Alternative hypothesis $H_1 : \mu \neq 350$ Perform 2-tail test at α % significance level. p-value $= 2 \times P(Z < -2.0027) = 0.0452$ To reject H_0, p-value $= 0.045209 < \alpha\%$ ∴ smallest significance level is 4.52%.	

(iv) **[3]**	Let μ be the population mean number of calories in Dyanmic Protein bars. Null hypothesis $H_0: \mu = 350$ Alternative hypothesis $H_1: \mu > 350$ At 4% level of significance, we do not reject H_0. For p-value > 0.04 or $z < 1.750686$, we do not reject H_0. $z = \dfrac{\bar{x} - \mu_0}{\sqrt{\frac{\sigma^2}{n}}} = \dfrac{352.2 - 350}{\sqrt{\frac{\sigma^2}{45}}} < 1.750686$  $1.750686\sqrt{\dfrac{\sigma^2}{45}} > 2.2$ $\sigma^2 > 45\left(\dfrac{2.2}{1.750686}\right)^2$ $\sigma^2 > 71.0626$ $\therefore \sigma^2 > 71.1$ (3 sf.)	

6.2 A hospital introduces a new medication regimen to help patients recover from a viral infection more quickly. Previously, the average recovery time was 7.2 days.

After adopting the new regimen, a random sample of 40 patients is observed, and their recovery times, x, in days, are recorded. The results are summarised below.

$$\sum(x-7.2)=-42 \quad \text{and} \quad \sum(x-7.2)^2=648$$

(a) Explain whether the hospital should carry out a one-tailed test or a two-tailed test. [1]

(b) Calculate unbiased estimates of the population mean and variance of the recovery times under the new regimen. [2]

(c) Carry out an appropriate test at the 5% significance level. You should state clearly the hypotheses for your test and define any parameters that you use. [4]

(d) Explain, in the context of the question, the meaning of 'at the 5% significance level'. [1]

(e) Explain whether it would have been sufficient for the hospital to take a random sample of 10 patients and record their recovery times under the new regimen in order to carry out the hypothesis test. [1]

It was discovered that an intern at the hospital had made an error in recording the recovery times. A random sample of another 30 patients was taken and their recovery times under the new regimen was taken and recorded by a senior researcher. The sample standard deviation was found to be 1.7 days and the sample mean time was denoted by k.

(f) A test at the 1% significance level concludes that the average recovery time did not improve from 7.2 days. Find the set of values of k. [3]

[VJC 2025 Prelim P2 Q12]

Solution		Comments
(a) **[1]**	The researchers should carry out a one-tailed test as they want to test whether the new regimen has **reduced mean recovery time.**	

(b) **[2]**

Let $y = x - 7.2$

Unbiased estimate of population mean, $\bar{x} = \bar{y} + 7.2 = \frac{-42}{40} + 7.2 = 6.15$

Unbiased estimate of population variance,

$s_x^2 = s_y^2 = \frac{1}{39}\left(648 - \frac{(-42)^2}{40}\right) = \frac{603.9}{39} = 15.4846 \approx 15.5$

(c) **[4]**

Let X days be the recovery time of a patient and μ days be the population mean of X.

$H_0 : \mu = 7.2$

$H_1 : \mu < 7.2$


Level of Significance: 5%

Test Statistic:

Since $n = 40$ is large, by Central Limit Theorem, $\bar{X}$ is approximately normal.

When H_0 is true, $Z = \frac{\bar{X} - 7.2}{S/\sqrt{n}} \sim N(0, 1)$ approximately.

Computation: $\bar{x} = 6.15$, $s_x^2 \approx 15.4846$, p-value $= 0.045744$ (or $z = -1.6876$)




Conclusion: Since p-value $= 0.0457 < 0.05$, H_0 is rejected at 5% level of significance. Hence there is sufficient evidence to conclude that the mean recovery time has reduced under the new regimen.

(d) **[1]**

A 5% significance level means that there is a probability of 0.05 that the test concludes that the mean recovery time has reduced when it is actually 7.2 days.

(e) **[1]**

No. Since the distribution of recovery time under the new regimen (X) is unknown, the sample size would have to be sufficiently large for Central Limit Theorem to be applied so that the **sample mean recovery time** is approximately normally distributed. A sample of 10 is not large enough.

(f) **[3]**	$H_0 : \mu = 7.2$ $H_1 : \mu < 7.2$ Level of Significance: 1% Test Statistic: Since $n = 30$ is large, by Central Limit Theorem, $\bar{X}$ is approximately normal. When H_0 is true, $Z = \frac{\bar{X} - 7.2}{S/\sqrt{n}} \sim N(0, 1)$ approximately. Computation: $s^2 = \frac{n}{n-1}(\text{sample variance}) = \frac{30}{29}(1.7)^2 = 2.9897$ Rejection Region: $z \leq -2.3263$  Since H_0 is not rejected, $\frac{k - 7.2}{\sqrt{2.9897}/\sqrt{30}} > -2.3263$ $k > 6.4656$ $\therefore \{k \in \mathbb{R} : k > 6.47\}$	

6.3 In a drinks factory, a machine is programmed to dispense 500 ml of green tea into empty bottles.

(a) After a routine check of the machine, the production manager suspects that the machine is dispensing more green tea than expected. A random sample of 50 bottles is taken for a hypothesis test at α% level of significance and the data is as follows:

Volume of green tea (correct to nearest ml)	498	499	500	501	502	503	504
Number of bottles	5	11	10	11	10	1	2





Taking $\alpha = 2$, test the production manager's suspicion and state the meaning of the p-value in the context of the question. [5]

(i) Determine the set of values of α for which the production manager's suspicion is valid. [1]

(ii) Explain why the production manager is able to carry out the hypothesis test without knowing anything about the distribution of the volume of green tea dispensed by the machine. [1]

(b) The machine is being recalibrated to dispense 500 ml of green tea. Another random sample of 50 bottles of green tea is taken and the mean and standard deviation of this sample are 502 ml and k ml respectively. A hypothesis test, at the 5% significance level, concluded that the recalibration is done accurately. Find the set of values that k can take. [5]

[JPJC 2024 Prelim P2 Q10]

Solution		**Comments**
(a) **[5]**	   Using GC, $\bar{x} = 500.42$ (exact) $s = 1.5265$ (to 5 s.f.)   	

Let X = vol. of green tea dispensed in a bottle and

μ = population mean green tea dispensed in a bottle

$H_0 : \mu = 500$

$H_1 : \mu > 500$


Under H_0 ,since n = 50 is large, by Central Limit Theorem,

$$\bar{X} \sim N\left(500\ ,\ \frac{1.5265^2}{50}\right) \text{ approximately}$$

Test statistic: $z = \dfrac{\bar{x} - 500}{\left(\dfrac{1.5265}{\sqrt{50}}\right)}$

$$= \frac{500.42 - 500}{\left(\dfrac{1.5265}{\sqrt{50}}\right)} = 1.9455 \approx 1.95 \text{ (to 3 s.f)}$$

From GC, p-value = $0.025856 \approx 0.0259$ (to 3 s.f)



Since p-value = 0.0259 > 0.02, we do not reject Ho at 2% level of significance and conclude that there is insufficient evidence that the production manager's suspicion is valid.

The p-value is 0.0259 and it means there is a probability of 0.0259 of observing a test statistic, $z \geq 1.95$, given that the population mean green tea dispensed in a bottle is 500ml.

(i) **[1]**

For the production manager's suspicion ($\mu > 500$) to be valid, H_0 is rejected. Hence,

p-value $= 0.025856 < \dfrac{\alpha}{100}$

$\therefore 2.59 < \alpha < 100$ (3 s.f)

(ii) **[1]**

Since the sample size = 50 is large enough, Central Limit Theorem can be applied for sample means to follow a normal distribution approximately.

(b) **[5]**

$s^2 = \dfrac{n}{n-1}\sigma_x^2 = \dfrac{50}{49}k^2$

$H_0 : \mu = 500$


$H_1 : \mu \neq 500$

Under H_0 ,since n = 50 is large, by Central Limit Theorem,

$$\bar{X} \sim N\left(500\ ,\ \frac{\left(\frac{50k^2}{49}\right)}{50}\right) = N\left(500\ ,\ \frac{k^2}{49}\right) \text{ approximately}$$

Test statistic: $z = \dfrac{\bar{x} - 500}{\sqrt{\frac{k^2}{49}}} = \dfrac{502 - 500}{\frac{k}{7}} = \dfrac{14}{k}$

For α = 0.05,



If recalibration is done accurately ($\mu = 500$), H_0 is not rejected. Hence,

$-1.95996 < z < 1.95996$

$-1.95996 < \dfrac{14}{k} < 1.95996$

$\dfrac{k}{14} < -0.51021$ (rejected) or $\dfrac{k}{14} > 0.51021$

$k > 7.1430$

$\therefore k > 7.14$ (to 3 s.f)

[Note: the official solution labels the parts 10(i), (ii), (iii), (b); they are shown here as (a), (i), (ii), (b) to match the question's parts. The calculator screens above are the official screenshots printed in the solution.]

6.4 EastHam is a football club in a competitive league. It is found that the average time taken by EastHam to score their first goal in a match is 50 minutes. The coach trialled a new play tactic and the time taken, x minutes, to score the first goal is recorded for a random sample of 35 matches in the season. The total time taken is found to be 1650 minutes and the variance of the time is 250 minutes^2.

The coach wants to test if the average time taken to score the first goal has decreased with the implementation of the new play tactic.

(a) Explain why the coach is able to carry out a hypothesis test without any assumption about the distribution of the time taken to score the first goal. [1]

(b) Find unbiased estimates of the population mean and variance. Give your answers correct to 3 decimal places. [2]


(c) Carry out the test at 10% level of significance. You should state your hypotheses and define any symbols you use. [4]


A fan of another football club in the same league, NewPalace, claims that the average time taken by NewPalace to score their first goal in a match is 40 minutes. The time taken by NewPalace to score the first goal in a match can be assumed to be normally distributed with variance 280 minutes^2.

(d) A random sample of 50 matches is taken and a hypothesis test is carried out on whether the fan's claim is valid. Find the range of values of the mean time of this sample for which the fan's claim would be rejected at 10% level of significance. [4]

[CJC 2025 Prelim P2 Q11]

Solution		**Comments**
(a) **[1]**	The sample mean time for 35 matches is approximated to have a normal distribution by the Central Limit Theorem, since sample size is sufficiently large. Hence there is no need to make any assumptions about the distribution of the time taken to score the first goal.	
(b) **[2]**	Unbiased estimate for population mean $= \frac{1650}{35} = 47.143$ (to 3d.p.) Unbiased estimate for population variance is $s^2 = \frac{35}{34}(250) = 257.353$ (to 3d.p.)	

(c) **[4]**	Let X be the random variable denoting time taken by EastHam to score the first goal in a match. $H_0 : \mu = 50$ $H_1 : \mu < 50$, where μ is the population mean time taken by EastHam to score the first goal in a match Under H_0, Since n is large, $\bar{X} \sim N\left(50, \frac{257.353}{35}\right)$ approximately by CLT $Z = \frac{\bar{X} - 50}{\sqrt{s^2/n}} \sim N(0, 1)$ Method ①: By GC, p-value $= 0.146$ (3 s.f.). Since p-value > 0.1, we do not reject H_0 and conclude that there is insufficient evidence at 10% significance level to say that average time taken by EastHam to score the first goal has decreased. Method ②: $z_{\text{test}} = \frac{\left(\frac{1650}{35}\right) - 50}{\sqrt{\frac{4375/17}{35}}} = -1.05366, \ z_{\text{test}} = -1.28155$  Since $z_{\text{test}} > z_{\text{critical}}$, we do not reject H_0 and conclude that there is insufficient evidence at 10% significance level to say that average time taken by EastHam to score the first goal has decreased.	

(d) **[4]**	Let Y be the random variable denoting time taken by NewPalace to score the first goal in a match. $H_0 : \mu = 40$ $H_1 : \mu \neq 40$ Under H_0, $\bar{Y} \sim N\left(40, \frac{280}{50}\right)$  Since H_0 is rejected, $\frac{\bar{y}-40}{\sqrt{280/50}} < -1.6449$ or $\frac{\bar{y}-40}{\sqrt{280/50}} > 1.6449$ $\bar{y} < 36.1$ (3 s.f.) or $\bar{y} > 43.9$ (3 s.f.) Hence, range of values of $\bar{y}$ is: $0 < \bar{y} < 36.1$ or $\bar{y} > 43.9$ [Note: the official answer key also contains Examiners' Feedback boxes; only the worked solutions are reproduced here. The second "$z_{test} = -1.28155$" is printed as such in the official solution (it is the critical value).]	

6.5 A food producer claims that the mean mass of a can of beans it produces is 425 g. Following customer feedback, the production manager wishes to test if the mean mass of a can of beans is indeed 425 g.

The production manager took a random sample of size 50 and the mass of each can, in x g, is recorded and the results are shown below:

$$\sum x = 21209, \quad \sum (x - 424.18)^2 = 522$$

(a) State what it means for a sample to be random in this context. [1]

(b) Find the unbiased estimates for the population mean and variance. [2]

(c) State the hypotheses for the manager's test, defining any parameters you use. Carry out the test at the 5% level of significance, giving your conclusion in the context of the question. [5]

The production manager wishes to test whether the mean mass of a can of beans has increased using the alternative manufacturing process. He finds that the mean mass of 55 randomly chosen cans is 426.5 g. He carries out a hypothesis test at 10% level of significance.

(d) Explain, with justification, how the population standard deviation of the mass of a can produced under the alternative process will affect the conclusion made by the production manager. [3]

[HCI 2025 Prelim P2 Q8]

Solution		**Comments**
(a) **[1]**	Every can in the population has an equal chance of being selected and is selected independently of the others.	
(b) **[2]**	Unbiased estimate for the population mean $\bar{x} = \frac{\sum x}{50} = \frac{21209}{50} = 424.18$ Unbiased estimate for the population variance $s^2 = \frac{50}{50-1} \frac{\sum (x - \bar{x})^2}{50}$ $= \frac{\sum (x - 424.18)^2}{49}$ $= \frac{522}{49}$ or 10.65306122	

(c) **[5]**	Let X be the mass (in grams) of a can of beans. Let μ and σ^2 be the population mean and variance of X. $H_0 : \mu = 425$ $H_1 : \mu \neq 425$ $n = 50, \bar{x} = 424.18, s^2 = \frac{522}{49}$ Under H_0, since $n = 50$ is large, by the Central Limit Theorem , $\bar{X} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$ approximately. Test statistic $Z = \frac{\bar{X} - \mu}{\sqrt{\frac{S^2}{n}}} \sim N(0, 1)$ approximately Level of significance: 5% Reject H_0 if p-value ≤ 0.05 Assuming H_0 is true, by G.C., p-value = **0.0757** (3 s.f.) Since p-value $= 0.0757 > 0.05$, we do not reject H_0 at 5% level of significance and conclude that there is insufficient evidence to say that the population mean mass of a can of beans is not 425 g.	

(d)
[3]

Let Y be the mass (in grams) of a can of beans produced using the alternative process.

Let μ and σ^2 be the population mean and variance of Y.

$H_0 : \mu = 425$

$H_1 : \mu > 425$


$n = 55, \bar{y} = 426.5$

Under H_0, since $n = 55$ is large, by the Central Limit Theorem, $\bar{Y} \sim N\left(\mu, \frac{\sigma^2}{n}\right)$ approximately.

Test statistic $Z = \frac{\bar{Y} - \mu}{\sqrt{\frac{\sigma^2}{n}}} \sim N(0, 1)$ approximately

Level of significance: 10%

Reject H_0 if $z \geq 1.281551567$



$$\frac{426.5 - 425}{\sqrt{\frac{\sigma^2}{55}}} \geq 1.281551567$$

Since $\sigma \geq 0$, $\sqrt{\sigma^2} \leq \frac{1.5\sqrt{55}}{1.281551567}$

$\Leftrightarrow \sigma \leq 8.680335632$

$\Leftrightarrow \sigma \leq 8.68$ (3 s.f.)

$\Leftrightarrow 0 < \sigma \leq 8.68$ (3 s.f.)

At 10% level of significance, if $0 < \sigma \leq 8.68$, then H_0 is rejected and the production manager may conclude that mean mass of a can of beans produced using the alternative process has increased.

7 Correlation and Linear Regression

7.1 An energy company is studying the relationship between the average daily temperature, x (in degrees Celsius), and the amount of electricity consumed, y (in megawatt-hours MWh). For a random sample of ten days, the records are shown in the table below.

Average daily temperature (x°C)	20	22	24	26	28	30	32	34	36	38
Electricity consumption (y MWh)	12.5	12.8	13.2	14.0	14.7	16.2	16.2	19.9	22.3	29.5

(a) Draw a scatter diagram for these values, labelling the axes clearly. [1]

It is thought that the electricity consumption and average daily temperature can be modelled by one of the formulae

$$y=a+bx^2, \qquad y=c+dx,$$

where a, b, c and d are constants.

(b) Find the value of the product moment correlation coefficient between

(i) x and y,

(ii) x^2 and y. [2]


(c) Using your answers in parts **(a)** and **(b)**, explain which of $y=a+bx^2$ and $y=c+dx$ is the better model and find the equation of a suitable regression line for this model. [3]

(d) Use the equation of your regression line to estimate the electricity consumption on a day with an average temperature of 37°C. Comment on the reliability of your estimate. [2]

In some regions, temperature is measured in degrees Fahrenheit instead of degrees Celsius. A temperature of C degrees Celsius is equivalent to F degrees Fahrenheit, where $F=\frac{9}{5}C+32$.

(e) Rewrite the equation of your regression line found in part **(c)** in terms of y and F, where F is the average daily temperature in degrees Fahrenheit. [1]

[VJC 2025 Prelim P2 Q11]

Solution		Comments
(a) **[1]**		
(b)(i)	From GC, $r=0.89056\approx 0.891$.	
(b)(ii)	From GC, $r=0.92229\approx 0.922$.	
(c) **[3]**	From the scatter diagram, it is observed that as x increases, y increases by increasing amounts, and the product moment correlation coefficient between x^2 and y is 0.922, which is closer to 1 than that between x and y, which is 0.891 . Hence $y=a+bx^2$ is the better model. $a=4.8291\approx 4.83$ $b=0.014074\approx 0.0141$ $y=4.83+0.0141x^2$	
(d) **[2]**	$y=4.8291+0.014074(37)^2$ ≈ 24.1 Since $x=37$ lies within the given range of x values, $20\le x\le 38$, and the value of $r\approx 0.922$ is close to 1, the estimated electricity consumption is reliable.	
(e) **[1]**	$F=\frac{9}{5}x+32\Rightarrow x=\frac{5}{9}(F-32)$ Replace x by $\frac{5}{9}(F-32)$: $y=4.8291+0.014074\left[\frac{5}{9}(F-32)\right]^2$ $=4.8291+0.0043438(F-32)^2$ $\approx 4.83+0.00434(F-32)^2$	

7.2 The yield per hectare, y kg, of a crop is believed to depend on the average rainfall, x mm, in the month of June. For 10 regions, records are kept of the values of x and y, and these are shown in the table below. The yield from the tenth region was accidentally deleted from the records after the data was analysed, and this is indicated by the value p.

Average rainfall (x mm)	149	110	188	135	156	140	168	118	122	174
Yield of crop (y kg)	13.8	6.5	15.2	12.2	14.4	12.2	14.7	9.5	9.9	p

Given that the equation of the regression line of y on x is $y=-2.5652+0.10168x$, show that $p=14.4$. [2]

(a) Draw a scatter diagram for these values, labelling the axes clearly. Calculate the product moment correlation coefficient between x and y. [2]

(b) It is thought that a model of the form $y=a+b\ln x$ may also be a suitable fit to the data. Calculate least square estimates of a and b, and find the value of the product moment correlation coefficient between y and $\ln x$. [3]

(c) Use your answers to parts **(a)** and **(b)** to explain which of


$$y=-2.5652+0.10168x \text{ or } y=a+b\ln x$$

is the better model. [2]

(d) Using an appropriate regression line, estimate the yield for a region that experienced 200 mm of rainfall in June. Comment on the reliability of your estimate. [2]

(e) In some regions, rainfall is measured in inches instead of in mm. Given that there are 25.4 mm in an inch, show how the regression line found in part **(b)** can be re-written so that it can be used when x, the average rainfall in June, is given in inches. [1]

[VJC 2024 Prelim P2 Q10]


Solution		Comments
[2]	$\sum x = 1460, \sum y = 108.4 + p, n = 10$ Since $(\bar{x}, \bar{y})$ lies on the regression line of y on x, $\bar{y} = -2.5652 + 0.10168\bar{x}$ $\frac{108.4+p}{10} = -2.5652 + 0.10168\frac{1460}{10}$ $= 12.28008$ $p = 14.4008$ ≈ 14.4 (shown)	
(a) **[2]**	 From GC, product moment correlation coefficient, $r = 0.92147 \approx 0.921$	
(b) **[3]**	From GC, the regression line of y on $\ln x$ is $y = -63.028 + 15.154 \ln x$ $a = -63.028 \approx -63.0$ $b = 15.154 \approx 15.2$ $r = 0.94660 \approx 0.947$	
(c) **[2]**	1) From the scatter diagram, it is observed that as x increases, y increases by decreasing amounts, and 2) product moment correlation coefficient between y and $\ln x$ is 0.947, which is closer to 1 than that of x and y, which is 0.921 . Hence $y = a + b \ln x$ is the better model.	
(d) **[2]**	$y = -63.028 + 15.154 \ln 200$ $= 17.263$ ≈ 17.3 Since $x = 200$ lies outside the given range of x values, $110 \le x \le 188$, the estimated yield may not be reliable.	

(e) **[1]**	Replace x by $25.4x$, $y = -63.028 + 15.154\ln(25.4x)$ $= -63.028 + 15.154[\ln 25.4 + \ln x]$ $= -14.009 + 15.154\ln x$ $= -14.0 + 15.2\ln x$ [Note: the official solution also has a "Comments" column (marking tips); only the Solutions column is reproduced here.]	

7.3 **(a)** A certain brand of breakfast cereal is sold in a variety of different sized packs. Details of the mass of cereal in each pack, m grams, and the price, p cents, are given in the table below.

m	24	250	550	1000	2000	5000
p	100	300	400	600	900	1200

A scatter diagram for the data is shown below.



(i) Explain what the scatter diagram tells you about the relationship between m and p. [1]

The following two models are proposed, where a, b, d and e are constants.

$$p = a + b \ln m \qquad\qquad p = d + e\sqrt{m}$$


(ii) Determine which of these models gives the better fit to the data. State the values of the constants and the product moment correlation coefficient in this case. [4]

(iii) A new pack containing 750 grams of cereal is introduced. Use the model you identified in part **(a)(ii)** to estimate the price of this pack, correct to the nearest 10 cents. Explain whether your answer is reliable. [2]

(b) Kai is investigating the relationship between the number, n, of a particular type of high-performance batteries in a pack and the price, \$$y$, of the pack she found in different stores. Her results are shown in the table below.

n	2	5	6	7	11
y	6	10	16	20	25

This information is shown in the scatter diagram below.



Kai decides to investigate whether the model $y=\frac{5}{2}n+1$ is a good fit for this data.

(i) Draw the line $y=\frac{5}{2}n+1$ on the scatter diagram above. [1]

(ii) For the model $y=\mathrm{f}(n)$, the residual for a point (a, b) is $b-\mathrm{f}(a)$.

(A) Mark the residuals for the points on the scatter diagram above. [1]

(B) Explain why Kai should use the sum of the squares of the residuals rather than the sum of the residuals when assessing the fit of the model. [1]

(C) Calculate the sum of the squares of the residuals for the line $y=\frac{5}{2}n+1$. [1]

Kai's friend points out that other lines parallel to Kai's line can be drawn which are a better fit for the data.

(iii) Explain how the sum of the squares of the residuals for a line that is a better fit for the data differs from the sum of the squares of the residuals for the line $y=\frac{5}{2}n+1$ found in part **(ii)(C)**. [1]

(iv) Find the range of values of c for which the line $y=\frac{5}{2}n+c$ is a better fit for the data than the line $y=\frac{5}{2}n+1$. [3]

[9758 A-Level 2024 P2 Q11]

Solution		Comments
(a)(i)	The scatter diagram tells us that as mass of a pack of cereal (m) increases, the price (p) increases at a decreasing rate.	

(a)(ii)

For $p = a + b\ln m$, product moment correlation coefficient $r_1 = 0.928513$

For $p = d + e\sqrt{m}$, product moment correlation coefficient $r_2 = 0.991438$

Since r_2 is closer to 1 than r_1, the model $\underline{p = d + e\sqrt{m}}$ gives a better fit to the data.

For this model, $d = 32.952567 = \underline{33.0}$ and $e = 17.269804 = \underline{17.3}$

$r = r_2 = \underline{0.991}$ (all answers to 3 s.f.)

(a)(iii)

$p = 32.952567 + 17.269804\sqrt{m}$

Sub. $m = 750$:


$p = 32.952567 + 17.269804\sqrt{750}$

$= 505.906$

$\therefore$ Estimated price $= \underline{510 \text{ cents}}$ (nearest 10 cents)

This estimate is reliable as the product moment correlation coefficient of 0.991 is very close to 1, indicating a strong positive linear correlation between $\sqrt{m}$ and p. Furthermore, the value $m = 750$ is within the data range $24 \le m \le 5000$, hence the estimate is an interpolation.

(b)(i)



(b)(ii)(A)

Point (a, b)	(2, 6)	(5, 10)	(6, 16)	(7, 20)	(11, 25)
f(a)	6	13.5	16	18.5	28.5
Residual $b - f(a)$	0	−3.5	0	1.5	−3.5

(b)(ii)(B)

As residuals may be positive or negative (or zero), they may cancel out if Kai simply sums the residuals. Squaring the residuals allows Kai to sum up positive numbers for a more accurate assessment of the fit of the model.

(b)(ii)(C)

Sum of squares of residuals for the line $y = \frac{5}{2}n + 1$ is $0^2 + (-3.5)^2 + 0^2 + 1.5^2 + (-3.5)^2 = \underline{26.75}$

(b)(iii) The sum of squares of residuals for a line that is a better fit will be less than the sum of squares of residuals for the line $y=\frac{5}{2}n+1$, i.e. less than 26.75.

(b)(iv) Consider $\mathrm{f}(n)=\frac{5}{2}n+c$.

Point (a, b)	(2, 6)	(5, 10)	(6, 16)	(7, 20)	(11, 25)
$\mathrm{f}(a)$	$5+c$	$12.5+c$	$15+c$	$17.5+c$	$27.5+c$
Residual $b-\mathrm{f}(a)$	$1-c$	$-2.5-c$	$1-c$	$2.5-c$	$-2.5-c$

Sum of squares of residuals for the line $y=\frac{5}{2}n+c$ is

$(1-c)^2+(-2.5-c)^2+(1-c)^2+(2.5-c)^2+(-2.5-c)^2$


$=2(1-c)^2+2(2.5+c)^2+(2.5-c)^2$

$=20.75+c+5c^2$

For this to be a better fit,

$20.75+c+5c^2<26.75$

$5c^2+c-6<0$



Using GC, range of values of c is $\underline{-1.2<c<1}$

[Note: SEAB publishes no worked solutions. This is the worked solution from the local file "2024 GCE A-Level H2 Mathematics 9758 Paper 2 Solutions" (tmp_ch10/AL2024_P2_with_sol.pdf); its author is not named on the document.]


7.4 Abi and Bhani find the fuel consumption for a car driven at different constant speeds. The table shows the fuel consumption, y kilometres per litre, for different constant speeds, x kilometres per hour.

x	40	45	50	55	60
y	22	20	18	17	16

(i) Abi decides to model the data using the line $y = 35 - \frac{1}{3}x$.

(a) On the grid

- draw a scatter diagram of the data,
- draw the line $y = 35 - \frac{1}{3}x$. [2]



(b) For a line of best fit $y = \mathrm{f}(x)$, the residual for a point (a, b) plotted on the scatter diagram is the vertical distance between $(a, \mathrm{f}(a))$ and (a, b). Mark the residual for each point on your diagram. [1]

(c) Calculate the sum of the squares of the residuals for Abi's line. [1]

(d) Explain why, in general, the sum of the squares of the residuals rather than the sum of the residuals is used. [1]

Bhani models the same data using a straight line passing through the points (40, 22) and (55, 17).

The sum of the squares of the residuals for Bhani's line is 1.

(ii) State, with a reason, which of the two models, Abi's or Bhani's, gives a better fit. [1]


(iii) State the coordinates of the point that the least squares regression line must pass through. [1]

(iv) Use your calculator to find the equation of the least squares regression line of y on x. State the value of the product moment correlation coefficient. [3]

(v) Use the equation of the regression line to estimate the fuel consumption when the speed is 30 kilometres per hour. Explain whether you would expect this value to be reliable. [2]

(vi) Cerie performs a similar experiment on a different car. She finds that the sum of the squares of the residuals for her line is 0. What can you deduce about the data points in Cerie's experiment? [1]

[9758 A-Level 2019 P2 Q10]

Solution		Comments
(i)(a)		
(i)(b)	The residuals are the distances labelled $e_1, \ldots, e_5$ in the diagram.	
(i)(c)	$\sum_{i=1}^{5}(e_i)^2 = \frac{1}{9}+0+\frac{1}{9}+\frac{1}{9}+1=\frac{4}{3}$ Key the eqn in Y_1 Define $L_3 = Y_{estimated}$ Define $L_4 = e_i^2$ Optional : Using List to calculate the Sum Go to List (Type 2nd → stat → MATH) Sum all the entries in L_4	
(i)(d)	Residuals can be positive or negative since they are calculated by taking (*the observed y-value*) – (*the estimated y-value*). Taking the sum of these residuals will result in them cancelling out and thus losing important information. Taking the sum of the squares of the residuals will avoid this problem.	

(ii) **[1]**	Sum of the squares of the residuals for Bhani's model $= 1 < \frac{4}{3}$ (Sum of the squares of the residuals for Abi's model), Bhani's model is a better fit.	
(iii) **[1]**	$(\bar{x}, \bar{y}) = (50,\ 18.6)$	
(iv) **[3]**	Regression line: $y = -0.3x + 33.6$ $r = -0.985$ (3 s.f.)	
(v) **[2]**	$x = 30$, $y = 24.6$ (kilometres per litre) $x = 30$ is outside the given range of x-values. The estimate obtained using extrapolation is unreliable.	
(vi) **[1]**	Sum of the squares of the residuals = 0 implies that ALL data points lie on the regression line, which is highly unlikely in a real-life experiment.	
	[Note: SEAB publishes no worked solutions. This is the Raffles Institution "Solution for 2019 A-Level Paper 2" (grail.moe item 19456). Its small GC screenshots for (i)(c) are illegible in the source and are described in words only. Question text transcribed from a tutor compilation (Paradigm "Correlation and Regression", Q7 N2019/P2/Q10), as no local copy of the SEAB question paper was found.]	