checking_form_in_manuscript


December 16, 2019

```
[1]: import numpy as np
  from sympy import integrate, exp, symbols, frac, Rational, erf
  from sympy.plotting import plot
  import matplotlib.pyplot as plt
  import lmfit

from sympy import init_printing
  init_printing()
```

The paper writes

[2]:
$$A + (1 - A) \left(-\frac{\sqrt{\pi} \operatorname{erf}\left(\frac{t}{\sigma}\right)}{2} + \frac{1}{2} \right)$$



[3]: <sympy.plotting.plot.Plot at 0x1a22342048>

The function for A=0.6 goes from 1.154 to 0.446

```
[5]: t_vals = np.linspace(-1, 1, 1000)
y = np.array([float(func_subs.subs(dict(t=t_val))) for t_val in t_vals])
```


```
[6]: dt = t_vals[1] - t_vals[0]
    dy_dt = np.diff(y)/dt

model = lmfit.models.GaussianModel()

result = model.fit(-dy_dt, x=t_vals[:-1])

fig, ax = plt.subplots()
    result.plot_fit(ax=ax)
    ax.set_xlabel('t')
    ax.set_ylabel('-dy/dt')
    print(result.fit_report())
```

```
[[Model]]
   Model(gaussian)
[[Fit Statistics]]
   # fitting method
                       = leastsq
   # function evals
                       = 31
   # data points
                       = 999
   # variables
                       = 3
   chi-square
                       = 7.4784e-17
   reduced chi-square = 7.5085e-20
   Akaike info crit
                       = -43988.6402
   Bayesian info crit = -43973.9199
[[Variables]]
                0.07071304 +/- 8.6584e-13 (0.00\%) (init = 1)
   sigma:
               -0.00100100 +/- 8.6584e-13 (0.00\%) (init = 0)
   center:
   amplitude: 0.70898154 +/- 7.5181e-12 (0.00\%) (init = 1)
                0.16651648 +/- 2.0389e-12 (0.00\%) == '2.3548200*sigma'
   fwhm:
   height:
                3.99986660 +/- 4.2415e-11 (0.00\%) ==
'0.3989423*amplitude/max(1.e-15, sigma)'
[[Correlations]] (unreported correlations are < 0.100)
   C(sigma, amplitude) = 0.577
```



when $\sigma = 1$ in the equaiton the fit gives $\sigma = 1/\sqrt(2)$