

# 1 Exercise 1

The implemented gradient descent procedure was tested on three functions:

- (a) A convex function:  $f(x) = -\text{Gaussian}(6, 4)$  (an upside-down Gaussian with mean 6 and variance 4), who's minimum value is at  $(6, -1)$ .
- (b) A scalar convex function of two variables:  $f(x, y) = (x + 2)^2 + y^2$ , who's minimum value is at  $(-2, 0)$ .
- (c) A non-convex function:  $f(x) = \sin(x + \frac{\pi}{2})$ , who has multiple minima at  $(n\pi, -1)$  for all values of  $n$ .

Figure 1 shows the result of calculating the gradient descent on functions (a) and (b) written above. The dots are the calculated minima from gradient descent calculated at multiple step sizes, initial guesses, and thresholds. For convex functions with one minimum, the threshold is the biggest indicator of how many steps the procedure takes to converge. The threshold also determines how close the result is to the real minimum - the smaller the threshold, the closer the result is to the real minimum. Because there is one global minimum, the initial guess does not affect the end result of the function or significantly change the number of iterations required to converge.




Figure 1: (a) Convex function with known minimum value and it's analytical derivative. The dots are the calculated minima according to gradient descent. (b) Contour plot of the gradient of a scalar function with vector-valued arguments. The dots are the calculated minima according to gradient descent.

Figure 2 shows the result of using gradient descent on function (c) with different parameters. The sequence of dots in each case represents the guesses that gradient descent takes to be the minimum. In general, the choice of starting point affects which local minimum gradient descent will find. If the starting point chosen is a local maximum, the procedure will not step towards a minimum because the gradient is zero at the maximum. If the step size is large, the procedure will overshoot the minimum and zig-zag back to the desired minimum, taking a large number of iterations to converge.

Approximating a function's gradient using central differences is very accurate if using an  $h$  of 1. Figure 3 shows graphs of either the gradient function or gradient vector field for the three functions tested above. It is hard to tell the difference between the analytical and numerical functions, so using central differences to numerically estimate the gradient is effective.




Figure 2: The solid line is the function, the dashed line is its analytical derivative, and the series of dots is the series of minimum guesses from gradient descent. (a) Initial guess: 5, step: 0.5, threshold: 0.1 converges in 9 iterations. (b) Initial guess: -2, step: 0.2, threshold: 0.1 converges in 20 iterations. (c) Initial guess: 4, step: 2, threshold: 0.1 converges in 59992 iterations. (d) Initial guess: 0, step: 0.5, threshold: 0.1 converges in 1 iteration, but does not converge to a minimum.




Figure 3: The analytical and numerical gradients for the functions (a) upside-down Gaussian, (b)  $f(x, y) = (x + 2)^2 + y^2$ , and (c)  $f(x) = \sin(x + \frac{\pi}{2})$ . The solid red line is the analytical gradient and the dashed blue line is the numerical gradient.

The BFGS optimizer in SciPy uses a more sophisticated method for estimating minima, so in general, the number of function evaluations needed before convergence is smaller. Table 1 shows a comparison of the gradient descent and BFGS optimizer, where in all cases the step size was 0.2 and the threshold was 0.01.

Function	Initial Guess	Algorithm	Evaluations	Min at
$f(x) = -\text{Gaussian}(6, 4)$	$x = 5$	GD	34	$x = 5.81$
		BFGS	6	$x = 6.00$
$f(x, y) = (x + 2)^2 + y^2$	$(x, y) = (5, 5)$	GD	13	$(x, y) = (-2, 0)$
		BFGS	4	$(x, y) = (-2, 0)$
$f(x) = \sin(x + \frac{\pi}{2})$	$x = 5$	GD	20	$x = 3.15$
		BFGS	6	$x = 3.14$

Table 1: Steps to convergence given a similar set of parameters. GD is gradient descent, BFGS is from SciPy. For gradient descent, all step sizes were 0.5 and thresholds were 0.01.

## 2 Exercise 2




Figure 4: Plots replicating Figure 1.4 in Bishop for different values of  $M$  using different methods. The blue is generated with the analytical solution of the SSE, the green with the gradient descent method on the SSE, and the red using Python’s BFGS optimizer.

	$M = 0$	$M = 1$	$M = 6$	$M = 9$		$M = 0$	$M = 1$	$M = 6$	$M = 9$		$M = 0$	$M = 1$	$M = 6$	$M = 9$
$\hat{w}_0$	0.186	0.820	0.353	0.349	$\hat{w}_0$	0.003	0.499	0.623	-0.014	$\hat{w}_0$	0.186	0.181	0.133	-0.077
$\hat{w}_1$		-1.268	2.619	232.411	$\hat{w}_1$		-0.502	5.577	6.018	$\hat{w}_1$		-0.791	4.974	6.080
$\hat{w}_2$			32.096	-5322.833	$\hat{w}_2$			-14.465	-12.820	$\hat{w}_2$			-0.150	-12.746
$\hat{w}_3$			-206.273	48577.404	$\hat{w}_3$			5.502	6.294	$\hat{w}_3$			4.964	6.398
$\hat{w}_4$			399.001	-231682.150	$\hat{w}_4$			0.475	-5.629	$\hat{w}_4$			-0.027	-5.494
$\hat{w}_5$			-332.709	640159.305	$\hat{w}_5$			0.453	4.425	$\hat{w}_5$			-0.020	4.586
$\hat{w}_6$			105.163	-1061992.54	$\hat{w}_6$			0.435	0.464	$\hat{w}_6$			-0.013	0.646
$\hat{w}_7$				1042586.68	$\hat{w}_7$				0.493	$\hat{w}_7$				0.692
$\hat{w}_8$				-557781.754	$\hat{w}_8$				0.516	$\hat{w}_8$				0.729
$\hat{w}_9$				125223.393	$\hat{w}_9$				0.533	$\hat{w}_9$				0.758

Figure 5: Table 1.1 from Bishop replicated using (a) analytical solutions, (b) gradient descent, and (c) the Scipy BFGS optimizer

	$M = 0$	$M = 1$	$M = 3$
SSE	3.774	2.237	0.350
$\nabla$ SSE analytic	$2.2 \times 10^{-16}$	$(1.3 \times 10^{-15}, -1.3 \times 10^{-15})$	$(-3.3 \times 10^{-14}, -6.4 \times 10^{-15}, -5.8 \times 10^{-15}, -1.7 \times 10^{-14})$
$\nabla$ SSE numeric	0	$(2.7 \times 10^{-15}, -1.3 \times 10^{-15})$	$(-3.4 \times 10^{-14}, -6.2 \times 10^{-15}, -7.7 \times 10^{-15}, -1.7 \times 10^{-14})$

Table 2: Values of the SSE and the analytical and numerical gradients for the given values of  $M$  calculated using the  $w$  coefficients calculated from the analytical solution.

There are three ways to calculate the maximum likelihood weight vector to minimize the  $L2$  norm - (a) analytically with matrices, (b) using gradient descent, and (c), using Scipy’s built-in optimizer. Figure 4 shows the plots from Bishop Figure 1.4 reconstructed using the analytical (blue), gradient descent (green), and Scipy BFGS (red) methods for different values of  $M$ . The coefficients from all three methods for the different values of  $M$  in Bishop’s Table 1.1

are given in Table 5. In general, the analytic computation matched Figure 1.4 in Bishop very well.

The analytic function to compute the gradient of the SSE was compared against the numeric central difference method in Table 2, using the weights computed by the analytic method for the various values of  $M$ . The methods match, so the analytic function is a valid one to use for further optimization.

When recreating Bishop's Figure 1.4 using gradient descent and the BFGS, the same initial guess was given, and the analytic formula of the SSE gradient was used, for simplicity and consistency. Choosing initial guesses, step sizes, and thresholds for gradient descent strongly influences the results. In general, the closer the initial guess is to the analytic solution, the more likely the gradient descent function will find it. This is because the gradient of the SSE has multiple zeros - if there is an optimization for 9 weights, then the SSE function is degree 9, which means it can at most have 4 local minima, according to Euler's theorem. This means that as in the discussion of Exercise 1 above, the gradient descent function will find different minima according to the start value. The threshold and step size were chosen to be 0.001 and 0.01 respectively, so that the gradient descent function came as close to the analytic solutions without taking too long to compute. The BFGS optimizer behaved like the gradient descent minimizer, except no threshold or step size needed to be given. However, it was strongly influenced by the initial guess, as discussed above.

### 3 Exercise 3




Figure 6: Predicted regressions using ridge regression for various values of  $M$  and  $\lambda$

Figure 6 shows the plots of predicted regression values using ridge regression for various values of  $M$  and  $\lambda$ . Because a value of  $\lambda = 0$  corresponds to no regularizer, the plot for  $M = 9$  looks like the plots from Exercise 2 above. As  $\lambda$  increases to 1 and greater for all values of  $M$ , the predicted regression becomes more flat, since the regularizer punishes having large coefficients. Small values of  $\lambda$  that are not 0 yield good results. There is no benefit to increasing  $M$  after a certain point when regressing with a regularizer.




Figure 7: Something

## 4 Exercise 4

This is a test.