

A meta-analysis of the impacts of best management practices on nonpoint source pollutants.

Michael Schramm^{1*}, Duncan Kikoyo¹, Janelle Wright¹, Shubham Jain¹

¹ Texas Water Resources Institute, Texas A&M AgriLife Research, College Station, Texas, USA

Correspondence*:

Michael Schramm

michael.schramm@ag.tamu.edu

2 ABSTRACT

3 Abstract length and content varies depending on article type. Refer to <http://www.frontiersin.org/about/AuthorGuidelines> for abstract requirement and length
4 according to article type.
5

6 **Keywords:** best management practice, water quality, nonpoint source pollution, fecal indicator bacteria, nutrients

1 INTRODUCTION

7 In the United States (U.S.), major progress towards water quality goals have been achieved through the
8 Clean Water Act of 1972. This progress has been largely attributed to investments and reductions in point
9 source discharges while reductions in nonpoint source pollutants remain a substantial challenge (Benham
10 et al., 2008; National Research Council, 2001; Schramm et al., 2022). Increased pollutant loads and
11 pollutant concentrations in runoff are a particular challenge when dealing with nonpoint source pollution
12 resulting from land use change. The impacts of land use change on hydrology and water quality are
13 well established (Allan, 2004; Carpenter et al., 1998; Bernhardt et al., 2008; Carey et al., 2013; Freeman
14 et al., 2019). Nonpoint source driven fecal indicator bacteria (FIB), nitrogen, phosphorus, and suspended
15 sediment remain major causes of water quality impairments in U.S. rivers and streams despite decades of
16 work. In 2017, the Environmental Protection Agency (EPA) estimated 41% or more of the nation's rivers
17 and streams rated poorly for biological condition due to excess nitrogen or phosphorus (EPA, 2017). FIB
18 remains the leading cause of water body impairment on the Clean Water Act 303(d) list in the United States
19 (EPA, 2017).

20 Best management practices (BMPs) have been the primary suite of tools for addressing nonpoint source
21 pollution. BMPs are structural or non-structural controls used to mitigate the effects of increased runoff
22 volume, pollutant loads, or pollutant concentrations emanating from diffuse nonpoint sources. BMPs
23 control the delivery of pollutants through a few possible mechanisms. Structural BMPs reduce and retard
24 total volume of runoff, thus reducing both the volume of water and pollutant load. Structural BMPs may
25 also provide a mechanism for physical, chemical, or biological removal of pollutant constituents suspended
26 or dissolved in runoff. Non-structural BMPs (such as nutrient management or livestock management) are
27 utilized to reduce the generation of pollutant runoff by avoiding pollutant generation during critical periods.

Practitioners rely extensively on mechanistic models to plan and evaluate BMP scenarios and resulting water quality. Lintern et al. (2020) found 43% of reviewed BMP effectiveness studies relied completely on modeled outputs, with modeled outputs almost always predicting water quality improvements following BMP implementation. However, field studies are much more likely to demonstrate mixed results including net releases of pollutants under certain conditions (Lintern et al., 2020; Liu et al., 2017). The disconnect between modeled outcomes and field studies might be attributed to (1) overly simplified or incorrect estimates of parameters that represent management practices (Ullrich and Volk, 2009; Fu et al., 2019; Lintern et al., 2020), (2) failure to incorporate uncertainty in estimates (Tasdighi et al., 2018; Fu et al., 2019; Lintern et al., 2020), and (3) assumption of static performance over time (Meals et al., 2010; Liu et al., 2017; Fu et al., 2019).

A source of uncertainty comes from the substantial variability of performance metrics reported in empirical BMP studies (Lintern et al., 2020). Previous empirical reviews generally describe high variability and uncertainty in nitrogen and phosphorus removal and consistent reduction in total suspended sediment concentrations across BMP types (Lintern et al., 2020; Liu et al., 2017; Koch et al., 2014; Clary et al., 2011; Barrett, 2008; Grudzinski et al., 2020). The review literature on the effects of BMPs on FIBs are sparse but generally find extremely high variance in performance across BMPs (Clary et al., 2011; Grudzinski et al., 2020). Accurately characterizing BMP treated runoff quality is challenging due to the high variability in BMP performance. However, influent water quality, seasonality, and BMP age provide some explanation to highly heterogeneous BMP efficiency data (Barrett, 2005; Liu et al., 2017).

A recent meta-analysis demonstrated the further linkages between local climatic conditions and BMP performance on nitrogen and phosphorus concentrations (Horvath et al., 2023).

2 METHODS

We conducted a systematic review of recent (2000-2022) literature to compile field studies documenting the effectiveness of best management practices on fecal indicator bacteria and nutrients concentrations. The systematic review followed guidance provided in the Collaboration for Environmental Evidence systematic review guidelines (Collaboration for Environmental Evidence, 2018). In order to maximize the number of studies included in the review, we included both peer-reviewed studies and unpublished white papers to reduce potential bias against negative results. The inclusion criteria filtered out (1) non-field studies, (2) modelling results, (3) studies that did not evaluate specific BMPs, (4) studies conducted outside of the U.S. or published in a language other than English. We ran search queries in Texas A&M Library Catalog, Web of Science, and Google Scholar. Although results from Google Scholar are not always replicable, we utilized the service to maximize search results for studies not published in academic journals and presumably increase the chance of identifying studies with negative effects. Fecal indicator bacteria study searches included the following query: “fecal indicator bacteria” OR “E. coli” OR “Escherichia coli” OR “enterococci” OR “enterococcus” AND “best management practices” OR “BMPs” AND “effectiveness” OR “performance”. Nutrient BMP studies utilized a similar query: “nutrient” OR “nitrogen” OR “phosphorus” OR “sediment” OR “TSS” AND “best management practices” OR “BMPs” AND “effectiveness” OR “performance”.

Results from each database were first filtered to remove duplicates. After removal of duplicates, each member of the research team ($n = 4$) was assigned a subset of studies to evaluate if they should be included. **include supplementary tables with inclusion criteria and data variables** Each study was reviewed by two team members and differences in opinion were collectively discussed and agreed upon before progressing. The remaining studies were split among team members for data extraction, again with at least

Table 1. Criteria applied for including or excluding studies within the review database.

| Attribute | Inclusion Criteria | Exclusion Criteria |
|----------------------|---|---|
| Study type | Journal articles, book chapters, conference papers, unpublished research reports, thesis and dissertations, organizational and agency white papers. | Synopsis or review studies, reports with reductions based on modeled or other estimated reductions (e.g. TMDLs, watershed plans, or modelling studies). |
| Outcomes | Field studies with measured effects on fecal indicator bacteria or nutrient concentrations. | Studies not explicitly linking reductions to a specific BMP or insufficient information to quantify reductions. |
| Geographical context | Studies conducted within the United States. | Studies outside of the United States. |
| Timeframe | Studies published from 2000 through 2022. | Studies published prior to 2000 or after 2022. |

70 two team members reviewing each study. If data was provided in figures, the data was extracted with the
 71 WebPlotDigitizer tool (Rohatgi, 2022).

72 Effect size calculation

73 Prior BMP reviews routinely report the effect of BMPs using efficiency values (see Agouridis et al., 2005;
 74 Kroger et al., 2012; Liu et al., 2015; Simpson and Weammert, 2009). Efficiency values are calculated as:

$$\text{Efficiency} = \frac{x_{control} - x_{experiment}}{x_{control}} \times 100,$$

75 where $x_{control}$ is the pre-treatment or control pollutant concentration and $x_{experiment}$ is the pollutant
 76 concentration measured after the BMP intervention. There are several statistical shortcomings (distributional
 77 asymmetry, skewness, and non-additive properties) when using efficiency to estimate overall effect sizes
 78 (Nuzzo, 2018; Cole and Altman, 2017). In comparison, the log ratio of means (ROM) provide preferable
 79 statistical properties for regression analysis (Osenberg et al., 1997; Hedges et al., 1999). ROM quantifies
 80 the difference in means between the control and experimental group (Hedges et al., 1999):

$$ROM_i = \ln\left(\frac{x_{i,control}}{x_{i,experiment}}\right) = \ln(x_{i,control}) - \ln(x_{i,experiment}),$$

81 where $x_{i,control}$ and $x_{i,experiment}$ are the mean pollutant concentrations for experiment i . The statistical
 82 properties of $\ln RR$ (normal distribution around zero and additive properties) are preferable to the more
 83 commonly reported efficiency metric (Osenberg et al., 1997; Hedges et al., 1999). Use of ROM required
 84 the exclusion of studies that only provided measures of BMP efficiency and not the underlying data used to
 85 derive the metric.

86 2.1 Statistical models

87 We used the “rma.mv” function in the *metafor* R package to fit multilevel random effects linear models
 88 with ROM as the effect variable (Viechtbauer, 2010; R Core Team, 2023). Our models specified a nested
 89 random effects term accounting for heterogeneity between effect sizes from the same study and for

Table 2. Study and effect variables extracted for review.

| Variable | Description |
|--------------------|---|
| Publication Year | Year the study was published |
| Parameter | The specific pollutant measured. |
| Runoff source | Dominant source of runoff (rop fields, livestock pasture, commerical, residential) |
| Source type | Major categorization of runoff source: agricultural or urban |
| BMP | BMP evaluated |
| BMP Classification | BMP description based on NRCS conservation practice standards and EPA BMP fact sheets |
| BMP Category | BMP categorization based on structural or management |
| BMP Subcategory | BMP subcategorization based on pollutant removal processes |
| Study scale | Spatial scale of the study area ("lot/field", "community", "watershed") |
| Location | Location name used in the study description |
| State | State where the study was conducted |
| Study area | Drainage area in hectares |
| Longitude | Approximated or reported latitude coordinate |
| Latitude | Approximated or reported longitude coordinate |
| Study years | Year or years when data were collected |
| N control | Number of control measurements |
| N experiment | Number of experimental measurements |
| X control | Mean concentration for control measurements |
| X experiment | Mean concentration for experimental measurements |
| SE control | Standard error of control measurements |
| SE experiment | Standard error of experimental measurements |
| Minimum control | Minimum control measurement |
| Minimum experiment | Minimum experiment measurement |
| Maximum control | Maximum control measurement |
| Maximum experiment | Maximum experiment measurement |
| SD control | Standard deviation of control measrements |
| SD experiment | Standard deviation of experimental measurements |
| Units | Units reported by the study |
| Percent reduction | BMP efficiency for studies that only reported efficiency |

90 heterogeneity between studies. A key feature of meta-analysis is the weighting of effects using sampling
 91 variance of individual effect sizes. 45% of fecal indicator bacteria, 68% of TN, 68% of TP, and 70% of TSS
 92 effect sizes were missing standard deviations required to estimate sampling variance. Removal of studies
 93 due to missing variance information can introduce substantial bias (Kambach et al., 2020), so we imputed
 94 missing standard deviations using random forest based multivariate imputation with chained equations
 95 using the *mice* R package (Buuren and Groothuis-Oudshoorn, 2011). Sampling variance was estimated
 96 with the variance estimator described in Doncaster et al. to reduce bias in small sample studies:

$$v(ROM) = \frac{\sum_{i=1}^K (CV_{control,i}^2)/K}{n_{control}} + \frac{\sum_{i=1}^K (CV_{experiment,i}^2)/K}{n_{experiment}},$$

Table 3. Summary table of moderator effects for performance of BMPs on fecal indicator bacteria concentrations.

| Moderator | Estimate ($\ln RR$) | 95% CI | SE | T-statistic | df | p-value |
|--------------------------|--------------------------|---------------|-------|-------------|----|---------|
| Intercept | 0.21 | [-2.07,2.48] | 1.06 | 0.19 | 14 | 0.85 |
| BMP Subcategories | | | | | | |
| Filtration | -0.39 | [-6.42,5.63] | 2.81 | -0.14 | 14 | 0.89 |
| Infiltration | -1.24 | [-3.50,1.01] | 1.13 | -1.10 | 64 | 0.28 |
| Livestock | -1.18 | [-3.25,0.89] | 0.96 | -1.23 | 14 | 0.24 |
| Treatment | -1.56 | [-2.63,-0.49] | 0.53 | -2.92 | 64 | <0.01 |
| log(Concentration) | 0.20 | [0.084,0.32] | 0.059 | 3.42 | 65 | <0.01 |

$$I^2_{\text{total}}=21.48, I^2_{\text{study}}=0, I^2_{\text{effect}}=21.48; R^2_{\text{marginal}}=0.46$$

97 where v represents the sampling variance, $CV_{control,i}$ and $CV_{experiment,i}^2$ are the coefficients of variation
 98 from the i th study for studies 1, 2, ..., K .

99 Our initial models included log transformed pre-treatment pollutant concentration, study scale (plot/field,
 100 community, watershed), study length (n years), BMP subcategory (), and pollutant concentration:BMP
 101 subcategory interactions were included as fixed effect terms. We used an information-theoretic approach
 102 to select the most parsimonious model from the subset of candidate models based on corrected Akaike
 103 information criterion (AIC_c) estimated with maximum likelihood (Cinar et al., 2021). The final model
 104 was selected from candidate models, which included all combination and subsets of the full model, by
 105 eliminating candidate models until the ΔAIC_c reached ≤ 2 (Burnham et al., 2011). We estimated regression
 106 coefficients of the selected model using restricted maximum likelihood. Relative heterogeneity between
 107 and within studies were calculated using the I^2 metric described in Nakagawa and Santos (2012). Marginal
 108 R^2 was used to describe the amount of variance explained by fixed effects (Nakagawa and Schielzeth,
 109 2013).

110 We tested for evidence of publication bias, in the form of small study effect, by using the extension of
 111 Egger's regression applied to the multilevel model framework that included adjusted sampling error as
 112 a moderator (Nakagawa et al., 2023). We did not identify evidence of publication bias in the surveyed
 113 studies for fecal indicator bacteria ($Z = -1.17, p = 0.25$), total nitrogen, total phosphorus, or total suspended
 114 solids; therefore, adjustments for publication bias were not included in the final models. We conducted
 115 a sensitivity analysis of the robustness of overall effect sizes to individual studies using leave-one-out
 116 analysis (Nakagawa et al., 2023). Outlier studies identified in the sensitivity analysis are highlighted in
 117 Table X and were removed prior to fitting the final models.

3 RESULTS

118 3.1 Summary of BMP literature

119 3.2 Regression results

120 Some evidence of a relationship between input concentrations and fecal indicator bacteria reductions. No
 121 evidence strong evidence that the type of BMP changed the overall relationship.

Table 4. Summary table of moderator effects for performance of BMPs on nitrogen removal.

| Moderator | Estimate ($\ln RR$) | 95% CI | SE | T-statistic | df | p-value |
|-------------------------|--------------------------|---------------|-------|-------------|-----|---------|
| Intercept | 0.95 | [0.38,1.52] | 0.28 | 3.41 | 28 | <0.01 |
| log(Concentration) | -0.034 | [-0.14,0.071] | 0.053 | -0.64 | 124 | 0.525 |
| Adjusted sampling error | -1.54 | [-3.30,0.22] | 0.89 | -1.74 | 124 | 0.085 |

$I^2_{\text{total}}=96.57$, $I^2_{\text{study}}=52.45$, $I^2_{\text{effect}}=44.12$; $R^2_{\text{marginal}}=0.05$

Table 5. Summary table of moderator effects for performance of BMPs on phosphorus removal.

| Moderator | Estimate ($\ln RR$) | 95% CI | SE | T-statistic | df | p-value |
|--------------------|--------------------------|--------------|-------|-------------|----|---------|
| Intercept | 0.40 | [-0.14,0.93] | 0.26 | 1.52 | 28 | 0.14 |
| log(Concentration) | 0.27 | [0.15,0.40] | 0.063 | 4.30 | 97 | <0.01 |

$I^2_{\text{total}}=15.54$, $I^2_{\text{study}}=0$, $I^2_{\text{effect}}=15.54$; $R^2_{\text{marginal}}=0.32$

122 3.3 Subsection 2

123 Frontiers requires figures to be submitted individually, in the same order as they are referred to in the
 124 manuscript. Figures will then be automatically embedded at the bottom of the submitted manuscript. Kindly
 125 ensure that each table and figure is mentioned in the text and in numerical order. Permission must be
 126 obtained for use of copyrighted material from other sources (including the web). Please note that it is
 127 compulsory to follow figure instructions. Figures which are not according to the guidelines will cause
 128 substantial delay during the production process.

4 DISCUSSION

129 Linking BMP performance to watershed scale improvements remains a challenge (Tomer and Locke,
 130 Melland et al, Meals). The successful implementation and maintenance of BMPs at watershed scales remains
 131 an area for more investigation (Liu et al., Lintern) Long-term trans-disciplinary monitoring projects not
 132 just on BMP performance but maintenance, management, and social aspects may elucidate unknown factors
 133 influencing the ability to scale BMP reductions to watershed scale improvements (Lintern et al.)

134 Runoff volume was not incorporated into our models, largely attributed to lack of available data. However,
 135 BMP size is typically scaled to accommodate runoff volumes of a particular return period volume (1-yr or
 136 2-yr flows for example) (Hoss et al.).

DATA AVAILABILITY STATEMENT

137 The data and R code used in this study are deposited in Zenodo: TBD.

DISCLOSURE/CONFLICT-OF-INTEREST STATEMENT

138 The authors declare that the research was conducted in the absence of any commercial or financial
 139 relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

140 MS: Conceptualization, Formal analysis, Funding acquisition, Methodology, Supervision, Writing—
141 original draft, Writing—review & editing. DK: Conceptualization, Formal analysis, Data collection,
142 Writing—original draft, Writing—review & editing. JW: Data collection, Writing—review & editing. SJ:
143 Data collection; Writing—review & editing.

ACKNOWLEDGMENTS

144 Funding: This project was supported by a state nonpoint source grant from the Texas State Soil and Water
145 Conservation Board.

SUPPLEMENTAL DATA

146 Supplementary Material should be uploaded separately on submission, if there are Supplementary Figures,
147 please include the caption in the same file as the figure. LaTeX Supplementary Material templates can be
148 found in the Frontiers LaTeX folder

REFERENCES

- 149 Agouridis, C. T., Workman, S. R., Warner, R. C., and Jennings, G. D. (2005). Livestock grazing
150 management impacts on stream water quality: A review. *Journal of the American Water Resources
151 Association* 41, 591–606. doi:10.1111/j.1752-1688.2005.tb03757.x
- 152 Allan, J. D. (2004). Landscapes and Riverscapes: The Influence of Land Use on Stream Ecosystems.
153 *Annual Review of Ecology, Evolution, and Systematics* 35, 257–284. doi:10.1146/annurev.ecolsys.35.
154 120202.110122
- 155 Barrett, M. E. (2005). Performance comparison of structural stormwater best management practices. *Water
156 Environment Research* 77, 78–86. doi:10.2175/106143005X41654
- 157 Barrett, M. E. (2008). Comparison of BMP performance using the International BMP Database. *Journal of
158 Irrigation and Drainage Engineering* 134, 556–561. doi:10.1061/(ASCE)0733-9437(2008)134:5(556)
- 159 Benham, B., Zeckoski, R., and Yagow, G. (2008). Lessons learned from TMDL implementation case
160 studies. *Water Practice* 2, 1–13. doi:10.2175/193317708X281370
- 161 Bernhardt, E. S., Band, L. E., Walsh, C. J., and Berke, P. E. (2008). Understanding, managing, and
162 minimizing urban impacts on surface water nitrogen loading. *Annals of the New York Academy of
163 Sciences* 1134, 61–96. doi:10.1196/annals.1439.014
- 164 Burnham, K. P., Anderson, D. R., and Huyvaert, K. P. (2011). AIC model selection and multimodel
165 inference in behavioral ecology: Some background, observations, and comparisons. *Behavioral Ecology
166 and Sociobiology* 65, 23–35. doi:10.1007/s00265-010-1029-6
- 167 Buuren, S. V. and Groothuis-Oudshoorn, K. (2011). Mice: Multivariate Imputation by Chained Equations
168 in R. *Journal of Statistical Software* 45. doi:10.18637/jss.v045.i03
- 169 Carey, R. O., Hochmuth, G. J., Martinez, C. J., Boyer, T. H., Dukes, M. D., Toor, G. S., et al. (2013).
170 Evaluating nutrient impacts in urban watersheds: Challenges and research opportunities. *Environmental
171 Pollution* 173, 138–149. doi:10.1016/j.envpol.2012.10.004
- 172 Carpenter, S. R., Caraco, N. F., Correll, D. L., Howarth, R. W., Sharpley, A. N., and Smith, V. H. (1998).
173 Nonpoint pollution of surface waters with phosphorus and nitrogen. *Ecological Applications* 8, 559–568.
174 doi:10.1890/1051-0761(1998)008[0559:NPOSWW]2.0.CO;2

- 175 Cinar, O., Umbanhowar, J., Hoeksema, J. D., and Viechtbauer, W. (2021). Using information-theoretic
176 approaches for model selection in meta-analysis. *Research Synthesis Methods* 12, 537–556. doi:10.
177 1002/jrsm.1489
- 178 Clary, J., Leisenring, M., Poresky, A., Earles, A., and Jones, J. (2011). BMP performance analysis
179 results for the International Stormwater BMP Database. In *World Environmental and Water Resources*
180 *Congress 2011* (Palm Springs, California, United States: American Society of Civil Engineers), 441–449.
181 doi:10.1061/41173(414)47
- 182 Cole, T. J. and Altman, D. G. (2017). Statistics Notes: What is a percentage difference? *BMJ* , j3663doi:10.
183 1136/bmj.j3663
- 184 Collaboration for Environmental Evidence (2018). *Guidelines and Standards for Evidence Synthesis in*
185 *Environmental Management*. Tech. Rep. Version 5.0
- 186 EPA (2017). *National Water Quality Inventory: Report to Congress*. Tech. Rep. EPA 841-R-16-011, U.S.
187 Environmental Protection Agency, Washington, D.C.
- 188 Freeman, L. A., Corbett, D. R., Fitzgerald, A. M., Lemley, D. A., Quigg, A., and Steppe, C. N. (2019).
189 Impacts of urbanization and development on estuarine ecosystems and water quality. *Estuaries and*
190 *Coasts* 42, 1821–1838. doi:10.1007/s12237-019-00597-z
- 191 Fu, B., Merritt, W. S., Croke, B. F., Weber, T. R., and Jakeman, A. J. (2019). A review of catchment-scale
192 water quality and erosion models and a synthesis of future prospects. *Environmental Modelling &*
193 *Software* 114, 75–97. doi:10.1016/j.envsoft.2018.12.008
- 194 Grudzinski, B., Fritz, K., and Dodds, W. (2020). Does riparian fencing protect stream water quality in
195 cattle-grazed lands? *Environmental Management* 66, 121–135. doi:10.1007/s00267-020-01297-2
- 196 Hedges, L. V., Gurevitch, J., and Curtis, P. S. (1999). The meta-analysis of response ratios in experimental
197 ecology. *Ecology* 80, 1150–1156. doi:10.1890/0012-9658(1999)080[1150:TMAORR]2.0.CO;2
- 198 Horvath, I. R., Zhang, K., Mayer, B. K., and Parolari, A. J. (2023). Effects of regional climate and
199 BMP type on stormwater nutrient concentrations in BMPs: A meta-analysis. *Environmental Science &*
200 *Technology* 57, 5079–5088. doi:10.1021/acs.est.2c05942
- 201 Kambach, S., Bruelheide, H., Gerstner, K., Gurevitch, J., Beckmann, M., and Seppelt, R. (2020).
202 Consequences of multiple imputation of missing standard deviations and sample sizes in meta-analysis.
203 *Ecology and Evolution* 10, 11699–11712. doi:10.1002/ece3.6806
- 204 Koch, B. J., Febria, C. M., Gevrey, M., Wainger, L. A., and Palmer, M. A. (2014). Nitrogen removal by
205 stormwater management structures: A data synthesis. *JAWRA Journal of the American Water Resources*
206 *Association* 50, 1594–1607. doi:10.1111/jawr.12223
- 207 Kroger, R., Perez, M., Walker, S., and Sharpley, A. (2012). Review of best management practice reduction
208 efficiencies in the Lower Mississippi Alluvial Valley. *Journal of Soil and Water Conservation* 67,
209 556–563. doi:10.2489/jswc.67.6.556
- 210 Lintern, A., McPhillips, L., Winfrey, B., Duncan, J., and Grady, C. (2020). Best management practices for
211 diffuse nutrient pollution: Wicked problems across urban and agricultural watersheds. *Environmental*
212 *Science & Technology* 54, 9159–9174. doi:10.1021/acs.est.9b07511
- 213 Liu, Y., Ahiablame, L. M., Bralts, V. F., and Engel, B. A. (2015). Enhancing a rainfall-runoff model to
214 assess the impacts of BMPs and LID practices on storm runoff. *Journal of Environmental Management*
215 147, 12–23. doi:10.1016/j.jenvman.2014.09.005
- 216 Liu, Y., Engel, B. A., Flanagan, D. C., Gitau, M. W., McMillan, S. K., and Chaubey, I. (2017). A review
217 on effectiveness of best management practices in improving hydrology and water quality: Needs and
218 opportunities. *Science of The Total Environment* 601–602, 580–593. doi:10.1016/j.scitotenv.2017.05.212

- 219 Meals, D. W., Dressing, S. A., and Davenport, T. E. (2010). Lag time in water quality response to best
220 management practices: A review. *Journal of Environment Quality* 39, 85. doi:10.2134/jeq2009.0108
- 221 Nakagawa, S. and Santos, E. S. A. (2012). Methodological issues and advances in biological meta-analysis.
222 *Evolutionary Ecology* 26, 1253–1274. doi:10.1007/s10682-012-9555-5
- 223 Nakagawa, S. and Schielzeth, H. (2013). A general and simple method for obtaining R^2 from generalized
224 linear mixed-effects models. *Methods in Ecology and Evolution* 4, 133–142. doi:10.1111/j.2041-210x.
225 2012.00261.x
- 226 Nakagawa, S., Yang, Y., Macartney, E. L., Spake, R., and Lagisz, M. (2023). Quantitative evidence
227 synthesis: A practical guide on meta-analysis, meta-regression, and publication bias tests for
228 environmental sciences. *Environmental Evidence* 12, 8. doi:10.1186/s13750-023-00301-6
- 229 National Research Council (2001). *Assessing the TMDL Approach to Water Quality Management*
230 (Washington, D.C.: National Academies Press). doi:10.17226/10146
- 231 Nuzzo, R. (2018). Percent differences: Another look. *PM&R* 10, 661–664. doi:10.1016/j.pmrj.2018.05.003
- 232 Osenberg, C. W., Sarnelle, O., and Cooper, S. D. (1997). Effect size in ecological experiments: The
233 application of biological models in meta-analysis. *The American Naturalist* 150, 798–812. doi:10.1086/
234 286095
- 235 R Core Team (2023). *R: A Language and Environment for Statistical Computing*. Vienna, Austria
- 236 Rohatgi, A. (2022). *WebPlotDigitizer*. Pacifica, California
- 237 Schramm, M., Gitter, A., and Gregory, L. (2022). Total Maximum Daily Loads and *Escherichia coli*
238 trends in Texas freshwater streams. *Journal of Contemporary Water Research & Education* 176, 36–49.
239 doi:10.1111/j.1936-704X.2022.3374.x
- 240 Simpson, D. T. and Weammert, S. (2009). *Developing Best Management Practice Definitions and
241 Effectiveness Estimates for Nitrogen, Phosphorus and Sediment in the Chesapeake Bay Watershed*. Tech.
242 rep., University of Maryland Mid-Atlantic Water Program
- 243 Tasdighi, A., Arabi, M., Harmel, D., and Line, D. (2018). A Bayesian total uncertainty analysis framework
244 for assessment of management practices using watershed models. *Environmental Modelling & Software*
245 108, 240–252. doi:10.1016/j.envsoft.2018.08.006
- 246 Ullrich, A. and Volk, M. (2009). Application of the Soil and Water Assessment Tool (SWAT) to predict
247 the impact of alternative management practices on water quality and quantity. *Agricultural Water
Management* 96, 1207–1217. doi:10.1016/j.agwat.2009.03.010
- 248 Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *Journal of Statistical
249 Software* 36. doi:10.18637/jss.v036.i03


FIGURE CAPTIONS

Figure 1. Distribution of studies identified in the systematic review by state and parameter.




Figure 2. Summary of (A) study scale, (B) dominant runoff source, and (C) BMPs identified in the systematic review.

