

Assessing linkages between watershed nutrient  
loading and water quality in a subtropical estuary  
with semiparametric models.

Michael Schramm<sup>1\*</sup>

<sup>1\*</sup>Texas Water Resources Institute, Texas A&M AgriLife Research,  
College Station, United States, 77840, TX, 1001 Holleman Dr. E..

Corresponding author(s). E-mail(s): [michael.schramm@ag.tamu.edu](mailto:michael.schramm@ag.tamu.edu);

**Abstract**

Lavaca Bay is a secondary embayment on the Texas coast displaying early signals of water quality degradation. This study applied a semiparametric approach to assess both watershed nutrient loads and responses in estuary water quality to nutrient loading and streamflow. Cross-validation indicated that, despite data constraints, semiparametric models performed well at nutrient load prediction. Based on these models, delivered annual nutrient loads varied substantially from year to year. In contrast, minimal changes in flow-normalized loads indicate that nutrient loadings were driven by natural variation in precipitation and runoff as opposed to changes in management of nonpoint sources. Models indicated there was no evidence of long-term changes in dissolved oxygen or chlorophyll-*a* within Lavaca Bay. However, site specific long-term increases in both organic and inorganic nitrogen are concerning. Further analysis found freshwater inflow was a strong driver of nutrient and chlorophyll-*a* concentrations within Lavaca Bay but there was no evidence that changes in watershed nutrient loading explained variation in dissolved oxygen or chlorophyll-*a* concentrations. In addition to providing a baseline assessment of watershed nutrient loading and water quality responses in the Lavaca Bay watershed, this study provides methodological support for the use of semiparametric methods in load regression models and estuary assessments.

**Keywords:** key, dictionary, word

047 *Remaining Items*

048

049 • keywords

050

051 • discussion

052

053 • formalize language (reduce first person active?)

054

055 • final language review

056

057 • springer doesn't want figures in subdirectory, need to knit to same directory as tex

058

059 file.

060

## 061 **1 Introduction**

062

063 Similar to many coastal areas globally, the coastal watersheds along the Texas Gulf

064

065 coast are facing pressures from increasing population, increases in point source and

066

067 non-point source pollution and alterations to freshwater flows that degrade water qual-

068 ity in downstream estuaries ([Bricker et al, 2008](#); [Kennicutt, 2017](#); [Bugica et al, 2020](#)).

069

070 Despite these escalating pressures, national scale assessments have classified coastal

071

072 estuaries in Texas as moderate or low risk for eutrophic conditions ([Bricker et al,](#)

073

074 [2008](#)). However, a suite of recent studies indicates that estuary water quality dynamics

075

076 in both agriculturally dominated and urban watersheds within Texas are expressing

077

078 conditions that are increasingly conducive to algal blooms and eutrophication ([Wetz](#)

079

080 [et al, 2016, 2017](#); [Bugica et al, 2020](#); [Chin et al, 2022](#)). With identification of several

081

082 localized areas of estuary water quality concern along the Texas coast ([Bugica et al,](#)

083

084 [2020](#)), localized studies are being prioritized to better inform management actions.

085

086 This project aims to provide an assessment of watershed nutrient loading and

087

088 resulting water quality responses in Lavaca Bay, Texas. Lavaca Bay is a secondary bay

089

090 in the larger Matagorda Bay system located roughly halfway between Houston, Texas

091

092 and Corpus Christi, Texas. Lavaca Bay faces substantial challenges associated with

093

094 legacy contamination but general water quality parameters such as dissolved oxygen

095

096

(DO), nutrients, and biological parameters have been well within state water quality standards. Despite largely meeting statewide water quality standards, there have been concerning recently identified declines in abundance, biomass, and diversity of benthic fauna in Lavaca Bay (Beseres Pollack et al, 2011). These declines are partially attributed to reductions in freshwater inflow and changes in estuary salinity and are indicative of an already stressed system (Beseres Pollack et al, 2011; Palmer and Montagna, 2015; Montagna et al, 2020). More recently, significant linear increases in total phosphorus (TP), orthophosphate, total Kjeldahl nitrogen (TKN), and chlorophyll-*a* concentrations were identified at monitoring sites within Lavaca Bay (Bugica et al, 2020). Although long-term changes in DO concentrations have not been identified, the trends in nutrient concentrations are concerning due to the role of nitrogen as a limiting factor for primary production in many Texas estuaries (Gardner et al, 2006; Hou et al, 2012; Dorado et al, 2015; Paudel et al, 2019; Wetz et al, 2017) and the ramifications that changes in nitrogen loadings could have for productivity and eutrophication in Lavaca Bay.

There are ongoing efforts between local, state, and federal agencies to address water quality impairments in the freshwater portions of the Lavaca Bay watershed (Jain and Schramm, 2021; Schramm et al, 2018; Berthold et al, 2021). However, at a statewide scale, these approaches have shown limited success and emphasize a need for improved efforts at assessing and linking management actions with downstream water quality to identify and replicate effective management actions across the state (Schramm et al, 2022). The identification and communication of changes and trends in water quality is complicated by the fact that trends are often non-linear and confounded by precipitation and runoff that hinder traditional analysis (Wazniak et al, 2007; Lloyd et al, 2014). The development and application of flexible statistical methods such as Weighted Regressions on Time, Discharge and Season (WRTDS, Hirsch et al, 2010) and Generalized Additive Models (GAMs, Wood, 2011) have provided effective tools

for researchers to quantify and communicate non-linear changes in river and estuary pollutant loadings.

WRTDS calculates a time series of in-stream concentrations or loads (daily, monthly, or annually) and flow-normalized estimates of concentrations and loads using locally weighted regression for unique combinations of time, discharge, and season. WRTDS has been widely used to assess and identify trends in riverine nutrients (Oelsner and Stets, 2019; Stackpoole et al, 2021), chlorides (Stets et al, 2018), and other pollutants of concern (Shoda et al, 2019). WRTDS has also been successfully adapted to assess trends in estuarine water quality concentrations (Beck et al, 2018).

While WRTDS is a statistical approach developed specifically for water quality applications, GAMs are a broadly applicable statistical method. GAMs are a semiparametric extension of generalized linear models where the linear predictor is represented as the sum of multiple unknown smooth functions and parametric linear predictors (Wood, 2011). Although the underlying parameter estimation procedure of GAMs is substantially different than WRTDS, both the functional form and results have been demonstrated to be similar when assessing nutrient concentration trends (Beck and Murphy, 2017). Water quality applications of GAMs have included include river and catchment nutrient concentration and load estimation (Wang et al, 2011; Kroon et al, 2012; Kuhnert et al, 2012; Robson and Dourdet, 2015; Hagemann et al, 2016; McDowell et al, 2021; Biagi et al, 2022), assessment of temporal trends of nutrients (Beck and Murphy, 2017; Murphy et al, 2019), phytoplankton (Bergbusch et al, 2021), and cyanobacteria (Hayes et al, 2020). Recently GAMs have also been used to link water quality responses in receiving water bodies to changes in nonpoint source nutrient inputs (Murphy et al, 2022). Beck and Murphy (2017) provides a substantial discussion on the differences (and similarities) between GAMs and WRTDS for water quality applications.

To provide actionable information for resource managers in Lavaca Bay, water quality conditions must be evaluated relative to changes in natural environmental drivers to better understand and manage potential human impacts. This study utilizes GAMs to develop estimates of delivered and flow-normalized nutrient loads and assess changes in loads delivered to Lavaca Bay. GAMs were chosen over other regression-based approaches for use in this study due to; (1) the ability to easily explore and incorporate different model terms; (2) the incorporation of non-linear smooth functions that do not require explicit a priori knowledge of the expected shape; and (3) inclusion of a link function that related the expected value of the response to linear predictors thus avoiding unneeded data transformations and bias corrections. The exploratory study also assesses the response of water quality parameters in Lavaca Bay over time and in response to freshwater inflow controlled for seasonality and to watershed nutrient loads that are controlled for environmentally driven variation.

## 2 Methods


### 2.1 Location and Data

Lavaca Bay is 190 km<sup>2</sup> with the majority of freshwater inflow provided by the Lavaca and Navidad River systems (Figure 1). The Garcitas-Arenosa, Placedo Creek, and Cox Bay watersheds provide additional freshwater inflows. The entire watershed land area is 8,149 km<sup>2</sup> and primarily rural. Watershed land cover and land use is 50% grazed pasture and rangeland, 20% cultivated cropland (primarily row crops such as corn, cotton, and sorghum), and 5% suburban/urban. Pasture and rangeland is concentrated in the Lavaca River watershed, while cultivated crops are generally located along the eastern tributaries of the Navidad river. The Lavaca and Navidad River watersheds are a combined 5,966 km<sup>2</sup>, or approximately 73% of the entire Lavaca Bay watershed area. Discharge from the Navidad River is regulated by Lake Texana which has been in operation since 1980. Lake Texana provides 0.210 km<sup>3</sup> of water storage and discharges

231 into the tidal section of the Navidad River which ultimately joins the tidal section of  
232 the Lavaca River 15 km upstream of the confluence with the Lavaca Bay.  
233

234 Daily discharges for the Lavaca River (USGS-08164000, Figure 1) were obtained  
235 from the United States Geologic Survey (USGS) National Water Information System  
236 using the *dataRetrieval* R package (De Cicco et al, 2022). Gaged daily discharges from  
237 the outlet of Lake Texana on the Navidad River (USGS-0816425) were provided by the  
238 Texas Water Development Board (TWDB) (April 21, 2022 email from R. Neupane,  
239 TWDB).  
240

241 Water quality sample data for the two freshwater and three estuary locations were  
242 obtained from the Texas Commission on Environmental Quality (TCEQ) Surface  
243 Water Quality Monitoring Information System. Data submitted through the system  
244 are required to be collected under Quality Assurance Project Plans and lab method  
245 procedures outlined by the TCEQ's procedures manual to ensure consistent collection  
246 and laboratory methods are applied between samples collected by different entities  
247 and under different projects. All sites had varying lengths of and availability of data.  
248 For freshwater locations, TP from January 2000 through December 2020 and nitrate-  
249 nitrogen ( $\text{NO}_3$ ) data from January 2005 through December 2020 were downloaded  
250 (Table 1). Less than 5-years of total nitrogen and TKN concentration data were avail-  
251 able at the freshwater sites and deemed insufficient to develop load estimation models  
252 (Horowitz, 2003; Snelder et al, 2017). The three estuary sites included an upper Lavaca  
253 Bay site near the outlet of the Lavaca River system (TCEQ-13563), a mid-Lavaca Bay  
254 site (TCEQ-13383), and the lower Lavaca Bay site near the mouth of the Bay (TCEQ-  
255 13384). For estuary locations, we obtained data for TP, Nitrite+Nitrate ( $\text{NO}_x$ ), TKN,  
256 chlorophyll-*a*, and DO concentrations from January 2005 through December 2020  
257 (Table 2).  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276



**Fig. 1** Map of Lavaca Bay and the contribution watershed. The freshwater sites are the most downstream freshwater stream locations with water quality and streamflow data used for nutrient load models. Water quality concentration data at the three Lavaca Bay sites were used to assess relationships between freshwater flows, loads and estuary water quality.

**Table 1** Summary of gauged streamflow and freshwater water quality samples between January 1, 2000 and December 31, 2020.

Station ID	Parameter	Mean	SD	N
USGS-08164000	TP (mg/L)	0.21	0.09	80
	NO <sub>3</sub> (mg/L)	0.18	0.24	74
	Mean Daily Streamflow (cfs)	332.78	1667.47	7671
USGS-08164525	TP (mg/L)	0.20	0.08	81
	NO <sub>3</sub> (mg/L)	0.29	0.26	62
	Mean Daily Streamflow (cfs)	666.14	2957.79	7671

**Table 2** Summary of estuary water quality samples collected between January 1, 2005 and December 31, 2020.

Station ID	Parameter	Mean	SD	N
TCEQ-13383	TP (mg/L)	0.11	0.05	47
	NO <sub>x</sub> (mg/L)	0.07	0.15	51
	TKN (mg/L)	0.94	0.49	45
	Chlorophyll- <i>a</i> (ug/L)	9.43	5.31	47
	DO (mg/L)	7.22	1.35	55
TCEQ-13384	TP (mg/L)	0.08	0.03	51
	NO <sub>x</sub> (mg/L)	0.06	0.08	52
	TKN (mg/L)	0.76	0.40	48
	Chlorophyll- <i>a</i> (ug/L)	8.22	6.44	46
	DO (mg/L)	7.51	1.32	54
TCEQ-13563	TP (mg/L)	0.13	0.06	50
	NO <sub>x</sub> (mg/L)	0.09	0.13	53
	TKN (mg/L)	0.94	0.37	49
	Chlorophyll- <i>a</i> (ug/L)	9.67	5.33	49
	DO (mg/L)	7.91	1.34	56

## 2.2 Estimating Watershed Based Nutrient Loads

Estimates of NO<sub>3</sub> and TP loads at the Lavaca River (USGS-08164000) and the outlet of Lake Texana on the Navidad River (USGS-08164525) were developed using GAMs relating nutrient concentration to river discharge, season, and time. Separate models were fit at each station for each parameter and used to predict nutrient concentrations for each day in the study period. GAMs were fit using the *mgcv* package in R which makes available multiple types of smooth functions with automatic smoothness selection (Wood, 2011). The general form of the model related NO<sub>3</sub> or TP concentration



to a long term tend, season, streamflow, and two different antecedent discharge terms,  
shown in Eq. 1:

$$g(\mu) = \alpha + f_1(ddate) + f_2(yday) + f_3(\log1p(Q)) + f_4(ma) + f_5(fa),$$

$$y \sim \mathcal{N}(\mu, \sigma^2) \quad (1)$$

where  $\mu$  is the conditional expected  $\text{NO}_3$  or TP concentration,  $g()$  is the log-link,  $\alpha$  is the intercept,  $f_n()$  are smoothing functions.  $y$  is the response variable ( $\text{NO}_3$  or TP concentration) modeled as normally distributed with mean  $\mu$  and standard deviation  $\sigma$ .  $ddate$  is the date converted to decimal notation,  $yday$  is numeric day of year (1-366), and  $\log1p(Q)$  is the natural log of mean daily streamflow plus 1.

Moving average ( $ma$ ) is an exponentially smoothed moving average that attempts to incorporate the influence of prior streamflow events on concentration at the current time period (Wang et al, 2011; Kuhnert et al, 2012; Zhang and Ball, 2017), using Eq. 2:

$$ma(\delta) = d\kappa_{i-1} + (1 - \delta)\hat{q}_{i-1},$$

$$\kappa_i = \sum_{m=1}^i \hat{Q}_m \quad (2)$$

where  $\delta$  is the discount factor (here, set equal to 0.95),  $\kappa_i$  is the cumulative flow ( $Q$ ) up to the  $i$ th day.

Flow anomaly ( $fa$ ) is a unitless term that represents how wet or dry the current time period is from a previous time period (Vecchia et al, 2009; Zhang and Ball, 2017). Long-term flow anomaly ( $ltfa$ ) is the streamflow over the previous year relative to the entire period (Zhang and Ball, 2017) and calculated using Eq. 3:

$$ltfa(t) = \bar{x}_{1\text{ year}}(t) - \bar{x}_{entire\ period} \quad (3)$$

and the short-term flow anomaly (*stfa*) calculated as the current day flow compared to the preceding 1-month streamflow using 4:

$$stfa(t) = x_{current\ day}(t) - \bar{x}_{1\ month}(t) \quad (4)$$

where  $x$  are the averages of log-transformed streamflow over the antecedent period (*1-year*, *1-month*, etc.) for time  $t$ . We used *ltfa* in  $\text{NO}_3$  models and *stfa* in TP models based on previous work demonstrating major improvements in  $\text{NO}_x$  regression models that incorporated *ltfa* and moderate improvements in TP regression models that incorporated *stfa* (Zhang and Ball, 2017). Moving averages and flow anomalies were calculated with the *adc* R package (Schramm, 2023).

The calculation of model terms for the Lake Texana site were modified because daily loads are not a function of natural stream flow processes alone, but of dam releases and nutrient concentrations at the discharge point of the lake.  $Q$ ,  $ma$ , and  $fa$  terms were calculated based on total gaged inflow from the 4 major tributaries to the lake. Thin-plate regression splines were used for  $ddate$ ,  $\log_{1p}(Q)$ ,  $fa$ , and  $ma$ . A cyclic cubic regression spline was used for  $yday$  to ensure the ends of the spline match (day 1 and day 366 are expected to match). First order penalties were applied to the smooths of flow-based variables which penalize departures from a flat function to help constrain extrapolations for high flow measurements.

Left-censored data were not uncommon in this dataset. Several methods are available to account for censored data. We transformed left-censored nutrient concentrations to one-half the detection limit. Although this simple approach can introduce bias (Hornung and Reed, 1990), we considered it acceptable because high concentrations and loadings are associated with high-flow events and low-flow/low-concentration events will account for a small proportion of total loadings (McDowell et al, 2021).

Daily loads were estimated as the predicted concentration multiplied by the daily streamflow. For the Navidad River (USGS-08164525) site, daily loads at the dam were calculated from the discrete daily concentration at the discharge point of the lake and corresponding reported daily discharge from the dam. Flow-normalized loads were estimated similar to WRTDS by setting flow-based covariates on each day of the year equal to each of the historical values for that day of the year over the study period (Hirsch et al, 2010). The flow-normalized estimate was calculated as the mean of all the predictions for each day considering all possible flow values. Standard deviations and 90% credible intervals were obtained by drawing samples from the multivariate normal posterior distribution of the fitted GAM (Wood, 2006; Marra and Wood, 2012; McDowell et al, 2021). GAM performance was evaluated using repeated 5-fold cross validation (Burman, 1989) and average Nash-Sutcliffe Efficiency (NSE), Pearson sample correlation ( $r$ ) and percent bias (PBIAS) metrics across folds were calculated for each model.

### 2.3 Linking Estuary Water Quality to Hydrology and Nutrient Loads

To test if changes in freshwater inflow and nutrient loading had explanatory effect on changes in estuary water quality a series of GAM models were fit at each site relating parameter concentration to temporal trends (Eq. 5), temporal trends and inflow (Eq. 6), and temporal trends, inflow, and nutrient loads (Eq. 7):

$$g(\mu) = \alpha + f_1(ddate) + f_2(yday), \quad (5)$$

$$g(\mu) = \alpha + f_1(ddate) + f_2(yday) + f_3(Q), \quad (6)$$

$$g(\mu) = \alpha + f_1(ddate) + f_2(yday) + f_3(Q) + f_4(Load) \quad (7)$$

**Table 3** Set of GAM models specified for each water quality parameter response.

Parameter	Model	Model Terms
TP	Temporal	$s(ddate) + s(yday)$
	Flow	$s(ddate) + s(yday) + s(Q)$
	Flow+Load	$s(ddate) + s(yday) + s(Q) + s(TP \text{ Load})$
NO <sub>x</sub>	Temporal	$s(ddate) + s(yday)$
	Flow	$s(ddate) + s(yday) + s(Q)$
	Flow+Load	$s(ddate) + s(yday) + s(Q) + s(NO_3 \text{ Load})$
Chlorophyll- <i>a</i>	Temporal	$s(ddate) + s(yday)$
	Flow	$s(ddate) + s(yday) + s(Q)$
	Flow+Load	$s(ddate) + s(yday) + s(Q) + s(TP \text{ Load}) + s(NO_3 \text{ Load})$
Dissolved Oxygen	Temporal	$s(ddate) + s(yday)$
	Flow	$s(ddate) + s(yday) + s(Q)$
	Flow+Load	$s(ddate) + s(yday) + s(Q) + s(TP \text{ Load}) + s(NO_3 \text{ Load})$
TKN	Temporal	$s(ddate) + s(yday)$
	Flow	$s(ddate) + s(yday) + s(Q)$

where  $\mu$  is the conditional expected response (nutrient concentration),  $g()$  is the log link, and response variable was modeled as Gamma distributed with mean  $\mu$  and scale  $\lambda$ .  $f_1(ddate)$  is decimal date smoothed with a thin-plate regression spline,  $f_2(yday)$  is the numeric day of year smoothed with a cyclic cubic regression spline,  $f_3(Q)$  is mean daily inflow (the combined measurements from Lavaca River and Navidad River) and  $f_4(Load)$  is the total NO<sub>3</sub> or TP watershed load. The set of models specified for each water quality response are in Table 3.

Because streamflow and nutrient loads are tightly correlated, freshwater inflow can mask signals from nutrient loads alone. Following the methodology implemented by Murphy et al (2022), both freshwater inflow and nutrient loads were preprocessed to account for season and streamflow respectively. Raw freshwater inflow values were replaced by seasonally adjusted log transformed inflow obtained from the residuals of a GAM model fit between season(day of year) and log transformed daily freshwater inflow. Raw nutrient loads were replaced with flow-adjusted values obtained from the residuals of a GAM model relating log transformed NO<sub>3</sub> or TP loads to log transformed daily inflow. Response residuals from the respective GAM models were used as  $Q$  and  $Load$  in Eq. 6 and Eq. 7.


This study used an information theoretic approach to evaluate evidence of model covariate effects on Lavaca Bay water quality. Model probabilities were calculated and compared using the AIC<sub>c</sub> scores between each group of temporal, inflow, and inflow+load models (Burnham et al, 2011). Improvements in model probabilities provide evidence that the terms explain additional variation in the response variable.

## 3 Results

### 3.1 Watershed Nutrient Loads


Predictive performance of nutrient loads ranged from “satisfactory” to “very good” based on standardized evaluation metrics of NSE,  $r$ , and PBIAS (Moriassi et al, 2015) calculated using 5-fold cross validation. Median goodness-of-fit metrics for NO<sub>3</sub> models in the Lavaca River were 0.34 NSE, 0.70  $r$ , and 2.00 PBIAS. Navidad River NO<sub>3</sub> models appeared to perform slightly better with 0.48 NSE and 0.87  $r$  but with higher bias at 10.90 PBIAS. Generally, TP models performed better than NO<sub>3</sub> models. Median goodness-of-fit metrics for TP in the Lavaca River were 0.81 NSE, 0.93  $r$ , and -7.20 PBIAS. Navidad River TP models has similar performance with 0.91 NSE, 0.99  $r$ , and -3.30 PBIAS. Density plots of metrics show similar distribution of values between sites for the same parameter, with the exception  $r$  values for NO<sub>3</sub> loads where Lavaca River had a much larger variance in values compared to the Navidad River (Figure 2). TP GAMS had higher average NSE and  $r$  values and lower variance in metric values compared to NO<sub>3</sub>.

Annual NO<sub>3</sub> and TP loads show considerable variation, generally following patterns in discharge (Figures 3, 4). Flow-normalized TP loads at both sites and flow-normalized NO<sub>3</sub> loads in the Lavaca River indicated watershed based loads did not change much over time when accounting for variation driven by streamflow (Figure 3). Flow-normalized loads in the Lavaca River showed small variation over time with some decreases in NO<sub>3</sub> loads since 2013.



**Fig. 2** Density plots of goodness-of-fit metrics (NSE,  $r$ , and PBIAS) from repeated 5-fold cross validation between predicted nutrient loads from GAM models and measured nutrient loads. Color indicates the tail probability calculated from the empirical cumulative distribution of the goodness-of-fit metrics.

Aggregated across both sites, the mean annual  $\text{NO}_3$  load 2005 through 2020 was 205,405 kg (126,867 kg - 341,569 kg, 90% CI). Annual  $\text{NO}_3$  loads ranged from 12,574 kg in 2011 to 794,510 kg in 2007. Total annual TP loads ranged from 7,839 kg in 2011 to 595,075 kg in 2007. Mean annual TP loading from 2005 through 2020 was 182,673 kg (152,227 kg - 219,310 kg, 90% CI). On average, the Navidad River accounted for 68% of  $\text{NO}_3$  loads and 59% of TP loads from 2005 through 2020. However, during periods of extreme drought the Lavaca River became the primary source of nutrient loading in the watershed with the Navidad River only accounting for 15% and 25% of  $\text{NO}_3$  and TP loads in 2011 (Figure 4).




**Fig. 3** Aggregated estimated annual and flow-normalized annual  $\text{NO}_3$  and TP loads for USGS-08164000 and USGS-08164525.

### 3.2 Linkages Between Water Quality and Watershed Flows and Loads

There is no evidence of long-term changes in TP or DO concentrations at any Lavaca Bay site (Figure 5). The upper-bay site, TCEQ-13563, shows evidence of a long-term linear increase in  $\text{NO}_x$  while chlorophyll-*a* decreased from 2005 through 2014 (Figure 5).  $\text{NO}_x$  concentration at the mid-bay site, TCEQ-13383, displayed an unusual periodic pattern that is indicative of a strong influence from inflow or precipitation. The temporal GAMs did not provide evidence of long-term trends in any of the water quality constituents at the lower-bay TCEQ-13384 site.


Freshwater inflow provided additional explanation for changes in TP and  $\text{NO}_x$  concentration at all of the Lavaca Bay sites according to  $\text{AIC}_c$  and model probability values (Table 4). TCEQ-13563, the site closest to the river outlet, was the only



**Fig. 4** Comparison of delivered annual loads and annual discharge at the Lavaca (USGS-08164000) and Navidad (USGS-08164525) Rivers.

site that had improvements in the explanations of DO and TKN concentration with the inclusion of inflow. Both TCEQ-13563 and TCEQ-13383, the mid-bay site, saw improvements in explanations for variations in chlorophyll-*a* with the inclusion of freshwater inflow. The addition of nutrient loads (both TP and NO<sub>3</sub>) terms did not provide additional explanation for changes in chlorophyll-*a* or DO concentrations. Inclusion of TP loads provided additional explanation of TP concentrations at the upper- and mid-bay sites, TCEQ-13563 and TCEQ-13383. Inclusion of NO<sub>3</sub> loads





**Fig. 5** Fitted splines (shaded regions indicate 90% confidence intervals) from the temporal estuary GAM display the marginal smoothed effect of date on TP (A), NO<sub>x</sub> (B), chlorophyll-*a* (C), TKN (D), and DO (E) concentrations at each site in Lavaca Bay.


only provided marginal improvements in the explanation of NO<sub>x</sub> concentration at the lower-bay TCEQ-13384 site.

GAMs showed that increases in freshwater inflow resulted in nearly linear increases in TP and NO<sub>x</sub> concentration at all three sites (Figure 6). At the upper-bay TCEQ-13563 site, GAMs showed that increases in freshwater inflow initially increased chlorophyll-*a* and DO concentration but concentrations leveled and potentially decreased at higher flows. The mid-bay TCEQ-13383 site showed a nearly linear increase in chlorophyll-*a* concentration in response to increases freshwater inflow. Freshwater flow did not have significant effects on chlorophyll-*a*, TKN, or DO at the lower-bay TCEQ-13384 site.


Increased TP loads resulted in nearly linear increases of TP concentration at the upper- and mid-bay sites, TCEQ-13563 and TCEQ-13383 respectively (Figure 7).

**Table 4** Estuary GAM AIC<sub>c</sub> values and associated model probabilities (in parenthesis). Models with the highest probability for each site and water quality parameter combination are bolded and italicized for emphasis.

Parameter	Site	Temporal	Inflow	Inflow + Load
TP	TCEQ-13383	-152.1 (0.03)	-156.1 (0.24)	<b>-158.2 (0.72)</b>
	TCEQ-13384	-194.4 (0.03)	<b>-200.2 (0.49)</b>	-200.2 (0.49)
	TCEQ-13563	-145.3 (0)	-156.6 (0.41)	<b>-157.3 (0.59)</b>
NO <sub>x</sub>	TCEQ-13383	-218.9 (0)	<b>-244.8 (0.5)</b>	-244.8 (0.5)
	TCEQ-13384	-263.4 (0)	-311.7 (0.48)	<b>-311.9 (0.52)</b>
	TCEQ-13563	-175.1 (0)	<b>-190.2 (0.5)</b>	-190.2 (0.5)
Chlorophyll- <i>a</i>	TCEQ-13383	279.7 (0.18)	<b>278.1 (0.41)</b>	278.1 (0.41)
	TCEQ-13384	<b>268.2 (0.33)</b>	268.2 (0.33)	268.2 (0.33)
	TCEQ-13563	289.5 (0.08)	<b>286.1 (0.46)</b>	286.1 (0.46)
TKN	TCEQ-13383	<b>42.2 (0.66)</b>	43.5 (0.34)	-
	TCEQ-13384	<b>34.3 (0.57)</b>	34.8 (0.43)	-
	TCEQ-13563	31.1 (0.22)	<b>28.7 (0.78)</b>	-
DO	TCEQ-13383	<b>146.4 (0.34)</b>	146.4 (0.34)	146.5 (0.32)
	TCEQ-13384	<b>135.9 (0.47)</b>	137 (0.27)	137 (0.27)
	TCEQ-13563	138.3 (0.25)	<b>137.2 (0.43)</b>	137.8 (0.32)



**Fig. 6** Fitted splines from estuary GAMs display the marginal smoothed effect of freshwater inflow (controlled for season) on TP (A), NO<sub>x</sub> (B), chlorophyll-*a* (C), TKN (D), and DO (E) concentrations at each site in Lavaca Bay.



**Fig. 7** Fitted splines from the nutrient loading GAMs display the marginal smoothed effect of TP and NO<sub>3</sub> loads on TP (A) and NO<sub>x</sub> (B) concentrations at each site in Lavaca Bay.

The relative effect size appeared to much smaller than the effect of freshwater inflow alone. Increased NO<sub>3</sub> loads only showed an effect at the lower-bay TCEQ-13384 site. The effect was quite small compared to streamflow and provided only small improvements to the model (Table 4). As noted above, nutrient loadings did not provide any explanation in changes in the remaining assessed water quality parameters.

## 4 Discussion

TP and NO<sub>3</sub> loadings from the Lavaca Bay watershed showed high inter-annual variability driven primarily by fluctuations in discharge. Notably, there were no indications of trends in flow-normalized NO<sub>3</sub> and TP loads in the Navidad River. Freshwater discharges in the Navidad River are regulated by the Palmetto Dam forming Lake Texana at the lower extent of the river. While the dominant agricultural land uses differ between the Lavaca (primarily grazed pasture and rangeland) and Navidad (mix of pasture and row crops) catchments, we did not have a reason to expect different flow normalized trends between the two systems from land use alone. Lentic nitrogen uptake and cycling may have regulating effects that mask changes in upstream nitrogen loadings. Additional nutrient data collection in the contributing tributaries

875 of Lake Texana is needed to fully assess the role of Lake Texana in regulating nutri-  
876 ent delivery to the Lavaca Bay system. However, these results suggest that there have  
877 been no changes in the  $\text{NO}_3$  or TP loading from the Navidad River system at the Lake  
878 Texana discharge point when accounting for variations in year to year discharge.  
879

881       There is some evidence of recent decreases in flow-normalized  $\text{NO}_3$  loads in the  
882 Lavaca River. Although there is no work directly correlating water quality planning  
883 and implementation efforts in the watershed to water quality outcomes, efforts to  
884 increase agricultural producer participation in the watershed have been ongoing since  
885 2016 ([Schramm et al, 2018](#); [Berthold et al, 2021](#)). The decrease in flow-normalized  $\text{NO}_3$   
886 loads could be a reflection of those collective efforts but the lack of evidence for similar  
887 changes in flow-normalized TP loads provide contrary support. The inconsistent flow-  
888 normalized trends may also reflect some of the weakness of the water quality dataset  
889 that is primarily composed of ambient water quality measurements.  
890

891       Cross-validation of the nutrient loading models highlights that predictions are  
892 prone to high bias, owing to the lack of targeted storm or flow biased measurements.  
893 The high biases are indicative that subsets of values were unable to capture the func-  
894 tional relationships with the flow based dependent variables. It was beyond the scope  
895 of the current study to evaluate the subsets of cross-validation data and scores. How-  
896 ever, the cross-validation procedure is indicative that more robust sampling would  
897 be beneficial for reducing prediction variance. Supplementary flow-biased monitoring  
898 targeting storm- or high-flow conditions is critical to improve model performance and  
899 strength of evidence produced by these models ([Horowitz, 2003](#); [Snelder et al, 2017](#)).  
900

901       Some prior studies have generated estimates of mean annual TP yields in the  
902 Lavaca River watershed (Table 5, [Dunn, 1996](#); [Rebich et al, 2011](#); [Omani et al, 2014](#);  
903 [Wise et al, 2019](#)). Although these studies differ in time periods and methodologies,  
904 they provide a sanity check for the reasonableness of the annual estimates generated  
905 in the current study. In a regional assessment of nutrient loading in river's along the  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920

Gulf of Mexico, [Dunn \(1996\)](#) used the LOADEST model to develop an estimated mean annual yield of 28.9 kg/km<sup>2</sup>. LOADEST is a multiple linear regression model that fits log transformed pollutant concentrations to long term, seasonal, and flow based predictors and includes methods for bias correction when exponentiating the response variable. While the annual yield estimated in [Dunn \(1996\)](#) is substantially lower than the estimate of 35.2 kg/km<sup>2</sup> in this study, it is important to note that both hydrology and land use was substantially different for a proportion of the study period. For almost the first ten years of the study period in [Dunn \(1996\)](#), the Navidad River was still undammed. More recently the eastern portion of the watershed has seen large conversions of rice fields to more traditional row crop production ([Kulat et al, 2019](#)). Studies of long-term precipitation patterns indicate both the total amount ([Dixon and Moore, 2011](#)) and variability of rainfall ([Mishra and Singh, 2010](#); [Fagnant et al, 2020](#)) has increased in the region which provide plausible reasons for increased runoff driven nutrient loading.

[Rebich et al \(2011\)](#) and [Wise et al \(2019\)](#) used SPARROW to provide recent assessments of regional catchment based loadings to the Gulf of Mexico (Table 5). SPARROW is a hybrid statistical-process model with the underlying nutrient load estimation methods based on the previously described LOADEST ([Schwarz et al, 2006](#)). The functional form of the LOADEST regression model is similar to the terms applied in the GAMs used in the current study. The only study to apply a mechanistic watershed model (SWAT) to estimate nutrient loadings in the Lavaca River watershed [Omani et al \(2014\)](#) developed estimated yields (42 kg/km<sup>2</sup>) similar to the two SPARROW models. The base periods used by [Rebich et al \(2011\)](#), [Omani et al \(2014\)](#) and [Wise et al \(2019\)](#) differ from the current study, making direct comparisons difficult. However, the ranges of estimated yields suggest the current estimates of TP loading are reasonable. Similar results would be expected given all of the previous studies have also relied primarily on ambient water quality measurements instead of flow and storm biased measurements.

**Table 5** Comparisons of previously published estimates of mean annual TP yield at the Lavaca River site.

Reported Yield (kg·km <sup>2</sup> ·year <sup>-1</sup> )	Approach	Time Period	Reference
35.2 (28.8, 43.3) <sup>a</sup>	GAM	2005-2020	This work
45.2	SPARROW	2000-2014	Wise et al (2019)
42	SWAT	1977-2005	Omani et al (2014)
20.81-91.58 <sup>b</sup>	SPARROW	1980-2002	Rebich et al (2011)
28.9	LOADEST	1972-1993	Dunn (1996)

<sup>a</sup> Mean of the annual point estimates and the lower and upper 95% credible intervals.

<sup>b</sup> Represents a binned value range from a choropleth map.

The non-linear estuary water quality trends identified in the current study differed slightly from previously identified trends (Bugica et al, 2020). This is not unexpected due to the different time periods, different methodology, and generally small slopes identified for most of the significant water quality parameters in prior work. Both DO and chlorophyll-*a* concentrations at all three Lavaca Bay sites were stable from 2005 through 2020. This is a positive outcome in comparison to other Texas estuaries that are facing larger demands for freshwater diversions, higher population growth, and more intense agricultural production which have resulted in more direct signs of eutrophication (Wetz et al, 2016; Bugica et al, 2020). Despite the stability of DO and chlorophyll-*a*, there are concerning site specific increases in NO<sub>x</sub> and TKN concentration over the same time period. These trends are especially concerning due to the nitrogen limitation identified in many Texas estuaries (Gardner et al, 2006; Hou et al, 2012; Dorado et al, 2015; Paudel et al, 2019; Wetz et al, 2017) and the relatively low ambient concentrations observed in Lavaca Bay.

The strong positive effect of freshwater inflow on NO<sub>x</sub>, TKN, and TP concentration are suggestive of nonpoint watershed sources, consistent with watershed uses and with other studies relating freshwater inflow with nutrient concentrations in Lavaca Bay and other estuaries (Russell et al, 2006; Caffrey et al, 2007; Peierls et al, 2012; Palmer and Montagna, 2015; Cira et al, 2021). Inflow had a non-linear relationship with TKN at the two upstream sites, with TKN increasing as freshwater inflow transitioned from

low to moderate levels. At higher freshwater inflows, the effect was attenuated, possibly indicating a flushing effect at higher freshwater inflow. No relationship between TKN and freshwater inflow were observed at TCEQ-13384 located in the lower reach of Lavaca Bay. The results suggest that tidal flushing maybe diluting TKN and act as a control on the effects of freshwater inflow in the lower reaches of Lavaca Bay. The results are also consistent with previous work that suggest processing of organic loads in the upper Lavaca Bay or tidal portions of the Lavaca River reduce transport of nutrients to the lower reaches of Lavaca Bay (Russell et al, 2006).

Freshwater inflow also displayed a strong positive effect on chlorophyll-*a* at the upper- and mid-bay sites. The upper-bay site, TCEQ-13563, showed decreases in chlorophyll-*a* at the highest freshwater inflow volumes. Freshwater flushing or increases in turbidity are associated with decreases in chlorophyll-*a* in other estuaries (Peierls et al, 2012; Cloern et al, 2014). No relationships between inorganic nitrogen or TP loadings with chlorophyll-*a* were observed. Due to the lack of TKN loading information, no assessment between organic nitrogen loads and chlorophyll-*a* were possible.

Although other studies have identified complex relationships between estuary nutrient concentrations, nutrient loading and chlorophyll-*a* concentrations in Texas estuaries (Örnólfssdóttir et al, 2004; Dorado et al, 2015; Cira et al, 2021; Tominack and Wetz, 2022), this study specifically used flow-adjusted freshwater derived nutrient loads to parse out contributions from changes in nutrient loadings while accounting for variations in load due to flow. Loading GAMs indicated no evidence of changes in flow-normalized TP loads in either river, and no changes in flow-normalized NO<sub>3</sub> loads in the Navidad River. The small changes in flow-normalized NO<sub>3</sub> loads in the Lavaca River are probably masked under most conditions by discharge from the Navidad River. Given the relatively small variation in flow-normalized loads, it can be expected that they would contribute little to the variance in downstream water quality.

1059     There was no evidence that adjusted freshwater inflow and nutrient loads had  
1060 effects on DO concentration in Lavaca Bay. The seasonality term in the temporal  
1061 GAM models explained a substantial amount of DO variation at all of the sites.  
1062 Responses of estuary metabolic processes and resulting DO concentrations can be  
1063 quite complicated and often locally specific ([Caffrey, 2004](#)). While the lack of total  
1064 nitrogen or TKN loading data hinders interpretation, the large seasonal effect on DO  
1065 concentration indicates physical factors (such as temperature, wind, and turbidity)  
1066 play an important role and should be included in future models. Prior work suggests  
1067 that Lavaca Bay may not be limited by nutrients alone, with high turbidity or nutrient  
1068 processing in upper portions of the Bay or intertidal river limiting production ([Russell  
1075 et al, 2006](#)). Finally, it is reasonable to assume that fluctuations in DO may not  
1076 occur immediately in response to nutrient pulses or freshwater inflow. Work has  
1077 shown that many water quality parameters may have lagged effects lasting days or  
1078 even months following storms and large discharge events ([Mooney and McClelland,  
1082 2012](#); [Wetz and Yoskowitz, 2013](#); [Bukaveckas et al, 2020](#); [Walker et al, 2021](#)). However,  
1083 this study only evaluated responses to loading and inflows occurring the day of water  
1084 quality observations.

1087     The GAM approach proved useful for both estimating loads and assessing down-  
1088 stream responses in water quality. Although we did not compare other models, it  
1089 is likely similar estimates of loadings would be obtained by methods suggest as  
1090 LOADEST, WRTDS, or SPARROW given the functionally similar dependent vari-  
1091 able structures. The underlying weakness in the estimates of loading in the current  
1092 study is the reliance on ambient water quality data used for statewide water quality  
1093 assessments. These data do not frequently include storm or flow biased samples which  
1094 is needed to improve model performance. Although there is existing work on the sam-  
1095 ples sizes required for reliable performance of both LOADEST ([Park and Engel, 2014](#))  
1096 and WRTDS ([Kumar et al, 2019](#)) models, similar work does not appear to have been  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104



extended to water quality applications of GAMs. Due to the concerning increases in eutrophication associated parameters in Lavaca Bay and other Texas estuaries (Bugica et al, 2020), and the increasing need to better link environmental outcomes with on the ground management actions (Schramm et al, 2022) there is a strong requirement for reliable estimates of pollutant loadings and responses along the Texas coast.

To date, state wide water quality monitoring programs have focused on collection of ambient condition data. A framework for establishing pollutant load monitoring programs across catchments that explicitly incorporate flow biased data is needed for assessing estuary health along the Texas coast. Additional efforts focused on identifying relevant effect sizes, sampling designs, and funding mechanisms that can support long term efforts are also needed to adequately design such a framework. Large long-term monitoring programs in and around the Chesapeake Bay, San Francisco Bay, and along the Mississippi River have proven extremely effective at informing management actions and tracking progress towards long-term pollutant reduction goals. Similar coordinated efforts across Texas coastal watersheds would provide useful for resource management efforts intended to protect biological and water quality integrity of Texas's estuaries.

## 5 Conclusion

Reliable estimates of nutrient loads for the Lavaca Bay were obtained using GAMs. However, additional flow-biased data collection efforts is needed to decrease the prediction variance and improve accuracy at critical high-flow loading events. While some ongoing projects will fill data gaps for total nitrogen and TKN loading, additional efforts are needed to coordinate data collection efforts specifically for load estimation across Texas estuaries. This study, consistent with others along the Texas coast, found strong effects of freshwater flow on nutrient and chlorophyll-*a* concentrations. DO concentrations, dominated by seasonal effects, did not show strong direct responses

1151 to freshwater flow. Small variance in flow-adjusted nutrient loads indicates that (1)  
1152 there have been limited changes in non-point sources of nutrients and (2) that there  
1153 isn't strong evidence that those small changes have had effects on chlorophyll-*a* or dis-  
1154 solved oxygen in Lavaca Bay. Although the study did not identify strong responses to  
1155 changes in nutrient loading, this does provide a baseline assessment for future water  
1156 quality management activities in the watershed.

1161 **Acknowledgments.** The author extends thanks to Dr. Mike Wetz (Harte Research  
1162 Institute, Texas A&M Corpus Christi), Chad Kinsfather and Partick Brzozowski  
1163 (Lavaca-Navidad River Authority), Brian Koch (Texas State Soil and Water Con-  
1164 servation Board), Bill Balboa (Matagorda Bay Foundation), Jason Pinchbeck (Texas  
1165 General Land Office) and the Lavaca Bay Foundation for supporting development of  
1166 this project and providing valuable feedback.

## 1173 **Declarations**

1175 **Funding.** This project was funded in part by a Texas Coastal Management Program  
1176 grant approved by the Texas Land Commissioner, providing financial assistance under  
1177 the Coastal Zone Management Act of 1972, as amended, awarded by the National  
1178 Oceanic and Atmospheric Administration (NOAA), Office for Coastal Management,  
1179 pursuant to NOAA Award No. NA21NOS4190136. The views expressed herein are  
1180 those of the author(s) and do not necessarily reflect the views of NOAA, the U.S.  
1181 Department of Commerce, or any of their subagencies.

1187 **Data Availability.** Reproducible code and datasets generated during this study are  
1188 available in the Zenodo repository, <https://doi.org/10.5281/zenodo.733075>.

## References

- Beck MW, Murphy RR (2017) Numerical and qualitative contrasts of two statistical models for water quality change in tidal waters. *JAWRA Journal of the American Water Resources Association* 53(1):197–219. <https://doi.org/10.1111/1752-1688.12489>
- Beck MW, Jabusch TW, Trowbridge PR, et al (2018) Four decades of water quality change in the upper San Francisco Estuary. *Estuarine, Coastal and Shelf Science* 212:11–22. <https://doi.org/10.1016/j.ecss.2018.06.021>
- Bergbusch NT, Hayes NM, Simpson GL, et al (2021) Unexpected shift from phytoplankton to periphyton in eutrophic streams due to wastewater influx. *Limnology and Oceanography* 66(7):2745–2761. <https://doi.org/10.1002/lno.11786>
- Berthold TA, Olsovsky T, Schramm MP (2021) Direct mailing education campaign impacts on the adoption of grazing management practices. *Journal of Contemporary Water Research & Education* 174:45–60. <https://doi.org/10.1111/j.1936-704X.2021.3360.x>
- Beseres Pollack J, Palmer T, Montagna P (2011) Long-term trends in the response of benthic macrofauna to climate variability in the Lavaca-Colorado Estuary, Texas. *Marine Ecology Progress Series* 436:67–80. <https://doi.org/10.3354/meps09267>
- Biagi K, Ross C, Oswald C, et al (2022) Novel predictors related to hysteresis and baseflow improve predictions of watershed nutrient loads: An example from Ontario’s lower Great Lakes basin. *Science of The Total Environment* 826:154023. <https://doi.org/10.1016/j.scitotenv.2022.154023>
- Bricker S, Longstaff B, Dennison W, et al (2008) Effects of nutrient enrichment in the nation’s estuaries: A decade of change. *Harmful Algae* 8(1):21–32. <https://doi.org/>

1243 [10.1016/j.hal.2008.08.028](https://doi.org/10.1016/j.hal.2008.08.028)  
1244  
1245 Bugica K, Sterba-Boatwright B, Wetz MS (2020) Water quality trends in Texas estu-  
1246 aries. *Marine Pollution Bulletin* 152:110903. [https://doi.org/10.1016/j.marpolbul.](https://doi.org/10.1016/j.marpolbul.2020.110903)  
1247 [2020.110903](https://doi.org/10.1016/j.marpolbul.2020.110903)  
1248  
1249  
1250  
1251 Bukaveckas PA, Tassone S, Lee W, et al (2020) The influence of storm events on  
1252 metabolism and water quality of riverine and estuarine segments of the James,  
1253 Mattaponi, and Pamunkey Rivers. *Estuaries and Coasts* 43(7):1585–1602. [https:](https://doi.org/10.1007/s12237-020-00819-9)  
1254 [//doi.org/10.1007/s12237-020-00819-9](https://doi.org/10.1007/s12237-020-00819-9)  
1255  
1256  
1257  
1258  
1259 Burman P (1989) A comparative study of ordinary cross-validation, v-fold cross-  
1260 validation and the repeated learning-testing methods. *Biometrika* 76(3):503–514.  
1261 <https://doi.org/10.1093/biomet/76.3.503>  
1262  
1263  
1264  
1265 Burnham KP, Anderson DR, Huyvaert KP (2011) AIC model selection and multimodel  
1266 inference in behavioral ecology: Some background, observations, and compar-  
1267 isons. *Behavioral Ecology and Sociobiology* 65(1):23–35. [https://doi.org/10.1007/](https://doi.org/10.1007/s00265-010-1029-6)  
1268 [s00265-010-1029-6](https://doi.org/10.1007/s00265-010-1029-6)  
1269  
1270  
1271  
1272 Caffrey JM (2004) Factors controlling net ecosystem metabolism in U.S. estuaries.  
1273 *Estuaries* 27(1):90–101. <https://doi.org/10.1007/BF02803563>  
1274  
1275  
1276 Caffrey JM, Chapin TP, Jannasch HW, et al (2007) High nutrient pulses, tidal mixing  
1277 and biological response in a small California estuary: Variability in nutrient concen-  
1278 trations from decadal to hourly time scales. *Estuarine, Coastal and Shelf Science*  
1279 71(3-4):368–380. <https://doi.org/10.1016/j.ecss.2006.08.015>  
1280  
1281  
1282  
1283  
1284 Chin T, Beecraft L, Wetz MS (2022) Phytoplankton biomass and community com-  
1285 position in three Texas estuaries differing in freshwater inflow regime. *Estuarine,*  
1286 *Coastal and Shelf Science* 277:108059. <https://doi.org/10.1016/j.ecss.2022.108059>  
1287  
1288

Cira EK, Palmer TA, Wetz MS (2021) Phytoplankton dynamics in a low-inflow estuary (Baffin Bay, TX) during drought and high-rainfall conditions associated with an El Niño event. <i>Estuaries and Coasts</i> 44(7):1752–1764. <a href="https://doi.org/10.1007/s12237-021-00904-7">https://doi.org/10.1007/s12237-021-00904-7</a>	1289 1290 1291 1292 1293 1294 1295
Cloern JE, Foster SQ, Kleckner AE (2014) Phytoplankton primary production in the world’s estuarine-coastal ecosystems. <i>Biogeosciences</i> 11(9):2477–2501. <a href="https://doi.org/10.5194/bg-11-2477-2014">https://doi.org/10.5194/bg-11-2477-2014</a>	1296 1297 1298 1299 1300 1301
De Cicco LA, Hirsch RM, Lorenz DL, et al (2022) dataRetrieval: R packages for discovering and retrieving water data available from Federal hydrologic web services. U.S. Geological Survey, <a href="https://doi.org/10.5066/P9X4L3GE">https://doi.org/10.5066/P9X4L3GE</a>	1302 1303 1304 1305 1306 1307
Dixon RW, Moore TW (2011) Trend detection in Texas temperature and precipitation. <i>Southwest Geography</i> 15:80–103	1308 1309 1310 1311
Dorado S, Booe T, Steichen J, et al (2015) Towards an understanding of the interactions between freshwater inflows and phytoplankton communities in a subtropical estuary in the Gulf of Mexico. <i>PLOS ONE</i> 10(7):e0130931. <a href="https://doi.org/10.1371/journal.pone.0130931">https://doi.org/10.1371/journal.pone.0130931</a>	1312 1313 1314 1315 1316 1317 1318 1319
Dunn D (1996) Trends in Nutrient Inflows to the Gulf of Mexico from Streams Draining the Conterminous United States, 1972-93. Water-Resources Investigations Report 96-4113, USGS, Austin, Texas, <a href="https://doi.org/10.3133/wri964113">https://doi.org/10.3133/wri964113</a>	1320 1321 1322 1323 1324 1325
Fagnant C, Gori A, Sebastian A, et al (2020) Characterizing spatiotemporal trends in extreme precipitation in Southeast Texas. <i>Natural Hazards</i> 104(2):1597–1621. <a href="https://doi.org/10.1007/s11069-020-04235-x">https://doi.org/10.1007/s11069-020-04235-x</a>	1326 1327 1328 1329 1330 1331 1332 1333 1334

1335 Gardner WS, McCarthy MJ, An S, et al (2006) Nitrogen fixation and dissimilatory  
 1336 nitrate reduction to ammonium (DNRA) support nitrogen dynamics in Texas estu-  
 1337 aries. *Limnology and Oceanography* 51(1part2):558–568. [https://doi.org/10.4319/](https://doi.org/10.4319/lo.2006.51.1_part.2.0558)  
 1338 [lo.2006.51.1\\_part.2.0558](https://doi.org/10.4319/lo.2006.51.1_part.2.0558)  
 1339  
 1340  
 1341  
 1342 Hagemann M, Asce SM, Kim D, et al (2016) Estimating Nutrient and Organic Carbon  
 1343 Loads to Water-Supply Reservoir Using Semiparametric Models. *J Environ Eng* p 9.  
 1344 [https://doi.org/10.1061/\(ASCE\)EE.1943-7870.0001077](https://doi.org/10.1061/(ASCE)EE.1943-7870.0001077)  
 1345  
 1346  
 1347  
 1348 Hayes NM, Haig HA, Simpson GL, et al (2020) Effects of lake warming on the seasonal  
 1349 risk of toxic cyanobacteria exposure. *Limnology and Oceanography Letters* 5(6):393–  
 1350 402. <https://doi.org/10.1002/lol2.10164>  
 1351  
 1352  
 1353  
 1354 Hirsch RM, Moyer DL, Archfield SA (2010) Weighted Regressions on Time, Dis-  
 1355 charge, and Season (WRTDS), with an application to Chesapeake Bay River inputs.  
 1356 *JAWRA Journal of the American Water Resources Association* 46(5):857–880.  
 1357 <https://doi.org/10.1111/j.1752-1688.2010.00482.x>  
 1358  
 1359  
 1360  
 1361 Hornung RW, Reed LD (1990) Estimation of average concentration in the presence of  
 1362 nondetectable values. *Applied Occupational and Environmental Hygiene* 5(1):46–51.  
 1363 <https://doi.org/10.1080/1047322X.1990.10389587>  
 1364  
 1365  
 1366  
 1367 Horowitz AJ (2003) An evaluation of sediment rating curves for estimating suspended  
 1368 sediment concentrations for subsequent flux calculations. *Hydrological Processes*  
 1369 17(17):3387–3409. <https://doi.org/10.1002/hyp.1299>  
 1370  
 1371  
 1372  
 1373 Hou L, Liu M, Carini SA, et al (2012) Transformation and fate of nitrate near  
 1374 the sediment–water interface of Copano Bay. *Continental Shelf Research* 35:86–94.  
 1375 <https://doi.org/10.1016/j.csr.2012.01.004>  
 1376  
 1377  
 1378  
 1379  
 1380

Jain S, Schramm MP (2021) Technical Support Document for One Total  
Maximum Daily Load for Indicator Bacteria in Lavaca River Above  
Tidal. Technical Report AS-221, Texas Commission on Environmen-  
tal Quality, Austin, Texas, URL [https://www.tceq.texas.gov/downloads/  
water-quality/tmdl/lavaca-river-above-tidal-rocky-creek-recreational-108/  
108-lavaca-river-addendum-tsd-2021-october-as-221.pdf](https://www.tceq.texas.gov/downloads/water-quality/tmdl/lavaca-river-above-tidal-rocky-creek-recreational-108/108-lavaca-river-addendum-tsd-2021-october-as-221.pdf)

Kennicutt MC (2017) Water Quality of the Gulf of Mexico. In: Ward CH (ed) Habitats  
and Biota of the Gulf of Mexico: Before the Deepwater Horizon Oil Spill. Springer  
New York, New York, NY, p 55–164, [https://doi.org/10.1007/978-1-4939-3447-8\\_2](https://doi.org/10.1007/978-1-4939-3447-8_2)

Kroon FJ, Kuhnert PM, Henderson BL, et al (2012) River loads of suspended solids,  
nitrogen, phosphorus and herbicides delivered to the Great Barrier Reef lagoon.  
Marine Pollution Bulletin 65(4-9):167–181. [https://doi.org/10.1016/j.marpolbul.  
2011.10.018](https://doi.org/10.1016/j.marpolbul.2011.10.018)

Kuhnert PM, Henderson BL, Lewis SE, et al (2012) Quantifying total suspended sedi-  
ment export from the Burdekin River catchment using the loads regression estimator  
tool. Water Resources Research 48(4). <https://doi.org/10.1029/2011WR011080>

Kulat MI, Mohtar RH, Olivera F (2019) Holistic Water-Energy-Food Nexus for  
Guiding Water Resources Planning: Matagorda County, Texas Case. Frontiers in  
Environmental Science 7:3. <https://doi.org/10.3389/fenvs.2019.00003>

Kumar S, Godrej A, Post H, et al (2019) The value of intensive sampling—a com-  
parison of fluvial loads. Water Resources Management 33(12):4303–4318. <https://doi.org/10.1007/s11269-019-02369-7>

Lloyd C, Freer J, Collins A, et al (2014) Methods for detecting change in hydrochemical  
time series in response to targeted pollutant mitigation in river catchments. Journal

1427 of Hydrology 514:297–312. <https://doi.org/10.1016/j.jhydrol.2014.04.036>  
 1428  
 1429 Marra G, Wood SN (2012) Coverage properties of confidence intervals for Generalized  
 1430 Additive Model components: Coverage properties of GAM intervals. Scandinavian  
 1431 Journal of Statistics 39(1):53–74. <https://doi.org/10.1111/j.1467-9469.2011.00760.x>  
 1432  
 1433 McDowell RW, Simpson ZP, Ausseil AG, et al (2021) The implications of lag times  
 1434 between nitrate leaching losses and riverine loads for water quality policy. Scientific  
 1435 Reports 11(1):16450. <https://doi.org/10.1038/s41598-021-95302-1>  
 1436  
 1437 Mishra AK, Singh VP (2010) Changes in extreme precipitation in Texas. Journal of  
 1438 Geophysical Research 115(D14):D14106. <https://doi.org/10.1029/2009JD013398>  
 1439  
 1440 Montagna PA, Cockett PM, Kurr EM, et al (2020) Assessment of the Relationship  
 1441 Between Freshwater Inflow and Biological Indicators in Lavaca Bay. Tech. Rep.  
 1442 Contract # 1800012268, Harte Research Institute, Texas A&M University-Corpus  
 1443 Christi, Corpus Christi, Texas  
 1444  
 1445 Mooney RF, McClelland JW (2012) Watershed export events and ecosystem responses  
 1446 in the Mission–Aransas National Estuarine Research Reserve, South Texas. Estu-  
 1447 aries and Coasts 35(6):1468–1485. <https://doi.org/10.1007/s12237-012-9537-4>  
 1448  
 1449 Moriasi DN, Gitau MW, Pai N, et al (2015) Hydrologic and Water Quality Mod-  
 1450 els: Performance Measures and Evaluation Criteria. Transactions of the ASABE  
 1451 58(6):1763–1785. <https://doi.org/10.13031/trans.58.10715>  
 1452  
 1453 Murphy RR, Perry E, Harcum J, et al (2019) A Generalized Additive Model approach  
 1454 to evaluating water quality: Chesapeake Bay case study. Environmental Modelling  
 1455 & Software 118:1–13. <https://doi.org/10.1016/j.envsoft.2019.03.027>  
 1456  
 1457  
 1458  
 1459  
 1460  
 1461  
 1462  
 1463  
 1464  
 1465  
 1466  
 1467  
 1468  
 1469  
 1470  
 1471  
 1472



Murphy RR, Keisman J, Harcum J, et al (2022) Nutrient improvements in Chesapeake Bay: Direct effect of load reductions and implications for coastal management.	1473
Environmental Science & Technology 56(1):260–270. <a href="https://doi.org/10.1021/acs.est.1c05388">https://doi.org/10.1021/acs.est.1c05388</a>	1474
	1475
	1476
	1477
	1478
	1479
Oelsner GP, Stets EG (2019) Recent trends in nutrient and sediment loading to coastal areas of the conterminous U.S.: Insights and global context. Science of The Total Environment 654:1225–1240. <a href="https://doi.org/10.1016/j.scitotenv.2018.10.437">https://doi.org/10.1016/j.scitotenv.2018.10.437</a>	1480
	1481
	1482
	1483
	1484
	1485
Omani N, Srinivasan R, Lee T (2014) Estimation of sediment and nutrient loads to bays from gauged and ungauged watersheds. Applied Engineering in Agriculture pp 869–887. <a href="https://doi.org/10.13031/aea.30.10162">https://doi.org/10.13031/aea.30.10162</a>	1486
	1487
	1488
	1489
	1490
	1491
Örnólfssdóttir EB, Lumsden S, Pinckney JL (2004) Nutrient pulsing as a regulator of phytoplankton abundance and community composition in Galveston Bay, Texas. Journal of Experimental Marine Biology and Ecology 303(2):197–220. <a href="https://doi.org/10.1016/j.jembe.2003.11.016">https://doi.org/10.1016/j.jembe.2003.11.016</a>	1492
	1493
	1494
	1495
	1496
	1497
	1498
	1499
Palmer TA, Montagna PA (2015) Impacts of droughts and low flows on estuarine water quality and benthic fauna. Hydrobiologia 753(1):111–129. <a href="https://doi.org/10.1007/s10750-015-2200-x">https://doi.org/10.1007/s10750-015-2200-x</a>	1500
	1501
	1502
	1503
	1504
	1505
Park Y, Engel B (2014) Use of pollutant load regression models with various sampling frequencies for annual load estimation. Water 6(6):1685–1697. <a href="https://doi.org/10.3390/w6061685">https://doi.org/10.3390/w6061685</a>	1506
	1507
	1508
	1509
	1510
	1511
Paudel B, Montagna PA, Adams L (2019) The relationship between suspended solids and nutrients with variable hydrologic flow regimes. Regional Studies in Marine Science 29:100657. <a href="https://doi.org/10.1016/j.rsma.2019.100657">https://doi.org/10.1016/j.rsma.2019.100657</a>	1512
	1513
	1514
	1515
	1516
	1517
	1518

1519 Peierls BL, Hall NS, Paerl HW (2012) Non-monotonic responses of phytoplankton  
 1520 biomass accumulation to hydrologic variability: A comparison of two coastal plain  
 1521 North Carolina estuaries. *Estuaries and Coasts* 35(6):1376–1392. [https://doi.org/](https://doi.org/10.1007/s12237-012-9547-2)  
 1522 [10.1007/s12237-012-9547-2](https://doi.org/10.1007/s12237-012-9547-2)  
 1523  
 1524  
 1525  
 1526 Rebich RA, Houston NA, Mize SV, et al (2011) Sources and delivery of nutrients to  
 1527 the northwestern Gulf of Mexico from streams in the south-central United States.  
 1528 JAWRA Journal of the American Water Resources Association 47(5):1061–1086.  
 1529 <https://doi.org/10.1111/j.1752-1688.2011.00583.x>  
 1530  
 1531  
 1532  
 1533  
 1534 Robson BJ, Dourdet V (2015) Prediction of sediment, particulate nutrient and  
 1535 dissolved nutrient concentrations in a dry tropical river to provide input to a mecha-  
 1536 nistic coastal water quality model. *Environmental Modelling & Software* 63:97–108.  
 1537 <https://doi.org/10.1016/j.envsoft.2014.08.009>  
 1538  
 1539  
 1540  
 1541 Russell MJ, Montagna PA, Kalke RD (2006) The effect of freshwater inflow on net  
 1542 ecosystem metabolism in Lavaca Bay, Texas. *Estuarine, Coastal and Shelf Science*  
 1543 68(1-2):231–244. <https://doi.org/10.1016/j.ecss.2006.02.005>  
 1544  
 1545  
 1546  
 1547 Schramm M (2023) Adc: Calculate Antecedant Discharge Conditions. URL [https:](https://CRAN.R-project.org/package=adc)  
 1548 [//CRAN.R-project.org/package=adc](https://CRAN.R-project.org/package=adc)  
 1549  
 1550  
 1551 Schramm M, Berthold A, Entwistle C, et al (2018) Lavaca River Watershed Pro-  
 1552 tection Plan. Technical Report TR-507, Texas Water Resources Institute, Col-  
 1553 lege Station, Texas, URL [https://twri.tamu.edu/publications/technical-reports/](https://twri.tamu.edu/publications/technical-reports/2018-technical-reports/tr-507/)  
 1554 [2018-technical-reports/tr-507/](https://twri.tamu.edu/publications/technical-reports/2018-technical-reports/tr-507/)  
 1555  
 1556  
 1557  
 1558  
 1559 Schramm M, Gitter A, Gregory L (2022) Total Maximum Daily Loads and *Escherichia*  
 1560 *coli* trends in Texas freshwater streams. *Journal of Contemporary Water Research*  
 1561 & Education 176:36–49. <https://doi.org/10.1111/j.1936-704X.2022.3374.x>  
 1562  
 1563  
 1564

Schwarz GE, Hoos AB, Alexander RB, et al (2006) The SPARROW Surface Water-	1565
Quality Model: Theory, Application and User Documentation. URL <a href="https://doi.org/10.3133/tm6B3">https://doi.</a>	1566
<a href="https://doi.org/10.3133/tm6B3">org/10.3133/tm6B3</a>	1567
	1568
	1569
	1570
Shoda ME, Sprague LA, Murphy JC, et al (2019) Water-quality trends in U.S. rivers,	1571
2002 to 2012: Relations to levels of concern. Science of The Total Environment	1572
650:2314–2324. <a href="https://doi.org/10.1016/j.scitotenv.2018.09.377">https://doi.org/10.1016/j.scitotenv.2018.09.377</a>	1573
	1574
	1575
	1576
Snelder T, McDowell R, Fraser C (2017) Estimation of catchment nutrient loads in	1577
New Zealand using monthly water quality monitoring data. JAWRA Journal of	1578
the American Water Resources Association 53(1):158–178. <a href="https://doi.org/10.1111/1752-1688.12492">https://doi.org/10.1111/</a>	1579
<a href="https://doi.org/10.1111/1752-1688.12492">1752-1688.12492</a>	1580
	1581
	1582
	1583
Stackpoole S, Sabo R, Falcone J, et al (2021) Long-term Mississippi River	1584
trends expose shifts in the river load response to watershed nutrient balances	1585
between 1975 and 2017. Water Resources Research 57(11). <a href="https://doi.org/10.1029/2021WR030318">https://doi.org/10.1029/</a>	1586
<a href="https://doi.org/10.1029/2021WR030318">2021WR030318</a>	1587
	1588
	1589
	1590
	1591
Stets E, Lee C, Lytle D, et al (2018) Increasing chloride in rivers of the contermi-	1592
nous U.S. and linkages to potential corrosivity and lead action level exceedances	1593
in drinking water. Science of The Total Environment 613–614:1498–1509. <a href="https://doi.org/10.1016/j.scitotenv.2017.07.119">https://doi.org/10.1016/j.scitotenv.2017.07.119</a>	1594
	1595
	1596
	1597
	1598
	1599
Tominack SA, Wetz MS (2022) Variability in phytoplankton biomass and community	1600
composition in Corpus Christi Bay, Texas. Estuaries and Coasts <a href="https://doi.org/10.1007/s12237-022-01137-y">https://doi.org/</a>	1601
<a href="https://doi.org/10.1007/s12237-022-01137-y">10.1007/s12237-022-01137-y</a>	1602
	1603
	1604
Vecchia AV, Gilliom RJ, Sullivan DJ, et al (2009) Trends in concentrations and use	1605
of agricultural herbicides for Corn Belt Rivers, 1996-2006. Environmental Science	1606
& Technology 43(24):9096–9102. <a href="https://doi.org/10.1021/es902122j">https://doi.org/10.1021/es902122j</a>	1607
	1608
	1609
	1610

1611 Walker LM, Montagna PA, Hu X, et al (2021) Timescales and magnitude of water  
 1612 quality change in three Texas estuaries induced by passage of Hurricane Harvey.  
 1613 Estuaries and Coasts 44(4):960–971. <https://doi.org/10.1007/s12237-020-00846-6>  
 1614  
 1615  
 1616 Wang YG, Kuhnert P, Henderson B (2011) Load estimation with uncertainties from  
 1617 opportunistic sampling data – A semiparametric approach. Journal of Hydrology  
 1618 396(1-2):148–157. <https://doi.org/10.1016/j.jhydrol.2010.11.003>  
 1619  
 1620  
 1621  
 1622 Wazniak CE, Hall MR, Carruthers TJB, et al (2007) Linking water quality to  
 1623 living resources in a Mid-Atlantic lagoon system, USA. Ecological Applications  
 1624 17(sp5):S64–S78. <https://doi.org/10.1890/05-1554.1>  
 1625  
 1626  
 1627  
 1628 Wetz MS, Yoskowitz DW (2013) An ‘extreme’ future for estuaries? Effects of extreme  
 1629 climatic events on estuarine water quality and ecology. Marine Pollution Bulletin  
 1630 69(1-2):7–18. <https://doi.org/10.1016/j.marpolbul.2013.01.020>  
 1631  
 1632  
 1633  
 1634 Wetz MS, Hayes KC, Fisher KV, et al (2016) Water quality dynamics in an urbaniz-  
 1635 ing subtropical estuary(Oso Bay, Texas). Marine Pollution Bulletin 104(1-2):44–53.  
 1636  
 1637 <https://doi.org/10.1016/j.marpolbul.2016.02.013>  
 1638  
 1639  
 1640 Wetz MS, Cira EK, Sterba-Boatwright B, et al (2017) Exceptionally high organic  
 1641 nitrogen concentrations in a semi-arid South Texas estuary susceptible to brown tide  
 1642 blooms. Estuarine, Coastal and Shelf Science 188:27–37. [https://doi.org/10.1016/j.](https://doi.org/10.1016/j.ecss.2017.02.001)  
 1643  
 1644  
 1645  
 1646  
 1647  
 1648 Wise DR, Anning DW, Miller OW (2019) Spatially referenced models of stream-  
 1649 flow and nitrogen, phosphorus, and suspended-sediment transport in streams of  
 1650 the southwestern United States. Scientific Investigations Report 2019-5106, U.S.  
 1651 Geological Survey, Reston, Virginia  
 1652  
 1653  
 1654  
 1655  
 1656

Wood SN (2006) On confidence intervals for generalized additive models based	1657
on penalized regression splines. Australian & New Zealand Journal of Statistics	1658
48(4):445–464. <a href="https://doi.org/10.1111/j.1467-842X.2006.00450.x">https://doi.org/10.1111/j.1467-842X.2006.00450.x</a>	1659
	1660
	1661
	1662
Wood SN (2011) Fast stable restricted maximum likelihood and marginal likelihood	1663
estimation of semiparametric generalized linear models: Estimation of Semipara-	1664
metric Generalized Linear Models. Journal of the Royal Statistical Society: Series	1665
B (Statistical Methodology) 73(1):3–36. <a href="https://doi.org/10.1111/j.1467-9868.2010.00749.x">https://doi.org/10.1111/j.1467-9868.2010.</a>	1666
<a href="https://doi.org/10.1111/j.1467-9868.2010.00749.x">00749.x</a>	1667
	1668
	1669
	1670
	1671
Zhang Q, Ball WP (2017) Improving riverine constituent concentration and flux esti-	1672
mation by accounting for antecedent discharge conditions. Journal of Hydrology	1673
547:387–402. <a href="https://doi.org/10.1016/j.jhydrol.2016.12.052">https://doi.org/10.1016/j.jhydrol.2016.12.052</a>	1674
	1675
	1676
	1677
	1678
	1679
	1680
	1681
	1682
	1683
	1684
	1685
	1686
	1687
	1688
	1689
	1690
	1691
	1692
	1693
	1694
	1695
	1696
	1697
	1698
	1699
	1700
	1701
	1702