

1 Linking watershed nutrient loading to 2 estuary water quality with semiparametric 3 models

4 Michael Schramm¹

5 ¹Texas Water Resources Institute, Texas A&M AgriLife Research, College Station, Texas,
6 United States

7 Corresponding author:

8 Michael Schramm¹

9 Email address: michael.schramm@ag.tamu.edu

10 ABSTRACT

11 Evaluating estuary water quality responses to reductions (or increases) in nutrient loading attributed to
12 on the ground management actions can be challenging due to the strong influence of environmental
13 drivers on nutrient loads and non-linear relationships. This study applied semiparametric approaches
14 to calculate watershed nutrient loads and assess responses in estuary water quality to seasonally-
15 adjusted freshwater inflow and flow-adjusted nutrient loads in Lavaca Bay, Texas. Lavaca Bay is a
16 secondary embayment on the Texas coast displaying early potential for eutrophication and water quality
17 degradation. Use of flow-adjusted nutrient loads allowed the study to evaluate the response in water
18 quality to changes in nutrient loads driven by non-natural variation. Cross-validation indicated that,
19 despite data constraints, semiparametric models performed well at nutrient load prediction. Based on
20 these models, delivered annual nutrient loads varied substantially from year to year. In contrast, minimal
21 changes in flow-normalized loads indicate that nutrient loadings were driven by natural variation in
22 precipitation and runoff as opposed to changes in management of nonpoint sources. Models indicated
23 no evidence of long-term changes in dissolved oxygen or chlorophyll-a within Lavaca Bay. However, site
24 specific long-term increases in both organic and inorganic nitrogen are concerning for their potential
25 to fuel eutrophication. Further analysis found freshwater inflow had strong influences on nutrient and
26 chlorophyll-a concentrations but there was no evidence that changes in watershed nutrient loading
27 explained additional variation in dissolved oxygen or chlorophyll-a concentrations. In addition to providing
28 a baseline assessment of watershed nutrient loading and water quality responses in the Lavaca Bay
29 watershed, this study provides methodological support for the use of semiparametric models in load
30 regression models and estuary assessments.

31 INTRODUCTION

32 Similar to many coastal areas globally, the coastal watersheds along the Texas Gulf coast are facing
33 pressures from growing populations, increases in point source and non-point source pollution and
34 alterations to freshwater flows that degrade water quality in downstream estuaries (Bricker et al., 2008;
35 Bugica et al., 2020; Kennicutt, 2017). Despite these escalating pressures, national scale assessments have
36 classified coastal estuaries in Texas as moderate or low risk for eutrophic conditions (Bricker et al., 2008).
37 However, a suite of recent studies indicates that estuary water quality dynamics in both agricultural and
38 urban dominated watersheds within Texas are expressing conditions that are increasingly conducive to
39 algal blooms and eutrophication (Bugica et al., 2020; Chin et al., 2022; Wetz et al., 2016, 2017). With
40 identification of several localized areas of estuary water quality concern along the Texas coast (Bugica
41 et al., 2020), localized studies are being prioritized to better inform management actions.

42 The goal of this project is to assess watershed nutrient loading and the resulting water quality responses
43 in Lavaca Bay, Texas. Lavaca Bay is a secondary bay in the larger Matagorda Bay system located roughly
44 halfway between Houston, Texas and Corpus Christi, Texas. Lavaca Bay faces substantial challenges
45 associated with legacy contamination but general water quality parameters such as dissolved oxygen (DO),
46 nutrients, and biological parameters have been well within state water quality standards. Despite largely

meeting state designated water quality thresholds, there have been concerning declines in abundance, biomass, and diversity of benthic fauna in Lavaca Bay (Beseres Pollack et al., 2011). These declines are partially attributed to reductions in freshwater inflow and changes in estuary salinity and are indicative of an already stressed system (Beseres Pollack et al., 2011; Montagna et al., 2020; Palmer & Montagna, 2015). More recently, significant linear increases in total phosphorus (TP), orthophosphate, total Kjeldahl nitrogen (TKN), and chlorophyll-*a* concentrations were identified at monitoring sites within Lavaca Bay (Bugica et al., 2020). These trends in nutrient concentrations are concerning due to the role of nitrogen as a limiting factor for primary production in many Texas estuaries (Dorado et al., 2015; Gardner et al., 2006; Hou et al., 2012; Paudel et al., 2019; Wetz et al., 2017) and the ramifications that changes in nitrogen loadings could have for productivity and eutrophication in Lavaca Bay.

There are ongoing efforts between local, state, and federal agencies to address water quality impairments in the freshwater portions of the Lavaca Bay watershed (Berthold et al., 2021; Jain & Schramm, 2021; Schramm et al., 2018). However, on a statewide scale, these approaches have shown limited success and emphasize a need for improved methods to assess and link management actions with downstream water quality (Schramm et al., 2022).

Such methods could help identify and replicate effective water quality management actions across the state. The identification and communication of changes and trends in water quality is complicated by the fact that trends are often non-linear and confounded by precipitation and runoff that hinder traditional analysis (Lloyd et al., 2014; Wazniak et al., 2007). The development and application of flexible statistical methods such as Weighted Regressions on Time, Discharge and Season (WRTDS, Hirsch et al., 2010) and Generalized Additive Models (GAMs, Wood, 2011) have provided effective tools for researchers to quantify and communicate non-linear changes in river and estuary pollutant loadings.

WRTDS calculates a time series of in-stream concentrations or loads (daily, monthly, or annually) and flow-normalized estimates of concentrations and loads using locally weighted regression for unique combinations of time, discharge, and season. WRTDS has been widely used to assess and identify trends in riverine nutrients (Oelsner & Stets, 2019; Stackpoole et al., 2021), chlorides (Stets et al., 2018), and other pollutants of concern (Shoda et al., 2019). WRTDS has also been successfully adapted to assess trends in estuary water quality concentrations (Beck et al., 2018).

While WRTDS is a statistical approach developed specifically for water quality applications, GAMs are a broadly applicable statistical method. GAMs are a semiparametric extension of generalized linear models where the linear predictor is represented as the sum of multiple unknown smooth functions and parametric linear predictors (Wood, 2011). Although the underlying parameter estimation procedure of GAMs are substantially different than WRTDS, both the functional form and results have been demonstrated to be similar when assessing nutrient concentration trends (Beck & Murphy, 2017). Water quality applications of GAMs have included river and catchment nutrient concentration and load estimation (Biagi et al., 2022; Hagemann et al., 2016; Kroon et al., 2012; Kuhnert et al., 2012; McDowell et al., 2021; Robson & Dourdet, 2015; Wang et al., 2011), and assessment of temporal trends of nutrients (Beck & Murphy, 2017; Murphy et al., 2019), phytoplankton (Bergbusch et al., 2021), and cyanobacteria (Hayes et al., 2020). Recently GAMs have also been used to link water quality responses in receiving water bodies to changes in nonpoint source nutrient inputs (Murphy et al., 2022). For a substantial discussion on the differences (and similarities) between GAMs and WRTDS for water quality applications readers are referred to Beck and Murphy (2017).

To provide actionable information for resource managers in Lavaca Bay, water quality conditions must be evaluated relative to changes in natural environmental drivers to better understand and manage potential human impacts. This study utilizes GAMs to develop estimates of delivered and flow-normalized nutrient loads and assess estuary water quality responses to changes in loads delivered to Lavaca Bay. GAMs were chosen over other regression-based approaches for use in this study due to: (1) the ability to easily explore and incorporate different model terms; (2) the incorporation of non-linear smooth functions that do not require explicit *a priori* knowledge of the expected shape; and (3) inclusion of a link function that related the expected value of the response to linear predictors thus avoiding unneeded data transformations and bias corrections. The exploratory study also assesses the response of water quality parameters in Lavaca Bay over time and in response to freshwater inflow controlled for seasonality and to watershed nutrient loads that are controlled for environmentally driven variation.

Table 1. Summary of gauged streamflow and freshwater water quality samples between January 1, 2000 and December 31, 2020.

Station ID	Parameter	Mean	SD	N
USGS-08164000	TP (mg/L)	0.21	0.09	80
	NO ₃ (mg/L)	0.18	0.24	74
	Mean Daily Streamflow (cfs)	332.78	1667.47	7671
USGS-08164525	TP (mg/L)	0.20	0.08	81
	NO ₃ (mg/L)	0.29	0.26	62
	Mean Daily Streamflow (cfs)	666.14	2957.79	7671

METHODS AND MATERIALS

Location and Data

Lavaca Bay is a 190 km² estuary with the majority of freshwater inflow provided by the Lavaca and Navidad River systems (Fig. 1). The Garcitas-Arenosa, Placedo Creek, and Cox Bay subwatersheds provide additional freshwater inflows. The entire watershed area is 8,149 km² and primarily rural. Watershed land cover and land use is 50% grazed pasture and rangeland, 20% cultivated cropland (primarily row crops such as corn, cotton, and sorghum), and 5% suburban/urban. Pasture and rangeland is concentrated in the Lavaca River watershed, while cultivated crops are generally located along the eastern tributaries of the Navidad river. The Lavaca and Navidad River watersheds are a combined 5,966 km², or approximately 73% of the entire Lavaca Bay watershed area. Discharge from the Navidad River is regulated by Lake Texana which has been in operation since 1980. Lake Texana provides 0.210 km³ of water storage and discharges into the tidal section of the Navidad River which ultimately joins the tidal section of the Lavaca River 15 km upstream of the confluence with the Lavaca Bay.

Daily discharges for the Lavaca River (USGS-08164000, Fig. 1) were obtained from the United States Geologic Survey (USGS) National Water Information System using the *dataRetrieval* R package (De Cicco et al., 2022). Gaged daily discharges from the outlet of Lake Texana on the Navidad River (USGS-0816425) were provided by the Texas Water Development Board (TWDB) (April 21, 2022 email from R. Neupane, TWDB).

Water quality data for the two freshwater and three estuary locations were obtained from the Texas Commission on Environmental Quality (TCEQ) Surface Water Quality Monitoring Information System. Data submitted through the system are required to be collected under Quality Assurance Project Plans and lab method procedures outlined by the TCEQ's procedures manual. These operating procedures ensure consistent collection and laboratory methods are applied between samples collected by different entities and under different projects. All sites had varying time periods and availability of water quality data. For freshwater locations, TP from January 2000 through December 2020 and nitrate-nitrogen (NO₃) data from January 2005 through December 2020 were downloaded (Table 1). Less than 5-years of total nitrogen and TKN concentration data were available at the freshwater sites and deemed insufficient to develop load estimation models (Horowitz, 2003; Snelder et al., 2017). The three estuary sites included an upper Lavaca Bay site near the outlet of the Lavaca River system (TCEQ-13563), a mid-Lavaca Bay site (TCEQ-13383), and the lower Lavaca Bay site near the mouth of the Bay (TCEQ-13384). For estuary locations, we obtained data for TP, Nitrite+Nitrate (NO_x), TKN, chlorophyll-*a*, and DO concentrations from January 2005 through December 2020 (Table 2).

Estimating Watershed Based Nutrient Loads

Estimates of NO₃ and TP loads at the Lavaca River (USGS-08164000) and the outlet of Lake Texana on the Navidad River (USGS-08164525) were developed using GAMs relating nutrient concentration to river discharge, season, and time. Separate models were fit at each station for each parameter and used to predict nutrient concentrations for each day in the study period. GAMs were fit using the *mgcv* package in R which makes available multiple types of smooth functions with automatic smoothness selection (Wood, 2011). The general form of the model related NO₃ or TP concentration to a long term trend, season, streamflow, and two different antecedent discharge terms, shown in Eq. 1:




Figure 1. Map of Lavaca Bay and the contribution watershed. The freshwater sites are the most downstream freshwater stream locations with water quality and streamflow data used for nutrient load models. Water quality concentration data at the three Lavaca Bay sites were used to assess relationships between freshwater flows, loads and estuary water quality.

Table 2. Summary of estuary water quality samples collected between January 1, 2005 and December 31, 2020.

Station ID	Parameter	Mean	SD	N
TCEQ-13383	TP (mg/L)	0.11	0.05	47
	NO _x (mg/L)	0.07	0.15	51
	TKN (mg/L)	0.94	0.49	45
	Chlorophyll- <i>a</i> (ug/L)	9.43	5.31	47
	DO (mg/L)	7.22	1.35	55
TCEQ-13384	TP (mg/L)	0.08	0.03	51
	NO _x (mg/L)	0.06	0.08	52
	TKN (mg/L)	0.76	0.40	48
	Chlorophyll- <i>a</i> (ug/L)	8.22	6.44	46
	DO (mg/L)	7.51	1.32	54
TCEQ-13563	TP (mg/L)	0.13	0.06	50
	NO _x (mg/L)	0.09	0.13	53
	TKN (mg/L)	0.94	0.37	49
	Chlorophyll- <i>a</i> (ug/L)	9.67	5.33	49
	DO (mg/L)	7.91	1.34	56

$$g(\mu) = \alpha + f_1(ddate) + f_2(yday) + f_3(\log(1 + Q)) + f_4(ma) + f_5(fa),$$

$$y \sim \mathcal{N}(\mu, \sigma^2)$$
(1)

where μ is the conditional expected NO₃ or TP concentration, $g(\cdot)$ is the log-link, α is the intercept, $f_n(\cdot)$ are smoothing functions. y is the response variable (NO₃ or TP concentration) modeled as normally distributed with mean μ and standard deviation σ . $ddate$ is the date converted to decimal notation, $yday$ is numeric day of year (1-366), and Q is the mean daily streamflow.

Moving average (ma) is an exponentially smoothed moving average that attempts to incorporate the influence of prior streamflow events on concentration at the current time period (Kuhnert et al., 2012; Wang et al., 2011; Zhang & Ball, 2017), using Eq. 2:

$$ma(t) = \frac{\sum_{j=1}^J d^{t-t_j} Q_{t_j}}{\sum_{j=1}^J d^{t-t_j}}$$
(2)

where t is the day of estimation, d is the discount factor (set to 0.95), J is the total number of daily discharge observations, $t - t_j$ is the lag time between the current day and some historical day of observation (t_j). Here, as d approaches zero, the time series becomes the daily observed values and as d gets closer to one, the time series becomes the mean flow.

Flow anomaly (fa) is a unitless term that represents how wet or dry the current time period is from a previous time period (Vecchia et al., 2009; Zhang & Ball, 2017). Long-term flow anomaly ($ltfa$) is the streamflow over the previous year relative to the entire period (Eq. 3, Zhang & Ball, 2017) and the short-term flow anomaly ($stfa$) calculated as the current day flow compared to the preceding 1-month streamflow (Eq. 4, Zhang & Ball, 2017):

$$ltfa(t) = \bar{x}_{1\text{year}}(t) - \bar{x}_{\text{entire period}},$$
(3)

$$stfa(t) = x_{\text{current day}}(t) - \bar{x}_{1\text{month}}(t)$$
(4)

where x are the averages of log-transformed streamflow over the antecedent period (1-year , 1-month , etc.) for time t . We used $ltfa$ in NO₃ models and $stfa$ in TP models based on previous work demonstrating major improvements in NO_x regression models that incorporated $ltfa$ and moderate improvements in TP

159 regression models that incorporated *stfa* (Zhang & Ball, 2017). Moving averages and flow anomalies
160 were calculated with the *adc* R package (Schramm, 2023).

161 The calculation of model terms for the Lake Texana site were modified because daily loads are not a
162 function of natural stream flow processes alone, but of dam releases and nutrient concentrations at the
163 discharge point of the lake. Q , ma , and fa terms were calculated based on total gaged inflow from the 4
164 major tributaries to the lake. Thin-plate regression splines were used for $ddate$, $\log_{1p}(Q)$, fa , and ma . A
165 cyclic cubic regression spline was used for $yday$ to ensure the ends of the spline match (day 1 and day 366
166 are expected to match). First order penalties were applied to the smooths of flow-based variables which
167 penalize departures from a flat function to help constrain extrapolations for high flow measurements.

168 Left-censored data were not uncommon in this dataset. Several methods are available to account
169 for censored data. We transformed left-censored nutrient concentrations to one-half the detection limit.
170 Although this simple approach can introduce bias (Hornung & Reed, 1990), we considered it accept-
171 able because high concentrations and loadings are associated with high-flow events and low-flow/low-
172 concentration events will account for a small proportion of total loadings (McDowell et al., 2021).

173 Daily loads were estimated as the predicted concentration multiplied by the daily streamflow. For
174 the Navidad River (USGS-08164525) site, daily loads at the dam were calculated from the discrete daily
175 concentration at the discharge point of the lake and corresponding reported daily discharge from the dam.
176 Flow-normalized loads were estimated similar to WRTDS by setting flow-based covariates on each day
177 of the year equal to each of the historical values for that day of the year over the study period (Hirsch
178 et al., 2010). The flow-normalized estimate was calculated as the mean of all the predictions for each day
179 considering all possible flow values. Standard deviations and 90% credible intervals were obtained by
180 drawing samples from the multivariate normal posterior distribution of the fitted GAM (Marra & Wood,
181 2012; McDowell et al., 2021; Wood, 2006). GAM performance was evaluated with repeated 5-fold cross
182 validation (Burman, 1989) using average Nash-Sutcliffe Efficiency (NSE), Pearson sample correlation (r)
183 and percent bias (PBIAS) metrics.

184 **Linking Estuary Water Quality to Hydrology and Nutrient Loads**

185 To test if changes in freshwater inflow and nutrient loading had explanatory effect on changes in estuary
186 water quality a series of GAM models were fit at each site relating parameter concentration to temporal
187 trends (Eq. 5), temporal trends and inflow (Eq. 6), and temporal trends, inflow, and nutrient loads (Eq. 7):

$$g(\mu) = \alpha + f_1(ddate) + f_2(yday), \quad (5)$$

$$g(\mu) = \alpha + f_1(ddate) + f_2(yday) + f_3(Q), \quad (6)$$

$$g(\mu) = \alpha + f_1(ddate) + f_2(yday) + f_3(Q) + f_4(Load) \quad (7)$$

188 where μ is the conditional expected response (nutrient concentration), $g()$ is the log link, and response
189 variable was modeled as Gamma distributed with mean μ and scale λ . $f_1(ddate)$ is decimal date
190 smoothed with a thin-plate regression spline, $f_2(yday)$ is the numeric day of year smoothed with a cyclic
191 cubic regression spline, $f_3(Q)$ is mean daily inflow (the combined measurements from Lavaca River and
192 Navidad River) and $f_4(Load)$ is the total NO_3 or TP watershed load. The set of models specified for each
193 water quality response are in Table 3.

194 Because streamflow and nutrient loads are tightly correlated, freshwater inflow can mask signals due
195 to changes in nutrient loads alone. Following the methodology implemented by Murphy et al. (2022), both
196 freshwater inflow and nutrient loads were preprocessed to account for season and streamflow respectively.
197 Raw freshwater inflow values were replaced by seasonally adjusted log transformed inflow obtained from
198 the residuals of a GAM model fit between season(day of year) and log transformed daily freshwater inflow.
199 Raw nutrient loads were replaced with flow-adjusted values obtained from the residuals of a GAM model
200 relating log transformed NO_3 or TP loads to log transformed daily inflow. Response residuals from the
201 respective GAM models were used as Q and $Load$ in Eq. 6 and Eq. 7.

202 This study used an information theoretic approach to evaluate evidence of model covariate effects
203 on Lavaca Bay water quality. Model probabilities were calculated and compared using the AIC_c scores
204 between each group of temporal, inflow, and inflow+load models (Burnham et al., 2011). Improvements
205 in model probabilities provide evidence that the terms explain additional variation in the response variable.

Table 3. Set of GAM models specified for each water quality parameter response.

Parameter	Model	Model Terms
TP	Temporal	$s(\text{ddate}) + s(\text{yday})$
	Flow	$s(\text{ddate}) + s(\text{yday}) + s(Q)$
	Flow+Load	$s(\text{ddate}) + s(\text{yday}) + s(Q) + s(\text{TP Load})$
NO _x	Temporal	$s(\text{ddate}) + s(\text{yday})$
	Flow	$s(\text{ddate}) + s(\text{yday}) + s(Q)$
	Flow+Load	$s(\text{ddate}) + s(\text{yday}) + s(Q) + s(\text{NO}_3 \text{ Load})$
Chlorophyll- <i>a</i>	Temporal	$s(\text{ddate}) + s(\text{yday})$
	Flow	$s(\text{ddate}) + s(\text{yday}) + s(Q)$
	Flow+Load	$s(\text{ddate}) + s(\text{yday}) + s(Q) + s(\text{TP Load}) + s(\text{NO}_3 \text{ Load})$
Dissolved Oxygen	Temporal	$s(\text{ddate}) + s(\text{yday})$
	Flow	$s(\text{ddate}) + s(\text{yday}) + s(Q)$
	Flow+Load	$s(\text{ddate}) + s(\text{yday}) + s(Q) + s(\text{TP Load}) + s(\text{NO}_3 \text{ Load})$
TKN	Temporal	$s(\text{ddate}) + s(\text{yday})$
	Flow	$s(\text{ddate}) + s(\text{yday}) + s(Q)$

RESULTS

Watershed Nutrient Loads

Predictive performance of nutrient loads ranged from “satisfactory” to “very good” based on standardized evaluation metrics of NSE, r , and PBIAS (Moriassi et al., 2015) calculated using 5-fold cross validation. Median goodness-of-fit metrics for NO₃ models in the Lavaca River were 0.34 NSE, 0.70 r , and 2.00 PBIAS. Navidad River NO₃ models appeared to perform slightly better with 0.48 NSE and 0.87 r but with higher bias at 10.90 PBIAS. Generally, TP models performed better than NO₃ models. Median goodness-of-fit metrics for TP in the Lavaca River were 0.81 NSE, 0.93 r , and -7.20 PBIAS. Navidad River TP models had similar performance with 0.91 NSE, 0.99 r , and -3.30 PBIAS. Density plots of metrics show similar distribution of values between sites for the same parameter, with the exception r values for NO₃ loads where Lavaca River had a much larger variance in values compared to the Navidad River (Fig. 2). TP GAMS had higher average NSE and r values and lower variance in metric values compared to NO₃.

Annual NO₃ and TP loads show considerable variation, generally following patterns in discharge (Fig. 3, Fig. 4). Flow-normalized TP loads at both sites and flow-normalized NO₃ loads in the Lavaca River indicated watershed based loads did not change much over time when accounting for variation driven by streamflow (Fig. 3). Flow-normalized loads in the Lavaca River showed small variation over time with some decreases in NO₃ loads since 2013.

Aggregated across both sites, the mean annual NO₃ load from 2005 through 2020 was 205,405 kg (126,867 kg - 341,569 kg, 90% CI). Annual NO₃ loads ranged from 12,574 kg in 2011 to 794,510 kg in 2007. Total annual TP loads ranged from 7,839 kg in 2011 to 595,075 kg in 2007. Mean annual TP loading from 2005 through 2020 was 182,673 kg (152,227 kg - 219,310 kg, 90% CI). On average, the Navidad River accounted for 68% of NO₃ loads and 59% of TP loads from 2005 through 2020. However, during periods of extreme drought the Lavaca River became the primary source of nutrient loading in the watershed with the Navidad River only accounting for 15% and 25% of NO₃ and TP loads in 2011 (Fig. 4).

Linkages Between Water Quality and Watershed Flows and Loads

There was no evidence of long-term changes in TP or DO concentrations at any Lavaca Bay site (Fig. 5). The upper-bay site, TCEQ-13563, had evidence of a long-term linear increase in NO_x while chlorophyll-*a* decreased from 2005 through 2014 (Fig. 5). NO_x concentration at the mid-bay site, TCEQ-13383, displayed an unusual periodic pattern that is indicative of a strong influence from inflow or precipitation. The temporal GAMS did not provide evidence of long-term trends in any of the water quality constituents at the lower-bay TCEQ-13384 site.




Figure 2. Density plots of goodness-of-fit metrics (NSE, r , and PBIAS) from repeated 5-fold cross validation between predicted nutrient loads from GAM models and measured nutrient loads. Color indicates the tail probability calculated from the empirical cumulative distribution of the goodness-of-fit metrics.




Figure 3. Aggregated estimated annual and flow-normalized annual NO_3 and TP loads for USGS-08164000 and USGS-08164525.




Figure 4. Comparison of delivered annual loads and annual discharge at the Lavaca (USGS-08164000) and Navidad (USGS-08164525) Rivers.




Figure 5. Fitted splines (shaded regions indicate 90% confidence intervals) from the temporal estuary GAM display the marginal smoothed effect of date on TP (A), NO_x (B), chlorophyll-*a* (C), TKN (D), and DO (E) concentrations at each site in Lavaca Bay.

Table 4. Estuary GAM AIC_c values and associated model probabilities (in parenthesis). Models with the highest probability for each site and water quality parameter combination are bolded and italicized for emphasis.

Parameter	Site	Temporal	Inflow	Inflow + Load
TP	TCEQ-13383	-152.1 (0.03)	-156.1 (0.24)	-158.2 (0.72)
	TCEQ-13384	-194.4 (0.03)	-200.2 (0.49)	-200.2 (0.49)
	TCEQ-13563	-145.3 (0)	-156.6 (0.41)	-157.3 (0.59)
NO _x	TCEQ-13383	-218.9 (0)	-244.8 (0.5)	-244.8 (0.5)
	TCEQ-13384	-263.4 (0)	-311.7 (0.48)	-311.9 (0.52)
	TCEQ-13563	-175.1 (0)	-190.2 (0.5)	-190.2 (0.5)
Chlorophyll- <i>a</i>	TCEQ-13383	279.7 (0.18)	278.1 (0.41)	278.1 (0.41)
	TCEQ-13384	268.2 (0.33)	268.2 (0.33)	268.2 (0.33)
	TCEQ-13563	289.5 (0.08)	286.1 (0.46)	286.1 (0.46)
TKN	TCEQ-13383	42.2 (0.66)	43.5 (0.34)	-
	TCEQ-13384	34.3 (0.57)	34.8 (0.43)	-
	TCEQ-13563	31.1 (0.22)	28.7 (0.78)	-
DO	TCEQ-13383	146.4 (0.34)	146.4 (0.34)	146.5 (0.32)
	TCEQ-13384	135.9 (0.47)	137 (0.27)	137 (0.27)
	TCEQ-13563	138.3 (0.25)	137.2 (0.43)	137.8 (0.32)

Freshwater inflow provided additional explanation for changes in TP and NO_x concentration at all of the Lavaca Bay sites according to AIC_c and model probability values (Table 4). TCEQ-13563, the site closest to the river outlet, was the only site that had improvements in the explanations of DO and TKN concentration with the inclusion of inflow. Both TCEQ-13563 and TCEQ-13383, the mid-bay site, saw improvements in explanations for variations in chlorophyll-*a* with the inclusion of freshwater inflow. The addition of nutrient loads (both TP and NO₃) terms did not provide additional explanation for changes in chlorophyll-*a* or DO concentrations. Inclusion of TP loads provided additional explanation of TP concentrations at the upper- and mid-bay sites, TCEQ-13563 and TCEQ-13383. Inclusion of NO₃ loads only provided marginal improvements in the explanation of NO_x concentration at the lower-bay TCEQ-13384 site.

Increases in freshwater inflow resulted in nearly linear increases in TP and NO_x concentration at all three sites (Fig. 6). At the upper-bay TCEQ-13563 site, increases in freshwater inflow initially increased chlorophyll-*a* and DO concentration but concentrations leveled and potentially decreased at higher flows. The mid-bay TCEQ-13383 site had nearly linear increases in chlorophyll-*a* concentration in response to increased freshwater inflow. There was no evidence of freshwater inflow effects on chlorophyll-*a*, TKN, or DO at the lower-bay TCEQ-13384 site.

Increased TP loads resulted in nearly linear increases of TP concentration at the upper- and mid-bay sites, TCEQ-13563 and TCEQ-13383 respectively (Fig. 7). The relative effect size appeared much smaller than the effect of freshwater inflow alone. Increased NO₃ loads only showed an effect at the lower-bay TCEQ-13384 site. The effect was quite small compared to streamflow and provided only small improvements to the model (Table 4). As noted above, nutrient loadings did not provide any explanation in changes in the remaining assessed water quality parameters.

DISCUSSION

Nutrient Loads

TP and NO₃ loadings from the Lavaca Bay watershed showed high inter-annual variability driven primarily by fluctuations in discharge. Notably, there were no indications of trends in flow-normalized NO₃ and TP loads in the Navidad River. In comparison, there was weak evidence of more recent decreases in flow-normalized NO₃ (but not TP) loads in the Lavaca River watershed. While the dominant agricultural land uses differ between the Lavaca (primarily grazed pasture and rangeland) and Navidad (mix of pasture and row crops) catchments, we did not have a reason to expect different flow normalized trends between




Figure 6. Fitted splines from estuary GAMs display the marginal smoothed effect of freshwater inflow (controlled for season) on TP (A), NO_x (B), chlorophyll-*a* (C), TKN (D), and DO (E) concentrations at each site in Lavaca Bay.




Figure 7. Fitted splines from the nutrient loading GAMs display the marginal smoothed effect of TP and NO₃ loads on TP (A) and NO_x (B) concentrations at each site in Lavaca Bay.

the two systems from land use alone. Freshwater discharges in the Navidad River are regulated by the Palmetto Bend Dam forming Lake Texana at the lower extent of the river. Lentic nitrogen uptake and cycling may have regulating effects that mask changes in upstream nitrogen loadings. Additional nutrient data collection in the contributing tributaries of Lake Texana is needed to fully assess the role of Lake Texana in regulating nutrient delivery to the Lavaca Bay system. However, these results suggest that there have been no changes in the NO_3 or TP loading from the Navidad River system at the Lake Texana discharge point when accounting for variations in year to year discharge.

The evidence of decreased Lavaca River NO_3 loading, although weak, is a potential positive sign for water quality managers working to implement practices that improve water quality in the freshwater sections of the Lavaca River watershed. Planning and implementation efforts to increase agricultural producer participation in water quality protection practices have been ongoing in the watershed since 2016 (Berthold et al., 2021; Schramm et al., 2018), however little work has been conducted to directly link these efforts with water quality outcomes. The decrease in flow-normalized NO_3 loads could be a reflection of those collective efforts but the lack of evidence for similar changes in flow-normalized TP loads provide contrary support. The inconsistent flow-normalized trends may also reflect some of the weakness of the water quality dataset that is primarily composed of ambient water quality measurements. The issues associated with the lack of flow-biased measurements is further discussed later in this section.

Some prior studies have generated estimates of mean annual TP yields in the Lavaca River watershed (Table 5, Dunn, 1996; Omani et al., 2014; Rebich et al., 2011; Wise et al., 2019). Although these studies differ in time periods and methodologies, they provide a sanity check for the reasonableness of the annual estimates generated in the current study. In a regional assessment of nutrient loading in river's along the Gulf of Mexico, Dunn (1996) used the LOADEST model to develop an estimated mean annual yield of 28.9 kg/km^2 . LOADEST is a multiple linear regression model that fits log transformed pollutant concentrations to long term, seasonal, and flow based predictors and includes methods for bias correction when exponentiating the response variable. While the annual yield estimated in Dunn (1996) is lower than the estimate of 35.2 kg/km^2 in this study, it is important to note that both hydrology and land use was different for a proportion of the study period. For almost the first ten years of the study period in Dunn (1996), the Navidad River was still undammed. More recently the eastern portion of the watershed has seen large conversions of rice fields to more traditional row crop production (Kulat et al., 2019). The apparent increase in mean TP yield might be suggestive of wetter recent conditions. To evaluate long-term trends in discharge, we fit a GAM relating log-transformed daily discharges on the Lavaca and Navidad Rivers to season and time (Fig. 8). The long-term trends in discharge indicate watershed discharges were at or above average from 1972 through the early and mid-1980s. In comparison watershed discharges since the mid-2000s are at or below average. The trend in mean streamflow alone is counter to expectations based on changes in nutrient yield estimates. Studies of long-term precipitation patterns indicate both the total amount (Dixon & Moore, 2011) and variability of rainfall (Fagnant et al., 2020; Mishra & Singh, 2010) has increased in the region which provide plausible reasons for increased runoff driven nutrient loading. Further assessments of the frequency and occurrence of periodic high flow events that mobilize sediments and are responsible for the bulk of loading volume would provide further insight to this discrepancy.

Rebich et al. (2011) and Wise et al. (2019) used SPARROW to provide a more recent assessment of regional catchment based loadings to the Gulf of Mexico (Table 5). SPARROW is a hybrid statistical-process model with the underlying nutrient load estimation methods based on the previously described LOADEST (Schwarz et al., 2006). The functional form of the LOADEST regression model is similar to the terms applied in the GAMs used in the current study. The only study to apply a mechanistic watershed model (SWAT) to estimate nutrient loadings in the Lavaca River watershed Omani et al. (2014) developed estimated yields (42 kg/km^2) similar to the two SPARROW models. The base periods used by Rebich et al. (2011), Omani et al. (2014) and Wise et al. (2019) differ from the current study, making direct comparisons difficult. It is probable that the lower than average discharges observed from 2010 through 2021 (Fig. 8) bias our estimates downward compared to other studies which included higher than average streamflow periods. Overall, the ranges of estimated yields among different studies along with the apparent large variability in streamflow driven loadings (Fig. 3, Fig. 4) suggest that the current estimates of TP loading are reasonable.




Figure 8. Measured daily discharges (log-transformed) and smoothed long-term trends for the Lavaca and Navidad Rivers from 1972 through 2001.

Table 5. Comparisons of previously published estimates of mean annual TP yield at the Lavaca River site.

Reported Yield ($\text{kg}\cdot\text{km}^2\cdot\text{year}^{-1}$)	Approach	Time Period	Reference
35.2 (28.8, 43.3) ^a	GAM	2005-2020	This work
45.2	SPARROW	2000-2014	Wise et al., 2019
42	SWAT	1977-2005	Omani et al., 2014
20.81-91.58 ^b	SPARROW	1980-2002	Rebich et al., 2011
28.9	LOADEST	1972-1993	Dunn, 1996

^a Mean of the annual point estimates and the lower and upper 90% credible intervals.

^b Represents a binned value range from a choropleth map.

Estuary Water Quality

The non-linear estuary water quality trends identified in the current study differed slightly from previously identified trends (Bugica et al., 2020). This is not unexpected due to the different time periods, different methodology, and generally small slopes identified for most of the significant water quality parameters in prior work. Both DO and chlorophyll-*a* concentrations at all three Lavaca Bay sites were stable from 2005 through 2020. This is a positive outcome in comparison to other Texas estuaries that are facing larger demands for freshwater diversions, higher population growth, and more intense agricultural production which have resulted in more direct signs of eutrophication (Bugica et al., 2020; Wetz et al., 2016). Despite the stability of DO and chlorophyll-*a*, there are concerning site specific increases in NO_x and TKN concentration over the same time period. These trends are especially concerning due to the nitrogen limitation identified in many Texas estuaries (Dorado et al., 2015; Gardner et al., 2006; Hou et al., 2012; Paudel et al., 2019; Wetz et al., 2017) and the relatively low ambient concentrations observed in Lavaca Bay.

The strong positive effect of freshwater inflow on NO_x, TKN, and TP concentration are suggestive of nonpoint watershed sources, consistent with watershed uses and with other studies relating freshwater inflow with nutrient concentrations in Lavaca Bay and other estuaries (Caffrey et al., 2007; Cira et al., 2021; Palmer & Montagna, 2015; Peierls et al., 2012; Russell et al., 2006). Inflow had a non-linear relationship with TKN at the two upstream sites, with TKN increasing as freshwater inflow transitioned from low to moderate levels. At higher freshwater inflows, the effect was attenuated, possibly indicating a flushing effect at higher freshwater inflow. No relationship between TKN and freshwater inflow were observed at TCEQ-13384 located in the lower reach of Lavaca Bay. The results suggest that tidal flushing might dilute TKN and act as a control on the effects of freshwater inflow in the lower reaches of Lavaca Bay. The results are also consistent with previous work that suggest processing of organic loads in the upper Lavaca Bay or tidal portions of the Lavaca River reduce transport of nutrients to the lower reaches of Lavaca Bay (Russell et al., 2006).

Freshwater inflow also displayed a strong positive effect on chlorophyll-*a* at the upper- and mid-bay sites. The upper-bay site, TCEQ-13563, showed decreases in chlorophyll-*a* at the highest freshwater inflow volumes. Freshwater flushing or increases in turbidity are associated with decreases in chlorophyll-*a* in other estuaries (Cloern et al., 2014; Peierls et al., 2012). No relationships between inorganic nitrogen or TP loadings with chlorophyll-*a* were observed. Due to the lack of TKN loading information, no assessment between organic nitrogen loads and chlorophyll-*a* were possible.

Although other studies have identified complex relationships between estuary nutrient concentrations, nutrient loading and chlorophyll-*a* concentrations in Texas estuaries (Cira et al., 2021; Dorado et al., 2015; Örnólfsson et al., 2004; Tominack & Wetz, 2022), this study specifically used flow-adjusted freshwater derived nutrient loads to parse out contributions from changes in nutrient loadings while accounting for variations in load due to flow. Loading GAMs indicated no evidence of changes in flow-normalized TP loads in either river, and no changes in flow-normalized NO₃ loads in the Navidad River. The small changes in flow-normalized NO₃ loads in the Lavaca River are probably masked under most conditions by discharge from the Navidad River. Given the relatively small variation in flow-normalized loads, it can be expected that they would contribute little to the variance in downstream water quality.

There was no evidence that adjusted freshwater inflow and nutrient loads had effects on DO concentration in Lavaca Bay. The seasonality term in the temporal GAM models explained a substantial amount of DO variation at all of the sites. Responses of estuary metabolic processes and resulting DO concentrations can be quite complicated and often locally specific (Caffrey, 2004). While the lack of total nitrogen or TKN loading data hinders interpretation, the large seasonal effect on DO concentration indicates physical factors (such as temperature, wind, and turbidity) play an important role and should be included in future models. Prior work suggests that Lavaca Bay may not be limited by nutrients alone, with high turbidity or nutrient processing in upper portions of the Bay or intertidal river limiting production (Russell et al., 2006). Finally, it is reasonable to assume that fluctuations in DO may not occur immediately in response to nutrient pulses or freshwater inflow. Work has shown that many water quality parameters may have lagged effects lasting days or even months following storms and large discharge events (Bukaveckas et al., 2020; Mooney & McClelland, 2012; Walker et al., 2021; Wetz & Yoskowitz, 2013). However, this study only evaluated responses to loading and inflows occurring the day of water quality observations.

Overall, this study suggests that DO and chlorophyll-*a* concentrations have been relatively stable in Lavaca Bay. Site-specific increases in TKN and NO_x concentrations are a cause of concern for increasing

377 risks of eutrophication within Lavaca Bay which might be currently attenuated by changes in freshwater
378 flow, turbidity, and other physical processes. While loading models indicate that there are large annual
379 fluctuations in NO_3 loads, these changes have been largely driven environmental conditions (changes in
380 runoff and river discharge). These models also provide evidence that estuary NO_x and TP concentrations
381 are strongly driven by freshwater inflow and to a lesser extent fluctuation in flow-adjusted riverine loadings.
382 Site-specific changes in the relationships between freshwater inflow and responses in both chlorophyll-*a*
383 and TKN concentrations are indicative nutrient processing and or tidal flushing effects moving from the
384 river discharge point to the mouth of Lavaca Bay. This study does not completely explain site specific
385 increases in NO_x and TKN concentrations in Lavaca Bay. The freshwater study sites did not quantify
386 nutrient loadings from tidal contribution areas or ungauaged watersheds. Nutrient contributions from
387 wastewater facility discharges, septic systems, and stormwater could be considerable contributors to
388 nutrient loadings in Lavaca Bay since they are not processed by a tidal river reach prior to entering the Bay.
389 The Garcitas-Arenosa Creek, Placedo Creek, and Cox Bay subwatersheds are currently undersampled
390 but compose approximately 27% of the watershed area. The contribution of nutrient loadings from these
391 undersampled areas is unknown.

392 **Limitations**

393 The GAM approach proved useful for both estimating loads and assessing downstream responses in water
394 quality. Although we did not compare other models, it is likely similar estimates of loadings would
395 be obtained by methods suggest as LOADEST, WRTDS, or SPARROW given the functionally similar
396 dependent variable structures. The underlying weakness in the estimates of loading in the current study is
397 the reliance on ambient water quality data used for statewide water quality assessments. Cross-validation
398 of the nutrient loading models highlights that predictions are prone to high bias, owing to the lack of
399 targeted storm or flow biased measurements. The high biases are indicative that subsets of values were
400 unable to capture the functional relationships with the flow based dependent variables. It was beyond
401 the scope of the current study to evaluate the subsets of cross-validation data and scores. However, the
402 cross-validation procedure is indicative that more robust sampling is needed. Supplementary flow-biased
403 monitoring targeting storm- or high-flow conditions is critical to improve model performance and strength
404 of evidence produced by these models (Horowitz, 2003; Snelder et al., 2017). Although there is existing
405 work on the samples sizes and sample design required for reliable performance of both LOADEST (Park
406 & Engel, 2014) and WRTDS (Kumar et al., 2019) models, similar work does not appear to have been
407 extended to water quality applications of GAMs.

408 Due to the concerning increases in eutrophication associated parameters in Lavaca Bay and other Texas
409 estuaries (Bugica et al., 2020), and the desire to quantify linkages between environmental outcomes and
410 on the ground management actions (Schramm et al., 2022) there is a strong need for reliable estimates of
411 pollutant loadings and responses along the Texas coast. Within Texas, statewide water quality monitoring
412 programs have focused on collection of ambient condition data. A framework for establishing pollutant
413 load monitoring programs across catchments that explicitly incorporate flow biased data is needed for
414 assessing nutrient loading and estuary health along the Texas coast.

415 Additional efforts focused on identifying relevant effect sizes, sampling designs, and funding mecha-
416 nisms that can support long term efforts are also needed to adequately design such a framework. Large
417 long-term monitoring programs in and around the Chesapeake Bay, San Francisco Bay, and along the
418 Mississippi River have proven extremely effective at informing management actions and tracking progress
419 towards long-term pollutant reduction goals. Similar coordinated efforts across Texas coastal watersheds
420 would provide useful for resource management efforts intended to protect the biological and water quality
421 integrity of Texas's estuaries.

422 **CONCLUSIONS**

423 The primary purpose of this study was to provide estimates of watershed nutrient loadings and assess
424 water quality responses to changes in nutrient loads. GAMs provided reliable estimates of watershed
425 NO_3 and TP loads. However, additional flow-biased data collection efforts are needed to decrease the
426 prediction variance and improve accuracy at critical high-flow loading events. While some ongoing
427 projects will fill data gaps for total nitrogen and TKN loading, additional efforts are needed to coordinate
428 data collection efforts specifically for load estimation across Texas estuaries. Despite these data gaps, this
429 study identified high annual fluctuations in nutrient loads driven primarily by discharge. No evidence was

identified to indicate that on the ground management had changed nutrient loading in the Navidad River subwatershed. There was weak evidence for recent reductions in flow-normalized NO₃ loading in the Lavaca River subwatershed although the results are at odds with flow-normalized trends in TP loads.

This study, consistent with others along the Texas coast, found strong effects of freshwater flow on nutrient and chlorophyll-*a* concentrations. DO concentrations, dominated by seasonal effects, did not show strong direct responses to freshwater flow. Small variances in flow-adjusted nutrient loads indicate that (1) there have been limited changes in non-point sources of nutrients and (2) that there is not strong evidence that those small changes have had effects on chlorophyll-*a* or dissolved oxygen in Lavaca Bay. Although this study did not identify changes in DO or chlorophyll-*a* concentrations in Lavaca Bay, site specific increases in NO_x and TKN are a cause for water quality concern. The study provides a baseline assessment for future water quality management activities in the watershed. In order to effectively track and link improvements or degradation of water quality conditions in Lavaca Bay and other coastal Texas watersheds with on the ground efforts, more robust sampling networks are needed to improve spatial coverage of undersampled areas and explicitly incorporate flow-biased sampling.

ACKNOWLEDGEMENTS

The author thanks Stephanie deVilleneuve for comments on an early draft and members of the Coastal Nutrient Input Repository project committee for supporting project development and their valuable input.

REFERENCES

- Beck, M. W., Jabusch, T. W., Trowbridge, P. R., & Senn, D. B. (2018). Four decades of water quality change in the upper San Francisco Estuary. *Estuarine, Coastal and Shelf Science*, 212, 11–22. <https://doi.org/10.1016/j.ecss.2018.06.021>
- Beck, M. W., & Murphy, R. R. (2017). Numerical and qualitative contrasts of two statistical models for water quality change in tidal waters. *JAWRA Journal of the American Water Resources Association*, 53(1), 197–219. <https://doi.org/10.1111/1752-1688.12489>
- Bergbusch, N. T., Hayes, N. M., Simpson, G. L., & Leavitt, P. R. (2021). Unexpected shift from phytoplankton to periphyton in eutrophic streams due to wastewater influx. *Limnology and Oceanography*, 66(7), 2745–2761. <https://doi.org/10.1002/lno.11786>
- Berthold, T. A., Olsovsky, T., & Schramm, M. P. (2021). Direct mailing education campaign impacts on the adoption of grazing management practices. *Journal of Contemporary Water Research & Education*, 174, 45–60. <https://doi.org/10.1111/j.1936-704X.2021.3360.x>
- Beseres Pollack, J., Palmer, T., & Montagna, P. (2011). Long-term trends in the response of benthic macrofauna to climate variability in the Lavaca-Colorado Estuary, Texas. *Marine Ecology Progress Series*, 436, 67–80. <https://doi.org/10.3354/meps09267>
- Biagi, K., Ross, C., Oswald, C., Soricetti, R., Thomas, J., & Wellen, C. (2022). Novel predictors related to hysteresis and baseflow improve predictions of watershed nutrient loads: An example from Ontario's lower Great Lakes basin. *Science of The Total Environment*, 826, 154023. <https://doi.org/10.1016/j.scitotenv.2022.154023>
- Bricker, S., Longstaff, B., Dennison, W., Jones, A., Boicourt, K., Wicks, C., & Woerner, J. (2008). Effects of nutrient enrichment in the nation's estuaries: A decade of change. *Harmful Algae*, 8(1), 21–32. <https://doi.org/10.1016/j.hal.2008.08.028>
- Bugica, K., Sterba-Boatwright, B., & Wetz, M. S. (2020). Water quality trends in Texas estuaries. *Marine Pollution Bulletin*, 152, 110903. <https://doi.org/10.1016/j.marpolbul.2020.110903>
- Bukaveckas, P. A., Tassone, S., Lee, W., & Franklin, R. B. (2020). The influence of storm events on metabolism and water quality of riverine and estuarine segments of the James, Mattaponi, and Pamunkey Rivers. *Estuaries and Coasts*, 43(7), 1585–1602. <https://doi.org/10.1007/s12237-020-00819-9>
- Burman, P. (1989). A comparative study of ordinary cross-validation, v-fold cross-validation and the repeated learning-testing methods. *Biometrika*, 76(3), 503–514. <https://doi.org/10.1093/biomet/76.3.503>
- Burnham, K. P., Anderson, D. R., & Huyvaert, K. P. (2011). AIC model selection and multimodel inference in behavioral ecology: Some background, observations, and comparisons. *Behavioral Ecology and Sociobiology*, 65(1), 23–35. <https://doi.org/10.1007/s00265-010-1029-6>

- 482 Caffrey, J. M. (2004). Factors controlling net ecosystem metabolism in U.S. estuaries. *Estuaries*, 27(1),
483 90–101. <https://doi.org/10.1007/BF02803563>
- 484 Caffrey, J. M., Chapin, T. P., Jannasch, H. W., & Haskins, J. C. (2007). High nutrient pulses, tidal mixing
485 and biological response in a small California estuary: Variability in nutrient concentrations
486 from decadal to hourly time scales. *Estuarine, Coastal and Shelf Science*, 71(3-4), 368–380.
487 <https://doi.org/10.1016/j.ecss.2006.08.015>
- 488 Chin, T., Beecraft, L., & Wetz, M. S. (2022). Phytoplankton biomass and community composition in three
489 Texas estuaries differing in freshwater inflow regime. *Estuarine, Coastal and Shelf Science*, 277,
490 108059. <https://doi.org/10.1016/j.ecss.2022.108059>
- 491 Cira, E. K., Palmer, T. A., & Wetz, M. S. (2021). Phytoplankton dynamics in a low-inflow estuary (Baffin
492 Bay, TX) during drought and high-rainfall conditions associated with an El Niño event. *Estuaries
493 and Coasts*, 44(7), 1752–1764. <https://doi.org/10.1007/s12237-021-00904-7>
- 494 Cloern, J. E., Foster, S. Q., & Kleckner, A. E. (2014). Phytoplankton primary production in the world's
495 estuarine-coastal ecosystems. *Biogeosciences*, 11(9), 2477–2501. <https://doi.org/10.5194/bg-11-2477-2014>
- 496
- 497 De Ciccio, L. A., Hirsch, R. M., Lorenz, D. L., Watkins, W., & Johnson, M. (2022). dataRetrieval: R
498 packages for discovering and retrieving water data available from Federal hydrologic web
499 services. <https://doi.org/10.5066/P9X4L3GE>
- 500 Dixon, R. W., & Moore, T. W. (2011). Trend detection in Texas temperature and precipitation. *Southwest
501 Geography*, 15, 80–103.
- 502 Dorado, S., Booe, T., Steichen, J., McInnes, A. S., Windham, R., Shepard, A., Lucchese, A. E. B.,
503 Preischel, H., Pinckney, J. L., Davis, S. E., Roelke, D. L., & Quigg, A. (2015). Towards an
504 understanding of the interactions between freshwater inflows and phytoplankton communities in
505 a subtropical estuary in the Gulf of Mexico. *PLOS ONE*, 10(7), e0130931. <https://doi.org/10.1371/journal.pone.0130931>
- 506
- 507 Dunn, D. (1996). *Trends in Nutrient Inflows to the Gulf of Mexico from Streams Draining the Conterminous
508 United States, 1972-93* (Water-Resources Investigations Report No. 96-4113). USGS. Austin,
509 Texas. <https://doi.org/10.3133/wri964113>
- 510 Fagnant, C., Gori, A., Sebastian, A., Bedient, P. B., & Ensor, K. B. (2020). Characterizing spatiotemporal
511 trends in extreme precipitation in Southeast Texas. *Natural Hazards*, 104(2), 1597–1621. <https://doi.org/10.1007/s11069-020-04235-x>
- 512
- 513 Gardner, W. S., McCarthy, M. J., An, S., Sobolev, D., Sell, K. S., & Brock, D. (2006). Nitrogen fixation
514 and dissimilatory nitrate reduction to ammonium (DNRA) support nitrogen dynamics in Texas
515 estuaries. *Limnology and Oceanography*, 51(1part2), 558–568. <https://doi.org/10.4319/lo.2006.51.1.part.2.0558>
- 516
- 517 Hagemann, M., Asce, S. M., Kim, D., & Park, M. H. (2016). Estimating Nutrient and Organic Carbon
518 Loads to Water-Supply Reservoir Using Semiparametric Models. *J. Environ. Eng.*, 9. [https://doi.org/10.1061/\(ASCE\)EE.1943-7870.0001077](https://doi.org/10.1061/(ASCE)EE.1943-7870.0001077)
- 519
- 520 Hayes, N. M., Haig, H. A., Simpson, G. L., & Leavitt, P. R. (2020). Effects of lake warming on the
521 seasonal risk of toxic cyanobacteria exposure. *Limnology and Oceanography Letters*, 5(6), 393–
522 402. <https://doi.org/10.1002/lol2.10164>
- 523
- 524 Hirsch, R. M., Moyer, D. L., & Archfield, S. A. (2010). Weighted Regressions on Time, Discharge,
525 and Season (WRTDS), with an application to Chesapeake Bay River inputs. *JAWRA Journal of
526 the American Water Resources Association*, 46(5), 857–880. <https://doi.org/10.1111/j.1752-1688.2010.00482.x>
- 527
- 528 Hornung, R. W., & Reed, L. D. (1990). Estimation of average concentration in the presence of nonde-
529 tectable values. *Applied Occupational and Environmental Hygiene*, 5(1), 46–51. <https://doi.org/10.1080/1047322X.1990.10389587>
- 530
- 531 Horowitz, A. J. (2003). An evaluation of sediment rating curves for estimating suspended sediment
532 concentrations for subsequent flux calculations. *Hydrological Processes*, 17(17), 3387–3409.
<https://doi.org/10.1002/hyp.1299>
- 533
- 534 Hou, L., Liu, M., Carini, S. A., & Gardner, W. S. (2012). Transformation and fate of nitrate near
535 the sediment–water interface of Copano Bay. *Continental Shelf Research*, 35, 86–94. <https://doi.org/10.1016/j.csr.2012.01.004>

- Jain, S., & Schramm, M. P. (2021). *Technical Support Document for One Total Maximum Daily Load for Indicator Bacteria in Lavaca River Above Tidal* (Technical Report No. AS-221). Texas Commission on Environmental Quality. Austin, Texas. <https://www.tceq.texas.gov/downloads/water-quality/tmdl/lavaca-river-above-tidal-rocky-creek-recreational-108/108-lavaca-river-addendum-tsd-2021-october-as-221.pdf>
- Kennicutt, M. C. (2017). Water Quality of the Gulf of Mexico. In C. H. Ward (Ed.), *Habitats and Biota of the Gulf of Mexico: Before the Deepwater Horizon Oil Spill* (pp. 55–164). Springer New York. https://doi.org/10.1007/978-1-4939-3447-8_2
- Kroon, F. J., Kuhnert, P. M., Henderson, B. L., Wilkinson, S. N., Kinsey-Henderson, A., Abbott, B., Brodie, J. E., & Turner, R. D. (2012). River loads of suspended solids, nitrogen, phosphorus and herbicides delivered to the Great Barrier Reef lagoon. *Marine Pollution Bulletin*, 65(4-9), 167–181. <https://doi.org/10.1016/j.marpolbul.2011.10.018>
- Kuhnert, P. M., Henderson, B. L., Lewis, S. E., Bainbridge, Z. T., Wilkinson, S. N., & Brodie, J. E. (2012). Quantifying total suspended sediment export from the Burdekin River catchment using the loads regression estimator tool. *Water Resources Research*, 48(4). <https://doi.org/10.1029/2011WR011080>
- Kulat, M. I., Mohtar, R. H., & Olivera, F. (2019). Holistic Water-Energy-Food Nexus for Guiding Water Resources Planning: Matagorda County, Texas Case. *Frontiers in Environmental Science*, 7, 3. <https://doi.org/10.3389/fenvs.2019.00003>
- Kumar, S., Godrej, A., Post, H., & Berger, K. (2019). The value of intensive sampling—a comparison of fluvial loads. *Water Resources Management*, 33(12), 4303–4318. <https://doi.org/10.1007/s11269-019-02369-7>
- Lloyd, C., Freer, J., Collins, A., Johnes, P., & Jones, J. (2014). Methods for detecting change in hydrochemical time series in response to targeted pollutant mitigation in river catchments. *Journal of Hydrology*, 514, 297–312. <https://doi.org/10.1016/j.jhydrol.2014.04.036>
- Marra, G., & Wood, S. N. (2012). Coverage properties of confidence intervals for Generalized Additive Model components: Coverage properties of GAM intervals. *Scandinavian Journal of Statistics*, 39(1), 53–74. <https://doi.org/10.1111/j.1467-9469.2011.00760.x>
- McDowell, R. W., Simpson, Z. P., Ausseil, A. G., Etheridge, Z., & Law, R. (2021). The implications of lag times between nitrate leaching losses and riverine loads for water quality policy. *Scientific Reports*, 11(1), 16450. <https://doi.org/10.1038/s41598-021-95302-1>
- Mishra, A. K., & Singh, V. P. (2010). Changes in extreme precipitation in Texas. *Journal of Geophysical Research*, 115(D14), D14106. <https://doi.org/10.1029/2009JD013398>
- Montagna, P. A., Cockett, P. M., Kurr, E. M., & Trungale, J. (2020). *Assessment of the Relationship Between Freshwater Inflow and Biological Indicators in Lavaca Bay* (tech. rep. No. Contract # 1800012268). Harte Research Institute, Texas A&M University-Corpus Christi. Corpus Christi, Texas.
- Mooney, R. F., & McClelland, J. W. (2012). Watershed export events and ecosystem responses in the Mission–Aransas National Estuarine Research Reserve, South Texas. *Estuaries and Coasts*, 35(6), 1468–1485. <https://doi.org/10.1007/s12237-012-9537-4>
- Moriasi, D. N., Gitau, M. W., Pai, N., & Daggupati, P. (2015). Hydrologic and Water Quality Models: Performance Measures and Evaluation Criteria. *Transactions of the ASABE*, 58(6), 1763–1785. <https://doi.org/10.13031/trans.58.10715>
- Murphy, R. R., Keisman, J., Harcum, J., Karrh, R. R., Lane, M., Perry, E. S., & Zhang, Q. (2022). Nutrient improvements in Chesapeake Bay: Direct effect of load reductions and implications for coastal management. *Environmental Science & Technology*, 56(1), 260–270. <https://doi.org/10.1021/acs.est.1c05388>
- Murphy, R. R., Perry, E., Harcum, J., & Keisman, J. (2019). A Generalized Additive Model approach to evaluating water quality: Chesapeake Bay case study. *Environmental Modelling & Software*, 118, 1–13. <https://doi.org/10.1016/j.envsoft.2019.03.027>
- Oelsner, G. P., & Stets, E. G. (2019). Recent trends in nutrient and sediment loading to coastal areas of the conterminous U.S.: Insights and global context. *Science of The Total Environment*, 654, 1225–1240. <https://doi.org/10.1016/j.scitotenv.2018.10.437>

- Omani, N., Srinivasan, R., & Lee, T. (2014). Estimation of sediment and nutrient loads to bays from gauged and ungauged watersheds. *Applied Engineering in Agriculture*, 869–887. <https://doi.org/10.13031/aea.30.10162>
- Örnólfsson, E. B., Lumsden, S., & Pinckney, J. L. (2004). Nutrient pulsing as a regulator of phytoplankton abundance and community composition in Galveston Bay, Texas. *Journal of Experimental Marine Biology and Ecology*, 303(2), 197–220. <https://doi.org/10.1016/j.jembe.2003.11.016>
- Palmer, T. A., & Montagna, P. A. (2015). Impacts of droughts and low flows on estuarine water quality and benthic fauna. *Hydrobiologia*, 753(1), 111–129. <https://doi.org/10.1007/s10750-015-2200-x>
- Park, Y., & Engel, B. (2014). Use of pollutant load regression models with various sampling frequencies for annual load estimation. *Water*, 6(6), 1685–1697. <https://doi.org/10.3390/w6061685>
- Paudel, B., Montagna, P. A., & Adams, L. (2019). The relationship between suspended solids and nutrients with variable hydrologic flow regimes. *Regional Studies in Marine Science*, 29, 100657. <https://doi.org/10.1016/j.rsma.2019.100657>
- Peierls, B. L., Hall, N. S., & Paerl, H. W. (2012). Non-monotonic responses of phytoplankton biomass accumulation to hydrologic variability: A comparison of two coastal plain North Carolina estuaries. *Estuaries and Coasts*, 35(6), 1376–1392. <https://doi.org/10.1007/s12237-012-9547-2>
- Rebich, R. A., Houston, N. A., Mize, S. V., Pearson, D. K., Ging, P. B., & Evan Hornig, C. (2011). Sources and delivery of nutrients to the northwestern Gulf of Mexico from streams in the south-central United States. *JAWRA Journal of the American Water Resources Association*, 47(5), 1061–1086. <https://doi.org/10.1111/j.1752-1688.2011.00583.x>
- Robson, B. J., & Dourdet, V. (2015). Prediction of sediment, particulate nutrient and dissolved nutrient concentrations in a dry tropical river to provide input to a mechanistic coastal water quality model. *Environmental Modelling & Software*, 63, 97–108. <https://doi.org/10.1016/j.envsoft.2014.08.009>
- Russell, M. J., Montagna, P. A., & Kalke, R. D. (2006). The effect of freshwater inflow on net ecosystem metabolism in Lavaca Bay, Texas. *Estuarine, Coastal and Shelf Science*, 68(1-2), 231–244. <https://doi.org/10.1016/j.ecss.2006.02.005>
- Schramm, M. (2023). *Adc: Calculate Antecedant Discharge Conditions*. Version 1.0.0. <https://CRAN.R-project.org/package=adc>
- Schramm, M., Berthold, A., Entwistle, C., & Peddicord, K. (2018, April). *Lavaca River Watershed Protection Plan* (Technical Report No. TR-507). Texas Water Resources Institute. College Station, Texas. <https://twri.tamu.edu/publications/technical-reports/2018-technical-reports/tr-507/>
- Schramm, M., Gitter, A., & Gregory, L. (2022). Total Maximum Daily Loads and *Escherichia coli* trends in Texas freshwater streams. *Journal of Contemporary Water Research & Education*, 176, 36–49. <https://doi.org/10.1111/j.1936-704X.2022.3374.x>
- Schwarz, G. E., Hoos, A. B., Alexander, R. B., & Smith, R. A. (2006). *The SPARROW Surface Water-Quality Model: Theory, Application and User Documentation* (6-B3). U.S. Geological Survey. <https://doi.org/10.3133/tm6B3>
- Shoda, M. E., Sprague, L. A., Murphy, J. C., & Riskin, M. L. (2019). Water-quality trends in U.S. rivers, 2002 to 2012: Relations to levels of concern. *Science of The Total Environment*, 650, 2314–2324. <https://doi.org/10.1016/j.scitotenv.2018.09.377>
- Snelder, T., McDowell, R., & Fraser, C. (2017). Estimation of catchment nutrient loads in New Zealand using monthly water quality monitoring data. *JAWRA Journal of the American Water Resources Association*, 53(1), 158–178. <https://doi.org/10.1111/1752-1688.12492>
- Stackpoole, S., Sabo, R., Falcone, J., & Sprague, L. (2021). Long-term Mississippi River trends expose shifts in the river load response to watershed nutrient balances between 1975 and 2017. *Water Resources Research*, 57(11). <https://doi.org/10.1029/2021WR030318>
- Stets, E., Lee, C., Lytle, D., & Schock, M. (2018). Increasing chloride in rivers of the conterminous U.S. and linkages to potential corrosivity and lead action level exceedances in drinking water. *Science of The Total Environment*, 613–614, 1498–1509. <https://doi.org/10.1016/j.scitotenv.2017.07.119>
- Tominack, S. A., & Wetz, M. S. (2022). Variability in phytoplankton biomass and community composition in Corpus Christi Bay, Texas. *Estuaries and Coasts*. <https://doi.org/10.1007/s12237-022-01137-y>
- Vecchia, A. V., Gilliom, R. J., Sullivan, D. J., Lorenz, D. L., & Martin, J. D. (2009). Trends in concentrations and use of agricultural herbicides for Corn Belt Rivers, 1996–2006. *Environmental Science & Technology*, 43(24), 9096–9102. <https://doi.org/10.1021/es902122j>

- Walker, L. M., Montagna, P. A., Hu, X., & Wetz, M. S. (2021). Timescales and magnitude of water quality change in three Texas estuaries induced by passage of Hurricane Harvey. *Estuaries and Coasts*, 44(4), 960–971. <https://doi.org/10.1007/s12237-020-00846-6>
- Wang, Y.-G., Kuhnert, P., & Henderson, B. (2011). Load estimation with uncertainties from opportunistic sampling data – A semiparametric approach. *Journal of Hydrology*, 396(1-2), 148–157. <https://doi.org/10.1016/j.jhydrol.2010.11.003>
- Wazniak, C. E., Hall, M. R., Carruthers, T. J. B., Sturgis, B., Dennison, W. C., & Orth, R. J. (2007). Linking water quality to living resources in a Mid-Atlantic lagoon system, USA. *Ecological Applications*, 17(sp5), S64–S78. <https://doi.org/10.1890/05-1554.1>
- Wetz, M. S., Cira, E. K., Sterba-Boatwright, B., Montagna, P. A., Palmer, T. A., & Hayes, K. C. (2017). Exceptionally high organic nitrogen concentrations in a semi-arid South Texas estuary susceptible to brown tide blooms. *Estuarine, Coastal and Shelf Science*, 188, 27–37. <https://doi.org/10.1016/j.ecss.2017.02.001>
- Wetz, M. S., Hayes, K. C., Fisher, K. V., Price, L., & Sterba-Boatwright, B. (2016). Water quality dynamics in an urbanizing subtropical estuary (Oso Bay, Texas). *Marine Pollution Bulletin*, 104(1-2), 44–53. <https://doi.org/10.1016/j.marpolbul.2016.02.013>
- Wetz, M. S., & Yoskowitz, D. W. (2013). An ‘extreme’ future for estuaries? Effects of extreme climatic events on estuarine water quality and ecology. *Marine Pollution Bulletin*, 69(1-2), 7–18. <https://doi.org/10.1016/j.marpolbul.2013.01.020>
- Wise, D. R., Anning, D. W., & Miller, O. W. (2019). *Spatially referenced models of streamflow and nitrogen, phosphorus, and suspended-sediment transport in streams of the southwestern United States* (Scientific Investigations Report No. 2019-5106). U.S. Geological Survey. Reston, Virginia.
- Wood, S. N. (2006). On confidence intervals for generalized additive models based on penalized regression splines. *Australian & New Zealand Journal of Statistics*, 48(4), 445–464. <https://doi.org/10.1111/j.1467-842X.2006.00450.x>
- Wood, S. N. (2011). Fast stable restricted maximum likelihood and marginal likelihood estimation of semiparametric generalized linear models: Estimation of Semiparametric Generalized Linear Models. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73(1), 3–36. <https://doi.org/10.1111/j.1467-9868.2010.00749.x>
- Zhang, Q., & Ball, W. P. (2017). Improving riverine constituent concentration and flux estimation by accounting for antecedent discharge conditions. *Journal of Hydrology*, 547, 387–402. <https://doi.org/10.1016/j.jhydrol.2016.12.052>