Some claims concerning a tetrahedron ABCP based on an acute triangle ABC

Scale and position the tetrahedron so that the coordinates of A, B and C are $(x_1, y_1, 0)$, $(x_2, y_2, 0)$, $(x_3, y_3, 0)$, respectively, with $x_j = \cos \phi_j$, $y_j = \sin \phi_j$ (j = 1,2,3), and $\phi_1 + \phi_2 + \phi_3 = 0$.

Define the following:

$$\angle A = \angle \text{CAB} , \ \angle B = \angle \text{ABC} , \ \angle C = \angle \text{BCA} , \ \alpha = \angle \text{BPC} , \ \beta = \angle \text{CPA} , \ \gamma = \angle \text{APB} ,$$

$$C_0 = \cos \alpha \cos \beta \cos \gamma , C_1 = \cos^2 \alpha , C_2 = \cos^2 \beta , C_3 = \cos^2 \gamma ,$$

$$L = 2 \left[(y_1 + y_2 + y_3)(1 - C_0) + (1 - x_1)y_1(C_1 - 1) + (1 - x_2)y_2(C_2 - 1) + (1 - x_3)y_3(C_3 - 1) \right] ,$$

$$R = 2 \left[(1 + x_1 + x_2 + x_3)(1 - C_0) + (1 + x_1)x_1(C_1 - 1) + (1 + x_2)x_2(C_2 - 1) + (1 + x_3)y_3(C_3 - 1) \right] ,$$

$$H = 1 + 2 C_0 - C_1 - C_2 - C_3 , \qquad E = L^2 + (R + H)^2 ,$$

$$D = E^2 + 18 \ EH^2 + 8 (R + H) H \left[(R + H)^2 - 3L^2 \right] - 27 H^4 .$$

The following (proven) constraints hold:

1.
$$H > 0$$
 (which is equivalent to $\alpha + \beta + \gamma < 2\pi$, $\alpha < \beta + \gamma$, $\beta < \alpha + \gamma$, $\gamma < \alpha + \beta$);

2.
$$\angle A + \beta + \gamma < 2\pi$$
, $\alpha + \angle B + \gamma < 2\pi$, $\alpha + \beta + \angle C < 2\pi$;

3.
$$\beta + \gamma - \alpha < 2 (\angle B + \angle C)$$
, $\gamma + \alpha - \beta < 2 (\angle C + \angle A)$, $\alpha + \beta - \gamma < 2 (\angle A + \angle B)$;

4.
$$\alpha < \angle A \rightarrow [\beta < \max \{ \angle B, \angle C + \alpha \} \land \gamma < \max \{ \angle C, \angle B + \alpha \}]$$
,
 $\beta < \angle B \rightarrow [\gamma < \max \{ \angle C, \angle A + \beta \} \land \alpha < \max \{ \angle A, \angle C + \beta \}]$,
 $\gamma < \angle C \rightarrow [\alpha < \max \{ \angle A, \angle B + \gamma \} \land \beta < \max \{ \angle B, \angle A + \gamma \}]$;

5.
$$\alpha < \angle A \rightarrow \cos \angle C \cos \beta + \cos \angle B \cos \gamma > 0$$
,
 $\beta < \angle B \rightarrow \cos \angle A \cos \gamma + \cos \angle C \cos \alpha > 0$,
 $\gamma < \angle C \rightarrow \cos \angle B \cos \alpha + \cos \angle A \cos \beta > 0$;

6.
$$(D > 0 \land \alpha < \angle A) \rightarrow (\beta < \angle B \lor \gamma < \angle C),$$

 $(D > 0 \land \beta < \angle B) \rightarrow (\gamma < \angle C \lor \alpha < \angle A),$
 $(D > 0 \land \gamma < \angle C) \rightarrow (\alpha < \angle A \lor \beta < \angle B).$

Also, experimental evidence strongly suggests that this collection of constraints is also sufficient for a suitable tetrahedron ABCP to exist for a given acute triangle ABC. At present though, this sufficiency claim is just a conjecture.