

5.

Throughout, G is a connected, reductive algebraic group over $k = \bar{\mathbf{F}}_q$ with a Frobenius map $F : G \rightarrow G$. We fix an F -stable Borel pair (B, T) and write $U = [B, B]$. We fix $\delta \geq 1$ so that F^δ acts trivially on W , and a section $w \mapsto \dot{w} : W \rightarrow N_{G^{F^\delta}}(T^{F^\delta})$. With these choices, $X_w \subseteq G/B$ and $\tilde{X}_w \subseteq G/U$ are F^δ -stable for all $w \in W$.

5.1.

Recall that in our running example where $G = \mathrm{SL}_2$ and F is standard, we can write $W = \{e, s\}$ with $e = \mathrm{id}$, and take $\delta = 1$. Last time, we computed the graded $\bar{\mathbf{Q}}_\ell[F]$ -modules formed by the compactly-supported ℓ -adic cohomologies of X_e and X_s :

$$H_c^*(X_e) \simeq \bar{\mathbf{Q}}_\ell^{\oplus(q+1)}, \quad H_c^*(X_s) \simeq \bar{\mathbf{Q}}_\ell^{\oplus q}[-1] \oplus \bar{\mathbf{Q}}_\ell[-2](-1).$$

Above $[-m]$ means “shift up by degree m ” and $(-m)$ means “twist the Frobenius action by a factor of q^m ”.

One more property of ℓ -adic cohomology that I could have added to the list from last time:

- (10) $H^0(X)$ is the vector space of $\bar{\mathbf{Q}}_\ell$ -valued functions on the set of connected components of X .

This gives another way to identify $H_c^*(X_e) = H_c^0(X_e)$, and by Poincaré duality, $H_c^2(X_s) \simeq H^0(X_s)^\vee[-2](-1)$. But it does more: It enables us to identify the G^F -actions on these vector spaces. It remains for us to identify the G^F -action on $H_c^1(X_s)$.

5.2.

As mentioned last time, it is easier in general to work with the virtual character $R_{w,\theta}$ than with the individual representations $H_c^i(\tilde{X}_w)[\theta]$. For any k -scheme of finite type X and automorphism $g : X \rightarrow X$, the *Lefschetz number* of g on $H_c^*(X)$ is defined to be

$$\mathcal{L}_X(g) = \sum_i (-1)^i \mathrm{tr}(g \mid H_c^i(X)).$$

The Lefschetz fixed-point formula tells us that if f is a Frobenius map, then $\mathcal{L}_X(f) = |X^f|$. At the same time,

$$\begin{aligned} \mathcal{L}_{X_w} &= R_{w,1}, \\ \mathcal{L}_{\tilde{X}_w} &= \sum_{\theta} R_{w,\theta} \end{aligned}$$

as functions on G^F .

The next result that we present, combining Exercise 4.7.4 and Theorem 4.4.12 in Geck, is a bridge between these two uses of Lefschetz number. Recall that $g : X \rightarrow X$ commutes with a Frobenius map $F : X \rightarrow X$ corresponding to some \mathbf{F}_q -rational structure $X = X_1 \otimes k$ if and only if g descends to X_1 , meaning $g = g_1 \otimes \text{id}$. Note that since X is of finite type, g is cut out by finitely many polynomials in finitely many variables. Thus, g is always defined over some finite subfield of k ; in other words, given g , we can always find some Frobenius that commutes with g .

Theorem 5.1. *Suppose that X is a smooth k -variety with Frobenius f , and $g : X \rightarrow X$ is an automorphism of finite order that commutes with f . Then:*

- (1) gf^m is a Frobenius map on X for all $m \geq 1$.
- (2) The formal series

$$\mathcal{L}_X(g, t) := - \sum_{m \geq 1} |X^{gf^m}| t^m$$

satisfies $\mathcal{L}_X(g) = \lim_{t \rightarrow \infty} \mathcal{L}_X(g, t)$.

Proof of (2) from (1). Since f and g commute, we can triangularize them simultaneously. Suppose that $(\lambda_{i,j})_j$, resp. $(\mu_{i,j})_j$, is the list of eigenvalues of f , resp. g , on $H_c^i(X)$. Since gf^m is a Frobenius map, the Lefschetz formula gives

$$|X^{gf^m}| = \sum_i (-1)^i \sum_j \mu_{i,j} \lambda_{i,j}^m,$$

from which

$$-\mathcal{L}_X(g, t) = \sum_{m,i,j} (-1)^i \mu_{i,j} \lambda_{i,j}^m t^m = \sum_{i,j} (-1)^i \mu_{i,j} \frac{\mu_{i,j} t}{1 - \mu_{i,j} t}.$$

Now observe that $\frac{\mu_{i,j} t}{1 - \mu_{i,j} t} \rightarrow -1$ as $t \rightarrow \infty$. □

Remark 5.2. The *Weil zeta series* of X with respect to f is defined by

$$Z_X(t) = \exp \left(\sum_{m \geq 1} |X^{f^m}| \frac{t^m}{m} \right),$$

where \exp is a formal exponential. We see that

$$\mathcal{L}_X(\text{id}, t) = -t \frac{d}{dt} \log Z_X(t).$$

In this sense, $\mathcal{L}(t, |g, X)$ is a mild generalization of the zeta series.

Corollary 5.3. *Keeping the hypotheses of Theorem 5.1, suppose that X is the union of disjoint subvarieties X' and X'' that are f -stable and g -stable. Then*

$$\mathcal{L}_X = \mathcal{L}_{X'} + \mathcal{L}_{X''}$$

as functions of g .

Previously, we sketched the reason why X_w and \tilde{X}_w form smooth varieties. If $g \in G^F$, then the action of g on G/B and G/U commutes with that of F , and hence, its action on X_w and \tilde{X}_w commutes with that of F^δ . So we can apply Theorem 5.1 and its corollary to the case where $X = X_w, \tilde{X}_w$, or some unions of these, and $f = F^\delta$ and $g \in G^F$.

Returning to the setup with $G = \mathrm{SL}_2$ and F standard, we deduce that

$$\mathcal{L}_{G/B} = \mathcal{L}_{X_e} + \mathcal{L}_{X_s} = R_{e,1} + R_{s,1}.$$

We also know the cohomology of G/B , since it is \mathbf{P}^1 :

$$H_c^*(G/B) \simeq H^*(G/B) \simeq \bar{\mathbf{Q}}_\ell \oplus \bar{\mathbf{Q}}_\ell[-2](-1),$$

Since $H^0(G/B)$ carries the trivial representation of G^F , the same is true of its Poincaré dual $H^2(G/B)$. Therefore $\mathcal{L}_{G/B}(g) = 2$ for all g .

From Mackey, we saw that the G^F -equivariant endomorphisms of $\mathbf{R}_{e,1} = H_c^*(X_e) = H_c^0(X_e)$ form a 2-dimensional algebra, which forces $\mathbf{R}_{e,1}$ to be a sum of two irreducible representations of G^F . But $\mathbf{R}_{e,1}$ is also the space of functions on X_e , which contains the trivial representation. So we must have

$$R_{e,1} = 1 + \mathrm{St} \quad \text{for some irreducible character } \mathrm{St}.$$

This is the Steinberg character mentioned previously. Finally,

$$R_{s,1} = \mathcal{L}_{G/B} - R_{e,1} = 2 - (1 + \mathrm{St}) = 1 - \mathrm{St}.$$

Since $\mathbf{R}_{s,1} = H_c^1(X_s) \oplus H_c^2(X_s)$, and $H_c^2(X_s)$ also carries the trivial character, we deduce that $H_c^1(X_s)$ carries the Steinberg character.

5.3.

Before we can describe $\mathbf{R}_{s,\theta} = H_c^*(X_s)[\theta]$ and $R_{s,\theta}$ for other θ , we should describe T^{sF} more explicitly. Taking T to be the diagonal torus given by

$$T(k) = \{t_a \mid a \in k^\times\}, \quad \text{where } t_a = \begin{pmatrix} a & \\ & a^{-1} \end{pmatrix},$$

we see that $s \cdot t_a = t_{a^{-1}}$. Therefore,

$$T^{sF} = \{t_a \in T \mid a^q = a^{-1}\} = \{t_a \in T \mid a^{q+1} = 1\}.$$

In particular, T^{sF} is cyclic of order $q + 1$.

Note that the condition $a^{q+1} = 1$ forces $a \in \mathbf{F}_{q^2}^\times$. Moreover, $a \in \mathbf{F}_q^\times$ only happens for $a = \pm 1$. These computations show that in general, the embedding of T into G does *not* restrict to an embedding of T^{sF} into $G^F = \mathrm{SL}_2(\mathbf{F}_q)$. Nonetheless:

Proposition 5.4. *Let G be any connected, smooth reductive algebraic group over k with Frobenius F . For any F -stable maximal torus $T \subseteq G$ and element $w \in W = N_G(T)/T$, we can find some $g \in G(k)$ such that $S := gTg^{-1}$ is F -stable and $S^F = gT^{wF}g^{-1}$. In particular, we get an embedding*

$$T^{wF} \xrightarrow{\sim} S^F \rightarrow G^F.$$

Proof. Lift w to an element $\dot{w} \in N_G(T)(k) \subseteq G(k)$. By Lang's theorem, we can find $g \in G(k)$ such that $\dot{w} = g^{-1}F(g)$. We see that

$$F(gTg^{-1}) = F(g)TF(g)^{-1} = g\dot{w}T\dot{w}^{-1}g^{-1} = gTg^{-1},$$

proving that gTg^{-1} is F -stable. Moreover, for all $t \in T(k)$, we see that $F(gtg^{-1}) = gtg^{-1} \iff \dot{w}F(t)\dot{w}^{-1} = t$. Thus $(gTg^{-1})^F = gT^{wF}g^{-1}$. \square

5.4.

To conclude our discussion of fixed-point formulas, we present two main results by Deligne–Lusztig, and explain their application to the discrete series of $\mathrm{SL}_2(\mathbf{F}_q)$. Geck omits their proofs in his Section 4.5.

The first result is Deligne–Lusztig Theorem 3.2. To motivate it, recall that any invertible matrix g over a field has a *Jordan decomposition* $g = g_s g_u = g_u g_s$, where g_s is diagonalizable (or *semisimple*) and g_u is unipotent. If the field characteristic is $p > 0$ and the (multiplicative) order of g is finite, then the order of g_s is coprime to p , while the order of g_u is a power of p .

Theorem 5.5 (Deligne–Lusztig). *Suppose that X is a smooth affine k -variety with Frobenius f , and $g : X \rightarrow X$ is an automorphism of finite order that commutes with f . Suppose that $g = g_s g_u = g_u g_s$, where $g_s : X \rightarrow X$, resp. $g_u : X \rightarrow X$, has order coprime to p , resp. a power of p . Then*

$$\mathcal{L}_X(g) = \mathcal{L}_{X^{g_s}}(g_u).$$

In the SL_2 example, this theorem implies that for any $t \in T^{sF}$, we have $\mathcal{L}_{\tilde{X}_s}(t) = \mathcal{L}_{\tilde{X}_s}(1)$. But T^{sF} acts freely on \tilde{X}_s , so the right-hand side vanishes whenever $t \neq 1$! By character theory, we deduce that as a representation of T^{sF} , the vector space $H_c^*(\tilde{X}_s)$ is a \oplus -power of the regular representation of T^{sF} . Since

T^{sF} is abelian, every character occurs in the latter with the same multiplicity. Therefore

$$\dim R_{s,\theta} = \dim R_{s,1} = 1 - q \quad \text{for all } \theta.$$

To actually determine how these characters decompose beyond the $\theta = 1$ case, we need more firepower.

The following result is Deligne–Lusztig Theorem 6.8. It is a geometric generalization of the Mackey-type formula we saw earlier. For the transporter scheme $N_G(S, S') = \{g \in G \mid S' = gSg^{-1}\}$ that we use below, see Milne Chapter 1, Section i.

Theorem 5.6 (Deligne–Lusztig). *Suppose that w', S' also satisfy the hypotheses on w, S in the setup of Proposition 5.4. Fix a character θ of S^F , resp. θ' of $(S')^F$, and identify it with a character of T^{wF} , resp. $T^{w'F}$. Then*

$$(R_{w,\theta}, R_{w',\theta'})_{G^F} = \frac{|N_G((S, \theta), (S', \theta'))^F|}{|S^F|},$$

where $N_G((S, \theta), (S', \theta'))$ is the subvariety of elements $g \in N_G(S, S')$ such that $\theta' = {}^g\theta$.

Corollary 5.7. *In the setup above,*

$$(R_{w,\theta}, R_{w,\theta})_{G^F} = |\{w \in W^F \mid {}^w\theta = \theta\}|.$$

The theorem statement hints that some features of the theory should really be stated directly in terms of S^F , rather than T^{wF} . We will return to this later.

In the SL_2 example, we have

$$(R_{s,\theta}, R_{s,\theta})_{G^F} = \begin{cases} 2 & \theta^2 = 1, \\ 1 & \text{else.} \end{cases}$$

In particular, $-R_{s,\theta}$ is an actual, irreducible representation of G^F whenever θ is a character of T^{sF} such that $\theta^2 \neq 1$. For q odd, there are $q - 1$ choices of such θ , which form $\frac{1}{2}(q - 1)$ conjugate pairs under s . Each pair contributes one new irreducible. The remaining two irreducibles of G^F are the summands of $\mathbf{R}_{s,\theta}$ for θ the order-2 character of T^{sF} . Taken together, these are all the *discrete series representations* of $\mathrm{SL}_2(\mathbf{F}_q)$.