

# ANNULAR WEBS AND A CONJECTURE OF HAIMAN

MINH-TÂM QUANG TRINH

**ABSTRACT.** Haiman conjectured that when traces corresponding to monomial symmetric functions are evaluated on the Hecke-algebra elements denoted  $C'_w$  by Kazhdan–Lusztig, the resulting polynomials have nonnegative coefficients. We show that recent work on annular webs implies this for permutations  $w$  that are 321-hexagon-avoiding. As a byproduct, we observe a precise analogy that matches the two bases in Kazhdan–Lusztig’s work with elementary and homogeneous symmetric functions.

## 1. INTRODUCTION

1.1. Let  $H_n(x)$  be the Iwahori–Hecke algebra of the symmetric group  $S_n$  over  $\mathbf{Z}[x^{\pm 1}]$ . As a quotient of the group algebra of a braid group, it has a standard basis  $\{\sigma_w\}_{w \in S_n}$ , consisting of the images of the positive permutation braids.

Kazhdan–Lusztig introduced two new bases for  $H_n(x)$  with remarkable properties [KL79]. Taking our  $x$  to be their  $q^{1/2}$ , we will focus on the basis that they denote by  $\{C'_w\}_w$ , but write  $b_w$  in place of  $C'_w$  for simplicity. When the elements  $b_w$  are expanded in the standard basis, the coefficients are Laurent polynomials in  $x$  with nonnegative integer coefficients. Up to rescaling, these are the celebrated Kazhdan–Lusztig polynomials for  $S_n$ . Their positivity can be proved through a geometric interpretation of  $H_n(x)$  in terms of sheaves on flag varieties.

The representation theory of  $S_n$  deforms to that of  $H_n(x)$ . In particular, each character  $\chi : S_n \rightarrow \bar{\mathbf{Q}}$  defines a  $\mathbf{Z}[x^{\pm 1}]$ -linear function  $\chi_x : H_n(x) \rightarrow \bar{\mathbf{Q}}(x)$  that still enjoys the trace property  $\chi(\alpha\beta) = \chi(\beta\alpha)$ . At the same time, the *irreducible* characters of  $S_n$  are indexed by integer partitions of  $n$ . Let  $\chi^\lambda$  be the irreducible character indexed by  $\lambda \vdash n$ . A geometric argument, similar to that used in the positivity of the Kazhdan–Lusztig polynomials, proves that  $\chi_x^\lambda(b_w) \in \mathbf{Z}_{\geq 0}[x^{\pm 1}]$  for all  $w$  and  $\lambda$ .

Haiman found evidence for a stronger positivity statement. Recall that for all  $\lambda, \mu$ , the Kostka number  $K_{\lambda, \mu}$  counts semistandard Young tableaux of shape  $\lambda$  and weight  $\mu$ . The Kostka numbers can be assembled into a unitriangular matrix of nonnegative integers. In particular, this matrix has an inverse with integer entries, so there are functions  $\phi_x^\mu : H_n(x) \rightarrow \mathbf{Z}[x^{\pm 1}]$  uniquely defined by requiring

$$(1.1) \quad \chi_x^\lambda = \sum_{\mu} K_{\lambda, \mu} \phi_x^\mu \quad \text{for all } \lambda \vdash n.$$

What follows is the main part of Conjecture 2.1 in [Hai93].

**Conjecture 1.1** (Haiman).  $\phi_x^\mu(b_w) \in \mathbf{Z}_{\geq 0}[x^{\pm 1}]$  for all  $w \in S_n$  and  $\mu \vdash n$ .

Abreu–Nigro observe that Conjecture 1.1 would imply several conjectures about the indifference graphs of Hessenberg functions in algebraic combinatorics: notably, the Stanley–Stembridge conjecture on the  $e$ -positivity of their chromatic symmetric functions, and Shareshian–Wachs’s generalization of this conjecture to chromatic quasi-symmetric functions [AN24].

1.2. This note will show how recent work of Queffelec–Rose and Gorsky–Wedrich on the diagrammatics of  $H_n(x)$  solves some cases of Conjecture 1.1.

For  $1 \leq i \leq n-1$ , let  $b_i = b_{s_i}$ , where  $s_i \in S_n$  is the transposition that swaps  $i$  and  $i+1$ . The main theorem is:

**Theorem 1.2.**  $\phi_x^\mu(b_{i_1} \cdots b_{i_\ell}) \in \mathbf{Z}_{\geq 0}[x^{\pm 1}]$  for any sequence of indices  $i_1, \dots, i_\ell$  that range between 1 and  $n-1$  inclusive, and any  $\mu \vdash n$ .

In what follows, suppose that  $w \in S_n$  is given by  $w = [w_1 w_2 \cdots w_n]$ , meaning it sends  $i$  to  $w_i$  for  $1 \leq i \leq n$ . Fix  $m \leq n$  and  $v = [v_1 v_2 \cdots v_m] \in S_m$ . We say that  $w$  is  $v_1 \cdots v_m$ -*avoiding* if and only if the sequence  $(w_1, \dots, w_n)$  does *not* contain a subsequence of size  $m$  whose elements have the same relative order as  $(v_1, \dots, v_m)$ . More formally, this means we cannot find indices  $1 \leq p_1 < \cdots < p_m \leq n$  such that  $w_{p_i} < w_{p_j}$  whenever  $i < j$  and  $v_i < v_j$ .

We write  $S_n^{v_1 \cdots v_m} \subseteq S_n$  for the set of  $v_1 \cdots v_m$ -avoiding elements. Following Billey–Warrington, we say that  $w$  is *321-hexagon-avoiding* if and only if

$$w \in S_n^{321} \cap S_n^{46718235} \cap S_n^{46781235} \cap S_n^{56718234} \cap S_n^{56781234}.$$

In [BW01], Billey–Warrington prove that the following conditions are equivalent:

- (1)  $w$  is 321-hexagon-avoiding.
- (2)  $b_w = b_{i_1} \cdots b_{i_\ell}$  whenever  $w = s_{i_1} \cdots s_{i_\ell}$  and  $\ell$  is the minimal length among such expressions.

Via this result, Theorem 1.2 implies:

**Corollary 1.3.** *Conjecture 1.1 holds when  $w$  is 321-hexagon-avoiding.*

1.3. The key observation is that Remark 4.21 of [GW23], a refinement of the annular web evaluation algorithm of [QR18], provides a counterpart to Theorem 1.2 (in fact, a slightly stronger statement) in the setting of MOY webs. The passage from Hecke-algebra traces to web diagrammatics is best explained by assembling the cocenters of all the Hecke algebras into a direct sum that we identify with Macdonald’s ring of symmetric functions  $\Lambda(x)$  over  $\mathbf{Z}[x^{\pm 1}]$ , after extending scalars. This defines a universal trace

$$\mathrm{tr} : \bigoplus_n H_n(x) \rightarrow \Lambda(x).$$

There is a natural candidate for the diagrammatic counterpart to  $\mathrm{tr}$ : the map  $\mathrm{ann}$  that sends a rectangular web to its annular closure.

For any  $\beta \in H_n(x)$ , the value of  $\phi_x^\mu(\beta)$  is just the  $\mu$ -th coefficient when we expand  $\mathrm{tr}(\beta)$  in the basis of complete homogeneous symmetric functions  $\{h_\mu\}_\mu$ : a fact already noted in [AN24]. Ultimately, we relate Theorem 1.2 to [GW23, Rem.

4.21] through a commutative diagram that relates  $\text{tr}$  to  $\text{ann}$ , and assigns simple webs to the  $b_i$  and  $h_\mu$ .

I do not know of any earlier statement of this commutative diagram in the literature, despite closely related results in the work of Aiston, Lukac, Morton, *et al.*: See [MM08] and the references there.

In fact, there is an analogous but inequivalent commutative diagram, where  $\{b_w\}_w$  is replaced by Kazhdan–Lusztig’s *other* basis for the Hecke algebra, and  $\{h_\mu\}_\mu$  is replaced by the basis of elementary symmetric functions  $\{e_\mu\}_\mu$ . The dichotomy between the two diagrams seems related to the dichotomy between the framed and unframed Khovanov–Rozansky functors in [GW23], though not exactly. We expand on this at the end of the note.

**1.4. Acknowledgments.** I thank Elijah Bodish, Mikhail Khovanov, and Paul Wedrich for helpful discussions.

## 2. HECKE ALGEBRAS AND SYMMETRIC FUNCTIONS

2.1. Recall that  $S_n$  forms a Coxeter group, in which the transpositions  $s_i$  for  $1 \leq i \leq n-1$  (see §1.2) form a system of simple reflections. With respect to this Coxeter presentation, let  $\ell_w$  denote the Bruhat length of  $w \in S_n$ , and let  $<$  be the Bruhat order on  $S_n$  [GP00, Ch. 1].

2.2. Formally, we take the *Iwahori–Hecke algebra* of  $S_n$  to be the  $\mathbf{Z}[x^{\pm 1}]$ -algebra  $H_n(x)$  spanned as a free module by elements  $\sigma_w$  for  $w \in S_n$ , modulo the following relations, where we set  $\sigma_i := \sigma_{s_i}$ :

$$(2.1) \quad \sigma_w \sigma_i = \begin{cases} \sigma_{ws_i} & ws_i > w, \\ \sigma_{ws_i} + (x - x^{-1})\sigma_w & ws_i < w. \end{cases}$$

There is an additive involution  $D : H_n(x) \rightarrow H_n(x)$  that sends  $x \mapsto x^{-1}$  and  $\sigma_w \mapsto \sigma_{w^{-1}}^{-1}$  for all  $w \in S_n$ .

2.3. Let  $\mathbf{K} = \mathbf{Q}(x)$ . It turns out that  $\mathbf{K} \otimes_{\mathbf{Z}[x^{\pm 1}]} H_n(x)$  is split as a  $\mathbf{K}$ -algebra. At the same time, there is an isomorphism of rings  $H_n(x)|_{x \rightarrow 1} \simeq \mathbf{Z}S_n$ . So by Tits deformation [GP00, Ch. 7], the semisimplicity of  $\mathbf{Q}S_n$  implies the semisimplicity of  $\mathbf{K} \otimes_{\mathbf{Z}[x^{\pm 1}]} H_n(x)$ , and moreover, there is a bijection between isomorphism classes of simple  $\mathbf{Q}S_n$ -modules and those of simple  $(\mathbf{K} \otimes_{\mathbf{Z}[x^{\pm 1}]} H_n(x))$ -modules.

This induces the assignment from characters  $\chi : S_n \rightarrow \bar{\mathbf{Q}}$  to  $\mathbf{Z}[x^{\pm 1}]$ -linear trace functions  $\chi_x : H_n(x) \rightarrow \bar{\mathbf{K}}$  described in the introduction. Explicitly,  $\chi_x(\beta)$  is the trace of  $\beta$  on the  $(\mathbf{K} \otimes_{\mathbf{Z}[x^{\pm 1}]} H_n(x))$ -module that corresponds to the  $\mathbf{Q}S_n$ -module with character  $\chi$ .

2.4. Kazhdan–Lusztig proved that for all  $w \in S_n$ , there is a unique  $D$ -invariant element  $b_w \in H_n(x)$  such that

$$b_w = \sum_{y \leq w} x^{\ell_y - \ell_w} P_{y,w}(x^2) \sigma_y$$

for some  $P_{y,w}(q) \in \mathbf{Z}[q]$  satisfying

$$(2.2) \quad \begin{aligned} P_{w,w}(q) &= 1, \\ \deg P_{y,w}(q) &\leq \frac{1}{2}(\ell_w - \ell_y - 1) \quad \text{for all } w, y \in S_n \text{ with } y \leq w. \end{aligned}$$

Let  $j$  be the additive involution that sends  $x \mapsto x^{-1}$  and  $\sigma_w \mapsto (-1)^{\ell_w} \sigma_w$ . Let  $c_w = j(b_w)$ . Then  $c_w$  is the unique  $D$ -invariant element of  $H_n(x)$  such that

$$c_w = \sum_{y \leq w} (-1)^{\ell_y} x^{\ell_w - \ell_y} P_{y,w}(x^{-2}) \sigma_y$$

for some  $P_{y,w}(q) \in \mathbf{Z}[q]$  satisfying (2.2). They turn out to be the same polynomials as before.

The sets  $\{b_w\}_{w \in S_n}$  and  $\{c_w\}_{w \in S_n}$  form bases for  $H_n(x)$  as a free  $\mathbf{Z}[x^{\pm 1}]$ -module, known as the two *Kazhdan–Lusztig bases* or *canonical bases*. The polynomials  $P_{y,w}(q)$  are the *Kazhdan–Lusztig polynomials* for  $S_n$ . Note that in [KL79],  $b_w$  and  $c_w$  are respectively denoted  $C'_w$  and  $-C_w$ .

Henceforth, we will write  $b_i, c_i$  in place of  $b_{s_i}, c_{s_i}$ . We can check that

$$(2.3) \quad \begin{aligned} b_i &= x^{-1} + \sigma_i &= x &+ \sigma_i^{-1}, \\ c_i &= x - \sigma_i &= x^{-1} - \sigma_i^{-1}. \end{aligned}$$

Just as  $\{\sigma_i\}_i$  generates  $H_n(x)$  as a  $\mathbf{Z}[x^{\pm 1}]$ -algebra, so do  $\{b_i\}_i$  and  $\{c_i\}_i$ .

2.5. Let  $\Lambda$  be the graded ring of symmetric functions over  $\mathbf{Z}$  in (countably) infinitely many variables. For background on  $\Lambda$ , we refer to [Mac15, Ch. I]. In this note, we will need the following elements of  $\Lambda$  indexed by integer partitions  $\lambda$ :

the Schur functions	$s_\lambda$ ,
the monomial symmetric functions	$m_\lambda = m_{\lambda_1} m_{\lambda_2} \dots$ ,
the complete homogeneous symmetric functions	$h_\lambda = h_{\lambda_1} h_{\lambda_2} \dots$ ,
the elementary symmetric functions	$e_\lambda = e_{\lambda_1} e_{\lambda_2} \dots$

Let  $\Lambda_n$  be the degree- $n$  component of  $\Lambda$ . The Schur functions  $s_\lambda$  with  $\lambda \vdash n$  form a basis for  $\Lambda$  as a free  $\mathbf{Z}$ -module; analogous statements hold with  $m_\lambda$  or  $h_\lambda$  or  $e_\lambda$  in place of  $s_\lambda$ .

Recall the Kostka numbers  $K_{\lambda,\mu} \in \mathbf{Z}$  from the introduction. As explained in [Mac15, §I.6], they relate the elements  $s_\lambda, m_\mu, h_\mu$  via the identities

$$(2.4) \quad \begin{aligned} s_\lambda &= \sum_{\mu} K_{\lambda,\mu} m_\mu, \\ h_\mu &= \sum_{\lambda} K_{\lambda,\mu} s_\lambda. \end{aligned}$$

The first identity shows that Haiman's character  $\phi_x^\mu$  is to the monomial symmetric function  $\mu_\mu$  as the irreducible character  $\chi_x^\lambda$  is to the Schur function  $s_\lambda$ .

Note that  $K_{\lambda,\lambda} = 1$  for all  $\lambda$ , and that  $K_{\lambda,\mu} = 0$  whenever  $\mu > \lambda$  in the dominance order on partitions. This makes precise the unitriangularity mentioned earlier.

2.6. Let  $\Lambda(x) = \mathbf{Z}[x^{\pm 1}] \otimes \Lambda$  and  $\Lambda_n(x) = \mathbf{Z}[x^{\pm 1}] \otimes \Lambda_n$  for all  $n$ . The map  $\mathrm{tr}$  mentioned in §1.3 is the sum of the  $\mathbf{Z}[x^{\pm 1}]$ -linear maps

$$\mathrm{tr}_n : H_n(x) \rightarrow \Lambda_n(x) \quad \text{defined by } \mathrm{tr}_n(\beta) = \sum_{\lambda \vdash n} \chi_x^\lambda(\beta) s_\lambda.$$

By construction,  $\mathrm{tr}_n(\alpha\beta) = \mathrm{tr}_n(\beta\alpha)$  for all  $\alpha, \beta$ . So the universal property of the cocenter of  $H_n(x)$  defines a  $\mathbf{Z}[x^{\pm 1}]$ -linear map from the cocenter into  $\Lambda(x)$ , which turns out to be an isomorphism of  $\mathbf{Z}[x^{\pm 1}]$ -modules.

Let  $\langle -, - \rangle : \Lambda(x) \times \Lambda(x) \rightarrow \mathbf{Z}[x^{\pm 1}]$  be the *Hall pairing*: the  $\mathbf{Z}[x^{\pm 1}]$ -linear pairing under which the Schur functions  $s_\lambda$  form an orthonormal basis. It lets us write

$$\chi_x^\lambda(\beta) = \langle \mathrm{tr}_n(\beta), s_\lambda \rangle \quad \text{for all } \lambda \vdash n.$$

By (1.1) and (2.4), we deduce:

$$(2.5) \quad \mathrm{tr}_n(\beta) = \sum_{\mu \vdash n} \phi_x^\mu(\beta) h_\mu.$$

Altogether, Theorem 1.2 is claiming that for any sequence of indices  $i_1, \dots, i_\ell$  that range between 1 and  $n-1$  inclusive, the expansion of  $\mathrm{tr}_n(b_{i_1} \cdots b_{i_\ell})$  in the complete homogeneous basis of  $\Lambda_n(x)$  will have coefficients in  $\mathbf{Z}_{\geq 0}[x^{\pm 1}]$ .

### 3. WEBS

3.1. We refer to [GW23, §2] for background on the Murakami–Ohtsuki–Yamada (MOY) web calculus. Note that their  $q$  is our  $x$ .

3.2. Let  $H_n^{\mathrm{mo}}(x)$  be the free  $\mathbf{Z}[x^{\pm 1}]$ -module generated by strictly upward-oriented web diagrams in a rectangle, connecting  $n$  inputs with label 1 at the bottom to  $n$  outputs with label 1 at the top, modulo the relations of the MOY bracket. It forms a  $\mathbf{Z}[x^{\pm 1}]$ -algebra under concatenation of diagrams. Murakami–Ohtsuki–Yamada showed that this algebra is isomorphic to  $H_n(x)$  [MOY98].

For  $1 \leq i \leq n-1$ , let  $\mathrm{can}_i \in H_n^{\mathrm{mo}}(x)$  denote the  $i$ th merge-split web. The notation  $\mathrm{can}$  is intended to suggest the adjective *canonical*. The precise statement proved in [MOY98], up to sign and up to use of Schur–Weyl duality, is that there is an isomorphism of  $\mathbf{Z}[x^{\pm 1}]$ -algebras

$$\Theta_b : H_n(x) \rightarrow H_n^{\mathrm{mo}}(x) \quad \text{defined by } \Theta_b(b_i) = \mathrm{can}_i.$$

To explain how this isomorphism appears in [GW23]: Recall that we can identify  $H_n(x)$  with the skein algebra of a rectangle with  $n$  inputs and  $n$  outputs, by sending  $\sigma_i$  to the  $i$ th simple twist. Decategorified, rectangular analogues of formulas (16) and (17) in [GW23] define two isomorphisms from this skein algebra to  $H_n^{\mathrm{mo}}(x)$ . Our map  $\Theta_b$  corresponds to their *framed* map (16).

3.3. Let  $\mathcal{C}^{\mathrm{mo}}(x)$  be the free  $\mathbf{Z}[x^{\pm 1}]$ -module generated by positively-oriented web diagrams in an annulus. It forms a commutative  $\mathbf{Z}[x^{\pm 1}]$ -algebra under nesting of diagrams.

For any  $n$  and  $\mu \vdash n$ , let  $o^\mu \in \mathcal{C}^{\mathrm{mo}}(x)$  be the diagram consisting of concentric essential circles with labels  $\mu_1, \mu_2, \dots$ . Note that by the commutativity of  $\mathcal{C}^{\mathrm{mo}}(x)$ ,

the order of these circles does not matter. The *annular web evaluation algorithm* of Queffelec–Rose [QR18, Lem. 5.2] shows that the set  $\{o_\mu\}_\mu$  forms a basis for  $\mathcal{C}^{\text{moy}}(x)$  as a free  $\mathbf{Z}[x^{\pm 1}]$ -module. The definition of the MOY bracket then implies, directly, that  $\mathcal{C}^{\text{moy}}(x)$  is freely generated *as an algebra* by the elements  $o_n := o_{(n)}$  corresponding to single, labeled essential circles. By placing  $o_n$  in degree  $n$ , we can further endow  $\mathcal{C}^{\text{moy}}(x)$  with the structure of a *graded* algebra.

At the same time, display (2.4), *resp.* (2.8), in [Mac15] implies that  $\Lambda(x)$  is freely generated as an algebra by the set  $\{e_n\}_n$ , *resp.* the set  $\{h_n\}_n$ . Thus, there are isomorphisms of graded  $\mathbf{Z}[x^{\pm 1}]$ -algebras

$$\Xi_h, \Xi_e : \Lambda(x) \rightarrow \mathcal{C}^{\text{moy}}(x) \quad \text{defined by } \Xi_e(e_\mu) = o_\mu \text{ and } \Xi_h(h_\mu) = o_\mu.$$

They differ precisely by the  $\mathbf{Z}[x^{\pm 1}]$ -algebra involution of  $\Lambda(x)$  that swaps  $h_\mu$  and  $e_\mu$ .

Prior to the introduction of webs, an analogous isomorphism for the skein algebra of the annulus was first established by Turaev [Tur88].

3.4. As in the literature on skein algebras, there is a  $\mathbf{Z}[x^{\pm 1}]$ -linear map

$$\text{ann} : \bigoplus_n H_n^{\text{moy}}(x) \rightarrow \mathcal{C}^{\text{moy}}(x)$$

called *annular closure*. It is defined graphically, by embedding a rectangle into an annulus as a sector, so that the upward orientation in the rectangle becomes the positive orientation in the annulus, then wrapping the  $n$  outputs of the rectangle around the annulus, without crossing, back to the  $n$  inputs.

Queffelec–Rose’s annular web evaluation algorithm originally treated  $\mathcal{C}^{\text{moy}}(x)$  as the triangulated Grothendieck group of the bounded homotopy category of a graded, linear category of annular foams between positively-oriented webs. Gorsky–Wedrich observed that it could be refined, by instead treating  $\mathcal{C}^{\text{moy}}(x)$  as the additive Grothendieck group of the *Karoubi* or *idempotent completion* of this foam category [GW23, Rem. 4.21]. The refinement shows:

**Theorem 3.1** (Positive Annular Web Evaluation). *The expansion of any single positively-oriented annular web in the basis  $\{o_\mu\}_\mu$  for  $\mathcal{C}^{\text{moy}}(x)$  will have coefficients in  $\mathbf{Z}_{\geq 0}[x^{\pm 1}]$ . In particular, this applies to  $\text{ann}(\text{can}_{i_1} \cdots \text{can}_{i_\ell})$  for any indices  $i_1, \dots, i_\ell$ .*

3.5. Together, (2.5) and Theorem 3.1 reduce Theorem 1.2 to the following result:

**Theorem 3.2.** *The following diagram commutes:*

$$\begin{array}{ccc} H_n(x) & \xrightarrow{\text{tr}} & \Lambda(x) \\ \Theta_b \downarrow & & \downarrow \Xi_h \\ H_n^{\text{moy}}(x) & \xrightarrow{\text{ann}} & \mathcal{C}^{\text{moy}}(x) \end{array}$$

*That is: For any  $\beta \in H_n(x)$  and  $\mu \vdash n$ , the value of  $\phi_x^\mu(\beta)$  is the coefficient of  $o_\mu$  when we expand  $\text{ann}(\Theta_b(\beta))$  in the basis  $\{o_\mu\}_\mu$ .*

*Proof.* **TODO**

□

3.6. Theorem 3.1 suggests a categorification of Conjecture 1.1.

Let  $\mathcal{C}$  be the category denoted  $\text{Kar}(\mathbf{AFoam}^+)$  in [GW23]: a Karoubi-complete, graded, linear category of foams between positively-oriented webs. Let  $\mathbf{H}_n$  be the analogous category where we replace the annulus by a rectangle with  $n$  inputs and  $n$  outputs. By work of Mackaay–Vaz [MV10],  $\mathbf{H}_n$  is a diagrammatic presentation of the category of Soergel bimodules for  $S_n$ , and hence, categorifies  $H_n(x)$ . Let  $\mathbf{B}_w$  be the indecomposable object of  $\mathbf{H}_n$  indexed by  $w \in S_n$ , so that the isomorphism from the Grothendieck group to  $H_n(x)$  sends  $[\mathbf{B}_w]$  to  $b_w$ . Let  $\mathbf{O}_\mu$  be the object of  $\mathcal{C}_n$  underlying the annular web  $o_\mu$ .

**Conjecture 3.3.** *For all  $w \in S_n$ , the annular closure of  $\mathbf{B}_w$  is isomorphic in  $\mathcal{C}$  to a direct sum of objects of the form  $\mathbf{O}_\mu$ .*

#### 4. THE OTHER COMMUTATIVE DIAGRAM

4.1. We claim that there is an isomorphism of  $\mathbf{Z}[x^{\pm 1}]$ -algebras

$$\Theta_c : H_n(x) \rightarrow H_n^{\text{moy}}(x) \quad \text{defined by } \Theta_c(c_i) = \text{can}_i,$$

analogous to but distinct from  $\Theta_b$ . Indeed, the relations (2.1) show that if either  $\alpha_i = b_i$  for all  $i$ , or  $\alpha_i = c_i$  for all  $i$ , then  $H_n(x)$  is generated by its subset  $\{\alpha_i\}_i$  modulo the relations

$$\begin{cases} \alpha_i \alpha_{i+1} \alpha_i - \alpha_i = \alpha_{i+1} \alpha_i \alpha_{i+1} - \alpha_{i+1}, \\ \alpha_i \alpha_j = \alpha_j \alpha_i & \text{for } |i - j| > 1, \\ \alpha_i^2 = (x + x^{-1}) \alpha_i. \end{cases}$$

We see that  $\Theta_c$  is the precomposition of  $\Theta_b$  with the  $\mathbf{Z}[x^{\pm 1}]$ -algebra involution of  $H_n(x)$  that swaps  $b_i$  and  $c_i$ . Note that this involution is not the map  $j$  from §2.4, since  $j$  is not  $\mathbf{Z}[x^{\pm 1}]$ -linear.

4.2. Moreover, we claim the following analogue of Theorem 3.2.

**Theorem 4.1.** *The following diagram commutes:*

$$\begin{array}{ccc} H_n(x) & \xrightarrow{\text{tr}} & \Lambda(x) \\ \Theta_c \downarrow & & \downarrow \Xi_e \\ H_n^{\text{moy}}(x) & \xrightarrow{\text{ann}} & \mathcal{C}^{\text{moy}}(x) \end{array}$$

*Proof.* First, recall that the involution of  $\Lambda(x)$  that swaps  $h_\mu$  and  $e_\mu$  also swaps  $s_\lambda$  and  $s_{\lambda^t}$ , where  $\lambda^t$  is the transpose of  $\lambda$  [Mac15, (3.8)]. So the map

$$\text{tr}_n^t : H_n(x) \rightarrow \Lambda_n(x) \quad \text{defined by } \text{tr}_n^t(\beta) = \sum_{\lambda \vdash n} \chi_x^\lambda(\beta) s_{\lambda^t}$$

satisfies  $\Xi_e \circ \text{tr}_n = \Xi_h \circ \text{tr}_n^t$ . Next, let  $\eta$  be the  $\mathbf{Z}[x^{\pm 1}]$ -algebra involution of  $H_n(x)$  that swaps  $b_i$  and  $c_i$ . Since  $\Theta_c = \Theta_b \circ \eta$ , it remains to show that  $\text{tr}^t = \text{tr} \circ \eta$ , then invoke Theorem 3.2.

Observe that  $\text{tr}_n^t(\beta) = \sum_{\lambda \vdash n} \chi_x^{\lambda^t}(\beta) s_\lambda$ . So we must show that  $\chi_x^{\lambda^t} = \chi_x^\lambda \circ \eta$  for all  $\lambda$ . Since  $\chi^{\lambda^t} = \varepsilon \chi^\lambda$ , where  $\varepsilon$  is the sign character of  $S_n$ , we are done by the following lemma.  $\square$

**Lemma 4.2.**  $(\varepsilon\chi)_x = \chi_x \circ \eta$  for all characters  $\chi$ .

*Proof.* By Proposition 9.4.1 of [GP00],

$$(\varepsilon\chi^\lambda)_x(\sigma_w) = (-1)^{\ell_w} \chi_x^\lambda(\sigma_w)|_{x \rightarrow x^{-1}}.$$

Using (2.3), we deduce that

$$\chi_x^{\lambda^t}(b_{i_1} \cdots b_{i_\ell}) = \chi_x^\lambda(c_{i_1} \cdots c_{i_\ell}) = \chi_x^\lambda(\eta(b_{i_1} \cdots b_{i_\ell}))$$

for any sequence of indices  $i_1, \dots, i_\ell$ . But the  $b_i$  generate  $H_n(x)$  as an algebra, so every element of  $H_n(x)$  is a linear combination of elements of the form  $b_{i_1} \cdots b_{i_\ell}$ .  $\square$

*Remark 4.3.* The sign character  $\varepsilon$ , the algebra involution  $\eta$ , and Lemma 4.2 all generalize beyond  $S_n$  to any finite Coxeter group.

#### 4.3. TODO

#### REFERENCES

- [AN24] Alex Corrêa Abreu and Antonio Nigro. An update on Haiman’s conjectures. *Forum Math. Sigma*, 12:Paper No. e86, 15, 2024.
- [BW01] Sara C. Billey and Gregory S. Warrington. Kazhdan-Lusztig polynomials for 321-hexagon-avoiding permutations. *J. Algebraic Combin.*, 13(2):111–136, 2001.
- [GP00] M. Geck and G. Pfeiffer. *Characters of finite Coxeter groups and Iwahori–Hecke algebras*, volume 21 of *London Mathematical Society Monographs. New Series*. The Clarendon Press, Oxford University Press, New York, 2000.
- [GW23] Eugene Gorsky and Paul Wedrich. Evaluations of annular Khovanov–Rozansky homology. *Math. Z.*, 303(1):Paper No. 25, 57, 2023.
- [Hai93] Mark Haiman. Hecke algebra characters and immanant conjectures. *J. Amer. Math. Soc.*, 6(3):569–595, 1993.
- [KL79] David Kazhdan and George Lusztig. Representations of Coxeter groups and Hecke algebras. *Invent. Math.*, 53(2):165–184, 1979.
- [Mac15] I. G. Macdonald. *Symmetric functions and Hall polynomials*. Oxford Classic Texts in the Physical Sciences. The Clarendon Press, Oxford University Press, New York, second edition, 2015. With contribution by A. V. Zelevinsky and a foreword by Richard Stanley.
- [MM08] H. R. Morton and P. M. G. Manchón. Geometrical relations and plethysms in the Homfly skein of the annulus. *J. Lond. Math. Soc. (2)*, 78(2):305–328, 2008.
- [MOY98] Hitoshi Murakami, Tomotada Ohtsuki, and Shuji Yamada. Homfly polynomial via an invariant of colored plane graphs. *Enseign. Math. (2)*, 44(3-4):325–360, 1998.
- [MV10] Marco Mackaay and Pedro Vaz. The diagrammatic Soergel category and  $\mathrm{sl}(N)$ -foams, for  $N \geq 4$ . *Int. J. Math. Math. Sci.*, pages Art. ID 468968, 20, 2010.
- [QR18] Hoel Queffelec and David E. V. Rose. Sutured annular Khovanov–Rozansky homology. *Trans. Amer. Math. Soc.*, 370(2):1285–1319, 2018.
- [Tur88] V. G. Turaev. The Conway and Kauffman modules of a solid torus. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)*, 167:79–89, 190, 1988.

DEPARTMENT OF MATHEMATICS, YALE UNIVERSITY, NEW HAVEN, CT 06520  
*Email address:* minh-tam.trinh@yale.edu