

naive idea behind Seifert–van Kampen:

if X is covered by open U_i 's [draw]
then can compute $\pi_1(X)$ from $\pi_1(U_i)$'s

but only works when all intersections of U_i 's are
path-connected

Thm (Seifert–van Kampen)

suppose $A, U_1, U_2 \subset X$ are open and $x \in A$

s.t. $X = U_1 \cup U_2$,
 $A = U_1 \cap U_2$
 A, U_1, U_2 are all path-connected

with inclusion maps

$$\begin{array}{ccccc} & i_1 & & U_1 & & j_1 \\ A & & & & & X \\ & i_2 & & U_2 & & j_2 \end{array}$$

then:

1) $j_{1,*}, j_{2,*}$ induce a surjective hom

$$\pi_1(U_1, x) * \pi_1(U_2, x) \rightarrow \pi_1(X, x)$$

2) via this hom, $\pi_1(X, x)$ is the largest quotient
of $\pi_1(U_1, x) * \pi_1(U_2, x)$ in which

$$i_{1,*}([y]) \sim i_{2,*}([y]) \text{ for all } [y] \in \pi_1(A, x)$$

Ex X the figure-eight,
 U_1, U_2 open thickenings of the S^1 's,
 x the intersection point
 [draw]

since $\pi_1(A, x)$ is trivial:

$$\begin{aligned}\pi_1(X, x) &\simeq \pi_1(U_1, x) * \pi_1(U_2, x) \\ &\simeq \pi_1(S^1, x) * \pi_1(S^1, x) \\ &\simeq \mathbb{Z} * \mathbb{Z}\end{aligned}$$

Ex $X = S^2$,
 U_1, U_2 open thickenings of opposed
 hemispheres,
 [draw]

here, $\pi_1(A, x) \simeq \pi_1(S^1, x) \simeq \mathbb{Z}$
but $\pi_1(U_1, x), \pi_1(U_2, x)$ trivial
so $\pi_1(X, x)$ also trivial

Rem Seifert–van Kampen alone
 cannot compute $\pi_1(S^1)$

(Munkres §53–54) let $o = (1, 0)$ in S^1

recall: $\omega_n(s) = (\cos(2\pi ns), \sin(2\pi ns))$

Thm $\Phi : \mathbb{Z}$ to $\pi_1(S^1, o)$ def by $\Phi(n) = [\omega_n]$
 is an isomorphism

Pf Step 1. Φ is a homomorphism
 Step 2. Φ is bijective

Step 1 must show that $\Phi(m + n) = \Phi(m) + \Phi(n)$,
meaning $[\omega_{m+n}] = [\omega_m * \omega_n]$

key idea: let $p : \mathbb{R} \rightarrow S^1$ be

$$p(x) = (\cos(2\pi x), \sin(2\pi x))$$

Lem for any $a, b \in \mathbb{Z}$ s.t. $b - a = n$, we have

$$\omega_n = p \circ \omega_{\{a, b\}}$$

where $\omega_{\{a, b\}} : [0, 1] \rightarrow \mathbb{R}$ is defined by

$$\omega_{\{a, b\}}(s) = (1 - s)a + sb$$

[draw]

Lem for any $a, b, c \in \mathbb{Z}$, we have

$$[\omega_{\{a, b\}} * \omega_{\{b, c\}}] = [\omega_{\{a, c\}}]$$

[draw]

therefore,

$$\begin{aligned} & [(p \circ \omega_{\{a, b\}}) * (p \circ \omega_{\{b, c\}})] \\ &= [p \circ (\omega_{\{a, b\}} * \omega_{\{b, c\}})] \text{ [Munkres 327]} \\ &= [p \circ \omega_{\{a, c\}}] \text{ [PS5, \#1]} \end{aligned}$$

giving $[\omega_m * \omega_n] = [\omega_{m+n}]$

before Step 2, we introduce the notion of covering map inspired by $p : \mathbb{R} \rightarrow S^1$