



# Affine Springer Fibers and Level-Rank Duality

---

Minh-Tâm Quang Trinh

Yale University



- 1 Springer Theory
- 2 Deligne–Lusztig Theory
- 3 Level-Rank Duality

Mainly about joint work with Ting Xue:

[arXiv:2311.17106](https://arxiv.org/abs/2311.17106)

See also the extended abstract on my website, which we have submitted to FPSAC '25.

- 1 Springer Theory
- 2 Deligne–Lusztig Theory
- 3 Level-Rank Duality

Mainly about joint work with Ting Xue:

[arXiv:2311.17106](https://arxiv.org/abs/2311.17106)

See also the extended abstract on my website, which we have submitted to FPSAC '25.

## 1 Springer Theory      Work over $\mathbf{C}$ .

$\mathbf{G}$     connected reductive group

$\mathbf{A}$     maximal torus

$W$     Weyl group

The *rational Cherednik algebra*  $D_c^{\text{rat}}$  is a deformation of  $\mathbf{C}W \ltimes \mathcal{D}(\mathbf{a})$  depending on a parameter  $c \in \mathbf{C}$ .

$$\begin{array}{ll}
 D_c^{\text{rat}} & \mathbf{U}\mathbf{g} \\
 \mathbf{C}[\mathbf{a}] \otimes \mathbf{C}W \otimes \mathbf{C}[\mathbf{a}^*] & \mathbf{U}\mathbf{n}_- \otimes \mathbf{U}\mathbf{a} \otimes \mathbf{U}\mathbf{n}_+ \\
 \Delta_c(\chi) & \Delta(\lambda) \\
 L_c(\chi) & L(\lambda)
 \end{array}$$

For  $c$  rational,  $D_c^{\text{rat}}$  can fail to be semisimple. This is the most interesting case.

# 1 Springer Theory      Work over $\mathbf{C}$ .

$\mathbf{G}$     connected reductive group

$\mathbf{A}$     maximal torus

$W$     Weyl group

The *rational Cherednik algebra*  $D_c^{\text{rat}}$  is a deformation of  $\mathbf{C}W \ltimes \mathcal{D}(\mathbf{a})$  depending on a parameter  $c \in \mathbf{C}$ .

$$\begin{array}{ll} D_c^{\text{rat}} & \mathbf{U}\mathfrak{g} \\ \mathbf{C}[\mathbf{a}] \otimes \mathbf{C}W \otimes \mathbf{C}[\mathbf{a}^*] & \mathbf{U}\mathfrak{n}_- \otimes \mathbf{U}\mathbf{a} \otimes \mathbf{U}\mathfrak{n}_+ \\ \Delta_c(\chi) & \Delta(\lambda) \\ L_c(\chi) & L(\lambda) \end{array}$$

For  $c$  rational,  $D_c^{\text{rat}}$  can fail to be semisimple. This is the most interesting case.

For  $c$  rational and *positive*,  $D_c^{\text{rat}}$ -modules from the geometry of *affine Springer fibers*.

$\mathbf{B}$                     Borel containing  $\mathbf{A}$   
 $\mathbf{I} \subseteq \mathbf{G}[[z]]$     Iwahori lifting  $\mathbf{B} \subseteq \mathbf{G}$

The affine Springer fiber over  $\gamma \in \mathfrak{g}((z))$  is

$$\mathcal{FL}_\gamma = \{g\mathbf{I} \in \mathbf{G}((z))/\mathbf{I} \mid \gamma \in \text{Lie}(g\mathbf{I}g^{-1})\}.$$

Note that  $\mathbf{G}((z))/\mathbf{I}$  is infinite-dimensional.

We say that  $\gamma$  is *regular semisimple* iff  $\mathbf{G}((z))_\gamma^\circ$  is a maximal torus.

Here  $\mathcal{FL}_\gamma$  is finite-dimensional!

But it varies wildly over  $\mathfrak{g}((z))^{\text{rs}} \subseteq \mathfrak{g}((z))$ .

# 1 Springer Theory      Work over $\mathbf{C}$ .

$\mathbf{G}$     connected reductive group  
 $\mathbf{A}$     maximal torus  
 $W$     Weyl group

The *rational Cherednik algebra*  $D_c^{\text{rat}}$  is a deformation of  $\mathbf{C}W \ltimes \mathcal{D}(\mathbf{a})$  depending on a parameter  $c \in \mathbf{C}$ .

$D_c^{\text{rat}}$	$\mathbf{U}\mathfrak{g}$
$\mathbf{C}[\mathbf{a}] \otimes \mathbf{C}W \otimes \mathbf{C}[\mathbf{a}^*]$	$\mathbf{U}\mathfrak{n}_- \otimes \mathbf{U}\mathbf{a} \otimes \mathbf{U}\mathfrak{n}_+$
$\Delta_c(\chi)$	$\Delta(\lambda)$
$L_c(\chi)$	$L(\lambda)$

For  $c$  rational,  $D_c^{\text{rat}}$  can fail to be semisimple. This is the most interesting case.

For  $c$  rational and *positive*,  $D_c^{\text{rat}}$ -modules from the geometry of *affine Springer fibers*.

**B**                      Borel containing **A**

**I**  $\subseteq$  **G**[[ $z$ ]]    Iwahori lifting **B**  $\subseteq$  **G**

The affine Springer fiber over  $\gamma \in \mathfrak{g}((z))$  is

$$\mathcal{FL}_\gamma = \{g\mathbf{I} \in \mathbf{G}((z))/\mathbf{I} \mid \gamma \in \text{Lie}(g\mathbf{I}g^{-1})\}.$$

Note that  $\mathbf{G}((z))/\mathbf{I}$  is infinite-dimensional.

We say that  $\gamma$  is *regular semisimple* iff  $\mathbf{G}((z))_\gamma^\circ$  is a maximal torus.

Here  $\mathcal{FL}_\gamma$  is finite-dimensional!

But it varies wildly over  $\mathfrak{g}((z))^{\text{rs}} \subseteq \mathfrak{g}((z))$ .

For  $c$  rational and *positive*,  $D_c^{\text{rat}}$ -modules from the geometry of *affine Springer fibers*.

$$\begin{array}{ll} \mathbf{B} & \text{Borel containing } \mathbf{A} \\ \mathbf{I} \subseteq \mathbf{G}[[z]] & \text{Iwahori lifting } \mathbf{B} \subseteq \mathbf{G} \end{array}$$

The affine Springer fiber over  $\gamma \in \mathfrak{g}((z))$  is

$$\mathcal{FL}_\gamma = \{g\mathbf{I} \in \mathbf{G}((z))/\mathbf{I} \mid \gamma \in \text{Lie}(g\mathbf{I}g^{-1})\}.$$

Note that  $\mathbf{G}((z))/\mathbf{I}$  is infinite-dimensional.

We say that  $\gamma$  is *regular semisimple* iff  $\mathbf{G}((z))_\gamma^\circ$  is a maximal torus.

Here  $\mathcal{FL}_\gamma$  is finite-dimensional!

But it varies wildly over  $\mathfrak{g}((z))^{\text{rs}} \subseteq \mathfrak{g}((z))$ .

Fix rational  $c = \frac{d}{m} > 0$  in lowest terms.

Let  $\mathbf{C}^\times \curvearrowright \mathbf{G}((z))$  according to

$$\boxed{c \cdot g(z) = \text{Ad}(c^{d\rho^\vee})g(c^m z).} \quad (\rho^\vee = \sum_\alpha \omega_\alpha^\vee)$$

(Oblomkov–Yun)  $\mathcal{FL}_\gamma$  is locally constant over

$$\mathfrak{g}_{d/m}^{\text{rs}} = \{\gamma \in \mathfrak{g}((z))^{\text{rs}} \mid c \cdot \gamma = c^d \gamma\},$$

and  $\mathbf{C}^\times \curvearrowright \mathcal{FL}_\gamma$  for such  $\gamma$ .

We say that  $\gamma$  is *homogeneous of slope*  $\frac{d}{m}$ .

**Example** Take  $\mathbf{G} = \mathbf{SL}_2$  and  $\mathbf{B}$  upper-triangular.

Then  $\begin{pmatrix} 1 & \\ & -1 \end{pmatrix}, \begin{pmatrix} & 1 \\ z & \end{pmatrix}, \begin{pmatrix} z & \\ & -z \end{pmatrix}$  have slopes  $0, \frac{1}{2}, 1$ .



Fix rational  $c = \frac{d}{m} > 0$  in lowest terms.

Let  $\mathbf{C}^\times \curvearrowright \mathbf{G}((z))$  according to

$$\boxed{c \cdot g(z) = \mathrm{Ad}(c^{d\rho^\vee})g(c^m z).} \quad (\rho^\vee = \sum_\alpha \omega_\alpha^\vee)$$

(Oblomkov–Yun)  $\mathcal{F}l_\gamma$  is locally constant over

$$\mathbf{g}_{d/m}^{\mathrm{rs}} = \{\gamma \in \mathbf{g}((z))^{\mathrm{rs}} \mid c \cdot \gamma = c^d \gamma\},$$

and  $\mathbf{C}^\times \curvearrowright \mathcal{F}l_\gamma$  for such  $\gamma$ .

We say that  $\gamma$  is *homogeneous of slope  $\frac{d}{m}$* .

**Example** Take  $\mathbf{G} = \mathbf{SL}_2$  and  $\mathbf{B}$  upper-triangular.

Then  $\begin{pmatrix} 1 & \\ & -1 \end{pmatrix}, \begin{pmatrix} & 1 \\ z & \end{pmatrix}, \begin{pmatrix} z & \\ & -z \end{pmatrix}$  have slopes  $0, \frac{1}{2}, 1$ .

(Oblomkov–Yun) Take  $\mathbf{G}$  simply-connected, simple.

For  $\gamma \in \mathfrak{g}_{d/m}^{\text{rs}}$  such that  $\mathcal{F}l_\gamma$  is proper:

- A perverse filtration  $\mathbf{P}_{\leq *}$  on  $H_{\mathbf{C}^\times}^*(\mathcal{F}l_\gamma)$ .

It arises from a Ngô-type global model.

- An action of  $D_{d/m}^{\text{rat}}$  on

$$\mathcal{E}_\gamma := \text{gr}_*^{\mathbf{P}} H_{\mathbf{C}^\times}^*(\mathcal{F}l_\gamma)^{\pi_0(\mathbf{G}_0, \gamma)}|_{\epsilon \rightarrow 1},$$

where  $\mathbf{G}_0 = (\mathbf{G}((z))^{\mathbf{C}^\times})^\circ$  and  $\epsilon \in H_{\mathbf{C}^\times}^2(\text{point})$ .

As a module,  $\mathcal{E}_\gamma$  contains  $L_{d/m}(\chi_{\text{triv}})$ .

Equality holds when  $m$  is the Coxeter number.

Fix rational  $c = \frac{d}{m} > 0$  in lowest terms.

Let  $\mathbf{C}^\times \curvearrowright \mathbf{G}((z))$  according to

$$\boxed{c \cdot g(z) = \text{Ad}(c^{d\rho^\vee})g(c^m z).} \quad (\rho^\vee = \sum_\alpha \omega_\alpha^\vee)$$

(Oblomkov–Yun)  $\mathcal{F}l_\gamma$  is locally constant over

$$\mathfrak{g}_{d/m}^{\text{rs}} = \{\gamma \in \mathfrak{g}((z))^{\text{rs}} \mid c \cdot \gamma = c^d \gamma\},$$

and  $\mathbf{C}^\times \curvearrowright \mathcal{F}l_\gamma$  for such  $\gamma$ .

We say that  $\gamma$  is *homogeneous of slope  $\frac{d}{m}$* .

Example Take  $\mathbf{G} = \mathbf{SL}_2$  and  $\mathbf{B}$  upper-triangular.

Then  $\begin{pmatrix} 1 & \\ & -1 \end{pmatrix}, \begin{pmatrix} & 1 \\ z & \end{pmatrix}, \begin{pmatrix} & z \\ & -z \end{pmatrix}$  have slopes  $0, \frac{1}{2}, 1$ .

(Oblomkov–Yun) Take  $\mathbf{G}$  simply-connected, simple.

For  $\gamma \in \mathfrak{g}_{d/m}^{\text{rs}}$  such that  $\mathcal{F}l_\gamma$  is proper:

- A perverse filtration  $\mathbf{P}_{\leq *}$  on  $H_{\mathbf{C}^\times}^*(\mathcal{F}l_\gamma)$ .

It arises from a Ngô-type global model.

- An action of  $D_{d/m}^{\text{rat}}$  on

$$\mathcal{E}_\gamma := \text{gr}_*^{\mathbf{P}} H_{\mathbf{C}^\times}^*(\mathcal{F}l_\gamma)^{\pi_0(\mathbf{G}_0, \gamma)}|_{\epsilon \rightarrow 1},$$

where  $\mathbf{G}_0 = (\mathbf{G}((z))^{\mathbf{C}^\times})^\circ$  and  $\epsilon \in H_{\mathbf{C}^\times}^2(\text{point})$ .

As a module,  $\mathcal{E}_\gamma$  contains  $L_{d/m}(\chi_{\text{triv}})$ .

Equality holds when  $m$  is the Coxeter number.

(Oblomkov–Yun) Take  $\mathbf{G}$  simply-connected, simple.

For  $\gamma \in \mathfrak{g}_{d/m}^{\text{rs}}$  such that  $\mathcal{F}l_\gamma$  is proper:

- A perverse filtration  $\mathbf{P}_{\leq *}$  on  $H_{\mathbf{G}^\times}^*(\mathcal{F}l_\gamma)$ .

It arises from a Ngô-type global model.

- An action of  $D_{d/m}^{\text{rat}}$  on

$$\mathcal{E}_\gamma := \text{gr}_*^{\mathbf{P}} H_{\mathbf{G}^\times}^*(\mathcal{F}l_\gamma)^{\pi_0(\mathbf{G}_0, \gamma)}|_{\epsilon \rightarrow 1},$$

where  $\mathbf{G}_0 = (\mathbf{G}((z))^{\mathbf{C}^\times})^\circ$  and  $\epsilon \in H_{\mathbf{G}^\times}^2(\text{point})$ .

As a module,  $\mathcal{E}_\gamma$  contains  $L_{d/m}(\chi_{\text{triv}})$ .

Equality holds when  $m$  is the Coxeter number.

**Problem** Give a formula for  $D_{d/m}^{\text{rat}} \curvearrowright \mathcal{E}_\gamma$  in general.

In practice, too hard. Replace with

$$\mathcal{E}_\gamma := \sum_i (-1)^i \text{gr}_*^{\mathbf{P}} H_{\mathbf{G}^\times}^i(\mathcal{F}l_\gamma)^{\pi_0(\mathbf{G}_0, \gamma)}|_{\epsilon \rightarrow 1}.$$

**Idea**  $D_{d/m}^{\text{rat}}$  commutes with monodromy of  $\mathcal{E}_\gamma$  over

$$\mathfrak{c}_{d/m}^{\text{rs}} \subseteq \mathfrak{g}_{d/m}^{\text{rs}},$$

a Kostant-type transverse slice to  $\mathbf{G}_0 \curvearrowright \mathfrak{g}_{d/m}^{\text{rs}}$ .

The monodromy seems to factor through an algebra from *Deligne–Lusztig theory*.

**Problem** Give a formula for  $D_{d/m}^{\text{rat}} \curvearrowright \mathcal{E}_\gamma$  in general.  
 In practice, too hard. Replace with

$$E_\gamma := \sum_i (-1)^i \text{gr}_*^{\text{P}} H_{\mathbf{C}^\times}^i(\mathcal{F}l_\gamma)^{\pi_0(\mathbf{G}_0, \gamma)}|_{\epsilon \rightarrow 1}.$$

**Idea**  $D_{d/m}^{\text{rat}}$  commutes with monodromy of  $\mathcal{E}_\gamma$  over

$$\mathbf{c}_{d/m}^{\text{rs}} \subseteq \mathfrak{g}_{d/m}^{\text{rs}},$$

a Kostant-type transverse slice to  $\mathbf{G}_0 \curvearrowright \mathfrak{g}_{d/m}^{\text{rs}}$ .

The monodromy seems to factor through an algebra from *Deligne–Lusztig theory*.

Deligne–Lusztig studied groups over *finite fields*. But up to Tate twist,

$$\mathrm{Gal}(\bar{\mathbf{F}}_q|\mathbf{F}_q) \simeq \hat{\mathbf{Z}} \simeq \mathrm{Gal}(\overline{\mathbf{C}((z))}|\mathbf{C}((z))).$$

Forms of  $\mathbf{G}$  are classified by Dynkin automorphisms in the same way over  $\mathbf{F}_q$  as over  $\mathbf{C}((z))$ .

Much of Oblomkov–Yun’s setup generalizes from  $\mathbf{G}$  to any of its forms  $\mathbf{G}_{\mathbf{C}((z))}$ .

The tori  $\mathbf{A}, \mathbf{G}_\gamma$  generalize to forms  $\mathbf{A}_{\mathbf{C}((z))}, \mathbf{G}_{\mathbf{C}((z)), \gamma}$ .

These have corresponding forms  $\mathbf{A}_{\mathbf{F}_q}, \mathbf{T}_{\mathbf{F}_q}$ .

Problem Give a formula for  $D_{d/m}^{\mathrm{rat}} \curvearrowright \mathcal{E}_\gamma$  in general. In practice, too hard. Replace with

$$E_\gamma := \sum_i (-1)^i \mathrm{gr}_*^{\mathrm{P}} \mathrm{H}_{\mathbf{C}^\times}^i(\mathcal{F}l_\gamma)^{\pi_0(\mathbf{G}_{0,\gamma})}|_{\epsilon \rightarrow 1}.$$

Idea  $D_{d/m}^{\mathrm{rat}}$  commutes with monodromy of  $\mathcal{E}_\gamma$  over

$$\mathbf{c}_{d/m}^{\mathrm{rs}} \subseteq \mathfrak{g}_{d/m}^{\mathrm{rs}},$$

a Kostant-type transverse slice to  $\mathbf{G}_0 \curvearrowright \mathfrak{g}_{d/m}^{\mathrm{rs}}$ .

The monodromy seems to factor through an algebra from *Deligne–Lusztig theory*.

Deligne–Lusztig studied groups over *finite fields*. But up to Tate twist,

$$\mathrm{Gal}(\bar{\mathbf{F}}_q|\mathbf{F}_q) \simeq \hat{\mathbf{Z}} \simeq \mathrm{Gal}(\overline{\mathbf{C}((z))}|\mathbf{C}((z))).$$

Forms of  $\mathbf{G}$  are classified by Dynkin automorphisms in the same way over  $\mathbf{F}_q$  as over  $\mathbf{C}((z))$ .

Much of Oblomkov–Yun’s setup generalizes from  $\mathbf{G}$  to any of its forms  $\mathbf{G}_{\mathbf{C}((z))}$ .

The tori  $\mathbf{A}, \mathbf{G}_\gamma$  generalize to forms  $\mathbf{A}_{\mathbf{C}((z))}, \mathbf{G}_{\mathbf{C}((z)), \gamma}$ .

These have corresponding forms  $\mathbf{A}_{\mathbf{F}_q}, \mathbf{T}_{\mathbf{F}_q}$ .

Deligne–Lusztig studied groups over *finite fields*. But up to Tate twist,

$$\mathrm{Gal}(\bar{\mathbf{F}}_q|\mathbf{F}_q) \simeq \hat{\mathbf{Z}} \simeq \mathrm{Gal}(\overline{\mathbf{C}((z))}|\mathbf{C}((z))).$$

Forms of  $\mathbf{G}$  are classified by Dynkin automorphisms in the same way over  $\mathbf{F}_q$  as over  $\mathbf{C}((z))$ .

Much of Oblomkov–Yun’s setup generalizes from  $\mathbf{G}$  to any of its forms  $\mathbf{G}_{\mathbf{C}((z))}$ .

The tori  $\mathbf{A}, \mathbf{G}_\gamma$  generalize to forms  $\mathbf{A}_{\mathbf{C}((z))}, \mathbf{G}_{\mathbf{C}((z)), \gamma}$ .

These have corresponding forms  $\mathbf{A}_{\mathbf{F}_q}, \mathbf{T}_{\mathbf{F}_q}$ .

2 Deligne–Lusztig Theory Work over  $\bar{\mathbf{F}}_q$  for good  $q$ .

$$\{\text{forms of } \mathbf{G} \text{ over } \mathbf{F}_q\} \leftrightarrow \{\text{Frobenii } F \curvearrowright \mathbf{G}\}$$

We say that  $\mathbf{G} = \mathbf{G}^F$  is a *finite group of Lie type*.

$F$ -stable Levis  $\mathbf{L} \subseteq \mathbf{G}$  correspond to Levis  $\mathbf{L} \subseteq G$ .

Deligne–Lusztig introduced varieties<sup>†</sup>  $\mathbf{Y}_{\mathbf{L}}^{\mathbf{G}}$  such that

$$G \curvearrowright H_c^*(Y_{\mathbf{L}}^{\mathbf{G}}) \curvearrowright L.$$

Induction map  $\mathbf{R}_{\mathbf{L}}^{\mathbf{G}} : K_0(L) \rightarrow K_0(G)$ :

$$\mathbf{R}_{\mathbf{L}}^{\mathbf{G}}(\lambda) = \sum_i (-1)^i H_c^i(Y_{\mathbf{L}}^{\mathbf{G}})[\lambda].$$

<sup>†</sup> Actually,  $\mathbf{Y}_{\mathbf{L}}^{\mathbf{G}}$  depends on a parabolic  $\mathbf{P} \supseteq \mathbf{L}$ .



2 Deligne–Lusztig Theory Work over  $\bar{\mathbf{F}}_q$  for good  $q$ .

$$\{\text{forms of } \mathbf{G} \text{ over } \mathbf{F}_q\} \leftrightarrow \{\text{Frobenii } F \curvearrowright \mathbf{G}\}$$

We say that  $G = \mathbf{G}^F$  is a *finite group of Lie type*.

$F$ -stable Levis  $\mathbf{L} \subseteq \mathbf{G}$  correspond to Levis  $L \subseteq G$ .

Deligne–Lusztig introduced varieties<sup>†</sup>  $Y_{\mathbf{L}}^{\mathbf{G}}$  such that

$$G \curvearrowright H_c^*(Y_{\mathbf{L}}^{\mathbf{G}}) \curvearrowright L.$$

Induction map  $R_L^{\mathbf{G}} : K_0(L) \rightarrow K_0(G)$ :

$$R_L^{\mathbf{G}}(\lambda) = \sum_i (-1)^i H_c^i(Y_{\mathbf{L}}^{\mathbf{G}})[\lambda].$$

<sup>†</sup> Actually,  $Y_{\mathbf{L}}^{\mathbf{G}}$  depends on a parabolic  $\mathbf{P} \supseteq \mathbf{L}$ .

(Broué–Malle) For  $m$ -regular maximal tori  $\mathbf{T}$ , a specific algebra  $H_T^G(\mathbf{q})$  such that

$$H_T^G(\zeta_m) = \bar{\mathbf{Q}} W_T^G, \quad \text{where } W_T^G = N_G(T)/T.$$

They conjecture:

$$1 \quad H_T^G(q) \otimes \bar{\mathbf{Q}}_\ell \simeq \text{End}_G(H_c^*(Y_{\mathbf{T}}^{\mathbf{G}})[1_T]).$$

2 As a virtual  $(G, H_T^G(q))$ -bimodule,

$$R_T^G(1_T) = \sum_{\substack{\rho \in \text{Irr}(G) \\ (\rho, R_T^G(1_T)) \neq 0}} \varepsilon_{T,\rho}(\rho \otimes \chi_{T,\rho,q})$$

where  $\varepsilon_{T,\rho} \in \{\pm 1\}$  and  $\chi_{T,\rho} \in \text{Irr}(W_T^G)$ .

(And  $\chi_{T,\rho,q} \in K_0(H_T^G(q))$  corresponds to  $\chi_{T,\rho}$ .)

2 Deligne–Lusztig Theory Work over  $\bar{\mathbf{F}}_q$  for good  $q$ .

$$\{\text{forms of } \mathbf{G} \text{ over } \mathbf{F}_q\} \leftrightarrow \{\text{Frobenii } F \curvearrowright \mathbf{G}\}$$

We say that  $G = \mathbf{G}^F$  is a *finite group of Lie type*.

$F$ -stable Levis  $\mathbf{L} \subseteq \mathbf{G}$  correspond to Levis  $L \subseteq G$ .

Deligne–Lusztig introduced varieties<sup>†</sup>  $Y_{\mathbf{L}}^{\mathbf{G}}$  such that

$$G \curvearrowright H_c^*(Y_{\mathbf{L}}^{\mathbf{G}}) \curvearrowright L.$$

Induction map  $R_L^G : K_0(L) \rightarrow K_0(G)$ :

$$R_L^G(\lambda) = \sum_i (-1)^i H_c^i(Y_{\mathbf{L}}^{\mathbf{G}})[\lambda].$$

<sup>†</sup> Actually,  $Y_{\mathbf{L}}^{\mathbf{G}}$  depends on a parabolic  $\mathbf{P} \supseteq \mathbf{L}$ .

(Broué–Malle) For  $m$ -regular maximal tori  $\mathbf{T}$ , a specific algebra  $H_T^G(\mathfrak{q})$  such that

$$H_T^G(\zeta_m) = \bar{\mathbf{Q}} W_T^G, \quad \text{where } W_T^G = N_G(T)/T.$$

They conjecture:

$$1 \quad H_T^G(q) \otimes \bar{\mathbf{Q}}_\ell \simeq \text{End}_G(H_c^*(Y_{\mathbf{T}}^G)[1_T]).$$

$$2 \quad \text{As a virtual } (G, H_T^G(q))\text{-bimodule,}$$

$$R_T^G(1_T) = \sum_{\substack{\rho \in \text{Irr}(G) \\ (\rho, R_T^G(1_T)) \neq 0}} \varepsilon_{T,\rho} (\rho \otimes \chi_{T,\rho,q})$$

where  $\varepsilon_{T,\rho} \in \{\pm 1\}$  and  $\chi_{T,\rho} \in \text{Irr}(W_T^G)$ .

(And  $\chi_{T,\rho,q} \in K_0(H_T^G(q))$  corresponds to  $\chi_{T,\rho} \cdot$ )

(Broué–Malle) For  $m$ -regular maximal tori  $\mathbf{T}$ , a specific algebra  $H_T^G(\mathbf{q})$  such that

$$H_T^G(\zeta_m) = \bar{\mathbf{Q}} W_T^G, \quad \text{where } W_T^G = N_G(T)/T.$$

They conjecture:

- 1  $H_T^G(q) \otimes \bar{\mathbf{Q}}_\ell \simeq \text{End}_G(H_c^*(Y_{\mathbf{T}}^G)[1_T]).$
- 2 As a virtual  $(G, H_T^G(q))$ -bimodule,

$$R_T^G(1_T) = \sum_{\substack{\rho \in \text{Irr}(G) \\ (\rho, R_T^G(1_T)) \neq 0}} \varepsilon_{T, \rho} (\rho \otimes \chi_{T, \rho, q})$$

where  $\varepsilon_{T, \rho} \in \{\pm 1\}$  and  $\chi_{T, \rho} \in \text{Irr}(W_T^G).$

(And  $\chi_{T, \rho, q} \in K_0(H_T^G(q))$  corresponds to  $\chi_{T, \rho}.$ )

Back to Springer.  $(\mathbf{A}_{\mathbf{F}_q}, \mathbf{T}_{\mathbf{F}_q} \leftrightarrow \mathbf{A}_{\mathbf{C}((z))}, \mathbf{G}_{\mathbf{C}((z)), \gamma})$

It turns out that  $\mathbf{A}$  and  $\mathbf{T}$  are 1- and  $m$ -regular.

Moreover,  $\pi_1(\mathbf{c}_{d/m}^{\text{rs}})$  is the braid group of  $W_T^G.$

Conjecture (T–Xue)

- 1  $\pi_1(\mathbf{c}_{d/m}^{\text{rs}}) \curvearrowright \mathcal{E}_\gamma$  factors through  $H_T^G(1).$
- 2 As a virtual  $(D_{d/m}^{\text{rat}}, H_T^G(1))$ -bimodule,<sup>†</sup>

$$E_\gamma = \sum_{\substack{\rho \in \text{Irr}(G) \\ (\rho, R_A^G(1_A)) \neq 0 \\ (\rho, R_T^G(1_T)) \neq 0}} \varepsilon_{T \rho} (\Delta_{d/m}(\chi_{A, \rho}) \otimes \chi_{T, \rho, 1}).$$

<sup>†</sup> In general,  $D_{d/m}^{\text{rat}}$  is defined using  $W_A^G.$

Back to Springer.  $(\mathbf{A}_{\mathbf{F}_q}, \mathbf{T}_{\mathbf{F}_q} \leftrightarrow \mathbf{A}_{\mathbf{C}((z))}, \mathbf{G}_{\mathbf{C}((z)), \gamma})$

It turns out that  $\mathbf{A}$  and  $\mathbf{T}$  are 1- and  $m$ -regular.

Moreover,  $\pi_1(\mathbf{c}_{d/m}^{\text{rs}})$  is the braid group of  $W_T^G$ .

Conjecture (T-Xue)

- 1  $\pi_1(\mathbf{c}_{d/m}^{\text{rs}}) \curvearrowright \mathcal{E}_\gamma$  factors through  $H_T^G(1)$ .
- 2 As a virtual  $(D_{d/m}^{\text{rat}}, H_T^G(1))$ -bimodule,<sup>†</sup>

$$E_\gamma = \sum_{\substack{\rho \in \text{Irr}(G) \\ (\rho, R_A^G(1_A)) \neq 0 \\ (\rho, R_T^G(1_T)) \neq 0}} \varepsilon_{T\rho}(\Delta_{d/m}(\chi_{A,\rho}) \otimes \chi_{T,\rho,1}).$$

<sup>†</sup> In general,  $D_{d/m}^{\text{rat}}$  is defined using  $W_A^G$ .

**Theorem (T–Xue)** True in these cases:

- $m$  is the (twisted) Coxeter number of  $\mathbf{G}_{\mathbf{C}((z))}$ .
- $(\mathbf{G}_{\mathbf{C}((z))}, m) = ({}^2A_2, 2), (C_2, 2), (G_2, 3), (G_2, 2)$ .

Under a conjecture of OY, true in further cases.

**Example** Take  $\mathbf{G}_{\mathbf{C}((z))}$  split,  $m$  its Coxeter number.

$\chi_{A,\rho}$  runs over characters  $\chi_{\wedge^k(\mathbf{a})}$  of  $W_A^G$ .

$\chi_{T,\rho}$  runs over *all* characters of  $W_T^G = \mathbf{Z}/m\mathbf{Z}$ .

In  $K_0(D_{d/m}^{\text{rat}})$ ,

$$\begin{aligned} [E_\gamma] &= \sum_k (-1)^k [\Delta_{d/m}(\chi_{\wedge^k(\mathbf{a})})] \\ &= [L_{d/m}(\chi_{\text{triv}})]. \end{aligned}$$

*Cf.* the BGG resolution of Berest–Etingof–Ginzburg.

Back to Springer.  $(\mathbf{A}_{\mathbf{F}_q}, \mathbf{T}_{\mathbf{F}_q} \leftrightarrow \mathbf{A}_{\mathbf{C}((z))}, \mathbf{G}_{\mathbf{C}((z)), \gamma})$

It turns out that  $\mathbf{A}$  and  $\mathbf{T}$  are 1- and  $m$ -regular.

Moreover,  $\pi_1(\mathbf{c}_{d/m}^{\text{rs}})$  is the braid group of  $W_T^G$ .

**Conjecture (T–Xue)**

- 1  $\pi_1(\mathbf{c}_{d/m}^{\text{rs}}) \curvearrowright \mathcal{E}_\gamma$  factors through  $H_T^G(1)$ .
- 2 As a virtual  $(D_{d/m}^{\text{rat}}, H_T^G(1))$ -bimodule,<sup>†</sup>

$$E_\gamma = \sum_{\substack{\rho \in \text{Irr}(G) \\ (\rho, R_A^G(1_A)) \neq 0 \\ (\rho, R_T^G(1_T)) \neq 0}} \varepsilon_{T\rho}(\Delta_{d/m}(\chi_{A,\rho}) \otimes \chi_{T,\rho,1}).$$

<sup>†</sup> In general,  $D_{d/m}^{\text{rat}}$  is defined using  $W_A^G$ .

Theorem (T–Xue) True in these cases:

- $m$  is the (twisted) Coxeter number of  $\mathbf{G}_{\mathbf{C}((z))}$ .
- $(\mathbf{G}_{\mathbf{C}((z))}, m) = ({}^2A_2, 2), (C_2, 2), (G_2, 3), (G_2, 2)$ .

Under a conjecture of OY, true in further cases.

Example Take  $\mathbf{G}_{\mathbf{C}((z))}$  split,  $m$  its Coxeter number.

$\chi_{A,\rho}$  runs over characters  $\chi_{\wedge^k(\mathbf{a})}$  of  $W_A^G$ .

$\chi_{T,\rho}$  runs over *all* characters of  $W_T^G = \mathbf{Z}/m\mathbf{Z}$ .

In  $K_0(D_{d/m}^{\text{rat}})$ ,

$$\begin{aligned} [E_\gamma] &= \sum_k (-1)^k [\Delta_{d/m}(\chi_{\wedge^k(\mathbf{a})})] \\ &= [L_{d/m}(\chi_{\text{triv}})]. \end{aligned}$$

*Cf.* the BGG resolution of Berest–Etingof–Ginzburg.

Theorem (T-Xue) True in these cases:

- $m$  is the (twisted) Coxeter number of  $\mathbf{G}_{\mathbf{C}((z))}$ .
- $(\mathbf{G}_{\mathbf{C}((z))}, m) = ({}^2A_2, 2), (C_2, 2), (G_2, 3), (G_2, 2)$ .

Under a conjecture of OY, true in further cases.

Example Take  $\mathbf{G}_{\mathbf{C}((z))}$  split,  $m$  its Coxeter number.

$\chi_{A,\rho}$  runs over characters  $\chi_{\wedge^k(\mathbf{a})}$  of  $W_A^G$ .

$\chi_{T,\rho}$  runs over *all* characters of  $W_T^G = \mathbf{Z}/m\mathbf{Z}$ .

In  $K_0(D_{d/m}^{\text{rat}})$ ,

$$\begin{aligned} [E_\gamma] &= \sum_k (-1)^k [\Delta_{d/m}(\chi_{\wedge^k(\mathbf{a})})] \\ &= [L_{d/m}(\chi_{\text{triv}})]. \end{aligned}$$

Cf. the BGG resolution of Berest–Etingof–Ginzburg.

3 Level-Rank Duality Compare  $E_\gamma$  given by

$$\sum_\rho \varepsilon_{T,\rho}(\Delta_{d/m}(\chi_{A,\rho}) \otimes \chi_{T,\rho,1})$$

with  $R_A^G(1_A) \otimes_{\bar{\mathbf{Q}}_\ell G} R_T^G(1_T)$  given by

$$\sum_\rho \varepsilon_{T,\rho}(\chi_{A,\rho,q} \otimes \chi_{T,\rho,q}).$$

The *Knizhnik–Zamolodchik functor*

$$\mathbf{KZ} : \text{Rep}(D_{d/m}^{\text{rat}}) \rightarrow \text{Rep}(H_A^G(\zeta_m))$$

sends  $\mathbf{KZ}(\Delta_{d/m}(\chi)) = \chi_{\zeta_m}$ . Thus an analogy:

$$\boxed{\mathbf{F}_q : (q, q) :: \mathbf{C}((z)) : (\zeta_m, 1)}$$

The symmetry between  $A$  and  $T$  led us to new discoveries about the Harish–Chandra theory of  $G$ .



3 Level-Rank Duality Compare  $E_\gamma$  given by

$$\sum_{\rho} \varepsilon_{T,\rho}(\Delta_{d/m}(\chi_{A,\rho}) \otimes \chi_{T,\rho,1})$$

with  $R_A^G(1_A) \otimes_{\bar{\mathbf{Q}}_\ell G} R_T^G(1_T)$  given by

$$\sum_{\rho} \varepsilon_{T,\rho}(\chi_{A,\rho,q} \otimes \chi_{T,\rho,q}).$$

The *Knizhnik–Zamolodchik functor*

$$\textcolor{red}{\mathbf{KZ}} : \text{Rep}(D_{d/m}^{\text{rat}}) \rightarrow \text{Rep}(H_A^G(\zeta_m))$$

sends  $\mathbf{KZ}(\Delta_{d/m}(\chi)) = \chi_{\zeta_m}$ . Thus an analogy:

$\mathbf{F}_q : (q, q) :: \mathbf{C}((z)) : (\zeta_m, 1)$

The symmetry between  $A$  and  $T$  led us to new discoveries about the Harish–Chandra theory of  $G$ .

Let  $\text{Uch}(G)$  be the set of *unipotent* irreps of  $G$ , which occur in  $R_T^G(1_T)$  for some maximal torus  $\mathbf{T}$ .

(Broué–Malle–Michel) Fix a positive integer  $l$ .

- $\mathbf{L} \subseteq \mathbf{G}$  is *l-split* iff  $\mathbf{L} = Z_{\mathbf{G}}(\mathbf{S})^\circ$ , where

$\mathbf{S}$  is a torus with  $|S|$  a power of  $\Phi_l(q)$ .

- $\lambda \in \text{Uch}(L)$  is *l-cuspidal* iff  $(\lambda, R_M^G(\mu)) = 0$  for any  $l$ -split  $M \neq L$ .

As we run over pairs  $(\mathbf{L}, \lambda)$  up to conjugacy,

$$\text{Uch}(G) = \coprod \text{Uch}(G)_{\mathbf{L}, \lambda},$$

where  $\text{Uch}(G)_{\mathbf{L}, \lambda} = \{\rho \mid (\rho, R_L^G(\lambda)) \neq 0\}$ .

For  $l = 1$ , these are classical *Harish–Chandra series*.

3 Level-Rank Duality Compare  $E_\gamma$  given by

$$\sum_{\rho} \varepsilon_{T, \rho}(\Delta_{d/m}(\chi_{A, \rho}) \otimes \chi_{T, \rho, 1})$$

with  $R_A^G(1_A) \otimes_{\bar{\mathbf{Q}}_\ell G} R_T^G(1_T)$  given by

$$\sum_{\rho} \varepsilon_{T, \rho}(\chi_{A, \rho, q} \otimes \chi_{T, \rho, q}).$$

The *Knizhnik–Zamolodchik functor*

$$\text{KZ} : \text{Rep}(D_{d/m}^{\text{rat}}) \rightarrow \text{Rep}(H_A^G(\zeta_m))$$

sends  $\text{KZ}(\Delta_{d/m}(\chi)) = \chi_{\zeta_m}$ . Thus an analogy:

$$\boxed{\mathbf{F}_q : (q, q) :: \mathbf{C}((z)) : (\zeta_m, 1)}$$

The symmetry between  $A$  and  $T$  led us to new discoveries about the Harish–Chandra theory of  $G$ .

Let  $\text{Uch}(G)$  be the set of *unipotent* irreps of  $G$ , which occur in  $R_T^G(1_T)$  for some maximal torus  $\mathbf{T}$ .

(Broué–Malle–Michel) Fix a positive integer  $l$ .

- $\mathbf{L} \subseteq \mathbf{G}$  is  *$l$ -split* iff  $\mathbf{L} = Z_{\mathbf{G}}(\mathbf{S})^\circ$ , where

$\mathbf{S}$  is a torus with  $|S|$  a power of  $\Phi_l(q)$ .

- $\lambda \in \text{Uch}(L)$  is  *$l$ -cuspidal* iff  $(\lambda, R_M^G(\mu)) = 0$  for any  $l$ -split  $M \neq L$ .

As we run over pairs  $(\mathbf{L}, \lambda)$  up to conjugacy,

$$\text{Uch}(G) = \coprod \text{Uch}(G)_{\mathbf{L}, \lambda},$$

where  $\text{Uch}(G)_{\mathbf{L}, \lambda} = \{\rho \mid (\rho, R_L^G(\lambda)) \neq 0\}$ .

For  $l = 1$ , these are classical *Harish-Chandra series*.

Let  $\text{Uch}(G)$  be the set of *unipotent* irreps of  $G$ , which occur in  $R_T^G(1_T)$  for some maximal torus  $\mathbf{T}$ .

(Broué–Malle–Michel) Fix a positive integer  $l$ .

- $\mathbf{L} \subseteq \mathbf{G}$  is *l-split* iff  $\mathbf{L} = Z_{\mathbf{G}}(\mathbf{S})^\circ$ , where

$\mathbf{S}$  is a torus with  $|S|$  a power of  $\Phi_l(q)$ .

- $\lambda \in \text{Uch}(L)$  is *l-cuspidal* iff  $(\lambda, R_M^G(\mu)) = 0$  for any  $l$ -split  $M \neq L$ .

As we run over pairs  $(\mathbf{L}, \lambda)$  up to conjugacy,

$$\text{Uch}(G) = \coprod \text{Uch}(G)_{\mathbf{L}, \lambda},$$

where  $\text{Uch}(G)_{\mathbf{L}, \lambda} = \{\rho \mid (\rho, R_L^G(\lambda)) \neq 0\}$ .

For  $l = 1$ , these are classical *Harish-Chandra series*.

Generalizing our discussion for maximal tori:

Broué–Malle define a Hecke algebra  $H_{L, \lambda}^G(\mathbf{q})$  such that

$$H_{L, \lambda}^G(\zeta_l) = \bar{\mathbf{Q}} W_{L, \lambda}^G, \text{ where } W_{L, \lambda}^G = N_G(L, \lambda)/L.$$

They conjecture:

- 1  $H_{L, \lambda}^G(q) \otimes \bar{\mathbf{Q}}_\ell = \text{End}_G(H_c^*(Y_{\mathbf{L}}^{\mathbf{G}})[\lambda]).$
- 2 As a virtual  $(G, H_{L, \lambda}^G(q))$ -bimodule,

$$R_L^G(\lambda) = \sum_{\rho \in \text{Uch}(G)_{\mathbf{L}, \lambda}} \varepsilon_{L, \lambda, \rho} (\rho \otimes \chi_{L, \lambda, \rho, q})$$

where  $\varepsilon_{L, \lambda, \rho} \in \{\pm 1\}$  and  $\chi_{L, \lambda, \rho} \in \text{Irr}(W_{L, \lambda}^G).$

Generalizing our discussion for maximal tori:

Broué–Malle define a Hecke algebra  $H_{L,\lambda}^G(\mathfrak{q})$  such that

$$H_{L,\lambda}^G(\zeta_l) = \bar{\mathbf{Q}} W_{L,\lambda}^G, \text{ where } W_{L,\lambda}^G = N_G(L, \lambda)/L.$$

They conjecture:

- 1  $H_{L,\lambda}^G(q) \otimes \bar{\mathbf{Q}}_\ell = \text{End}_G(H_c^*(Y_{\mathbf{L}}^G)[\lambda]).$
- 2 As a virtual  $(G, H_{L,\lambda}^G(q))$ -bimodule,

$$R_L^G(\lambda) = \sum_{\rho \in \text{Uch}(G)_{\mathbf{L},\lambda}} \varepsilon_{L,\lambda,\rho}(\rho \otimes \chi_{L,\lambda,\rho,q})$$

where  $\varepsilon_{L,\lambda,\rho} \in \{\pm 1\}$  and  $\chi_{L,\lambda,\rho} \in \text{Irr}(W_{L,\lambda}^G).$

Via the *decomposition map*

$$\chi \mapsto \chi_{\zeta_m} : \text{Irr}(W_{L,\lambda}^G) \rightarrow K_0(H_{L,\lambda}^G(\zeta_m)),$$

we partition  $\text{Irr}(W_{L,\lambda}^G)$  into *blocks*, describing how  $H_{L,\lambda}^G(\zeta_m)$  fails to be semisimple.

**Conjecture (T–Xue)** Fix  $l, m$ .

Fix an  $l$ -cuspidal  $(\mathbf{L}, \lambda)$  and  $m$ -cuspidal  $(\mathbf{M}, \mu)$ .

1 The set

$$\{\chi_{L,\lambda,\rho} \mid \rho \in \text{Uch}(G)_{\mathbf{L},\lambda} \cap \text{Uch}(G)_{\mathbf{M},\mu}\},$$

resp.  $\{\chi_{M,\mu,\rho} \mid \rho \in \text{Uch}(G)_{\mathbf{L},\lambda} \cap \text{Uch}(G)_{\mathbf{M},\mu}\},$

is a union of  $H_{L,\lambda}^G(\zeta_m)$ -, resp.  $H_{M,\mu}^G(\zeta_l)$ -blocks.

2 The indexing induces a matching of blocks.

Generalizing our discussion for maximal tori:

Broué–Malle define a Hecke algebra  $H_{L,\lambda}^G(\mathbf{q})$  such that

$$H_{L,\lambda}^G(\zeta_l) = \bar{\mathbf{Q}} W_{L,\lambda}^G, \text{ where } W_{L,\lambda}^G = N_G(L, \lambda)/L.$$

They conjecture:

$$1 \quad H_{L,\lambda}^G(q) \otimes \bar{\mathbf{Q}}_\ell = \text{End}_G(H_c^*(Y_{\mathbf{L}}^G)[\lambda]).$$

$$2 \quad \text{As a virtual } (G, H_{L,\lambda}^G(q))\text{-bimodule,}$$

$$R_L^G(\lambda) = \sum_{\rho \in \text{Uch}(G)_{\mathbf{L},\lambda}} \varepsilon_{L,\lambda,\rho} (\rho \otimes \chi_{L,\lambda,\rho,q})$$

where  $\varepsilon_{L,\lambda,\rho} \in \{\pm 1\}$  and  $\chi_{L,\lambda,\rho} \in \text{Irr}(W_{L,\lambda}^G)$ .

Via the *decomposition map*

$$\chi \mapsto \chi_{\zeta_m} : \text{Irr}(W_{L,\lambda}^G) \rightarrow K_0(H_{L,\lambda}^G(\zeta_m)),$$

we partition  $\text{Irr}(W_{L,\lambda}^G)$  into *blocks*, describing how  $H_{L,\lambda}^G(\zeta_m)$  fails to be semisimple.

**Conjecture (T-Xue)** Fix  $l, m$ .

Fix an  $l$ -cuspidal  $(\mathbf{L}, \lambda)$  and  $m$ -cuspidal  $(\mathbf{M}, \mu)$ .

1 The set

$$\begin{aligned} & \{\chi_{L,\lambda,\rho} \mid \rho \in \text{Uch}(G)_{\mathbf{L},\lambda} \cap \text{Uch}(G)_{\mathbf{M},\mu}\}, \\ \text{resp. } & \{\chi_{M,\mu,\rho} \mid \rho \in \text{Uch}(G)_{\mathbf{L},\lambda} \cap \text{Uch}(G)_{\mathbf{M},\mu}\}, \end{aligned}$$

is a union of  $H_{L,\lambda}^G(\zeta_m)$ -, resp.  $H_{M,\mu}^G(\zeta_l)$ -blocks.

2 The indexing induces a matching of blocks.

Via the *decomposition map*

$$\chi \mapsto \chi_{\zeta_m} : \text{Irr}(W_{L,\lambda}^G) \rightarrow K_0(H_{L,\lambda}^G(\zeta_m)),$$

we partition  $\text{Irr}(W_{L,\lambda}^G)$  into *blocks*, describing how  $H_{L,\lambda}^G(\zeta_m)$  fails to be semisimple.

Conjecture (T–Xue) Fix  $l, m$ .

Fix an  $l$ -cuspidal  $(\mathbf{L}, \lambda)$  and  $m$ -cuspidal  $(\mathbf{M}, \mu)$ .

1 The set

$$\{\chi_{L,\lambda,\rho} \mid \rho \in \text{Uch}(G)_{\mathbf{L},\lambda} \cap \text{Uch}(G)_{\mathbf{M},\mu}\},$$

$$\text{resp. } \{\chi_{M,\mu,\rho} \mid \rho \in \text{Uch}(G)_{\mathbf{L},\lambda} \cap \text{Uch}(G)_{\mathbf{M},\mu}\},$$

is a union of  $H_{L,\lambda}^G(\zeta_m)$ -, resp.  $H_{M,\mu}^G(\zeta_l)$ -blocks.

2 The indexing induces a matching of blocks.

Theorem (T–Xue) (1), (2) are compatible with block sizes for essentially all  $G, l, m$  with  $G$  exceptional.

Conjecture (T–Xue) In the preceding setup:

3 Via KZ functors, the bijection in (2) lifts to a derived equivalence between category-O blocks of appropriate rational Cherednik algebras.

Theorem (T–Xue) (1), (2), (3) hold for  $G = \text{GL}_n$  when  $l, m$  are coprime.



Theorem (T–Xue) (1), (2) are compatible with block sizes for essentially all  $G, l, m$  with  $G$  exceptional.

Conjecture (T–Xue) In the preceding setup:

3 Via KZ functors, the bijection in (2) lifts to a derived equivalence between category- $\mathcal{O}$  blocks of appropriate rational Cherednik algebras.

Theorem (T–Xue) (1), (2), (3) hold for  $G = \mathrm{GL}_n$  when  $l, m$  are coprime.

Note that  $W_{L,\lambda}^{\mathrm{GL}_n} \simeq S_N \ltimes \mathbf{Z}_l^N$  for some  $N$ , *etc.*

$$\mathrm{Rep}(H_{L,\lambda}^{\mathrm{GL}_n}(\zeta_m)) \quad \text{and} \quad \mathrm{Rep}(H_{M\mu}^{\mathrm{GL}_n}(\zeta_l))$$

can be interpreted in terms of *higher-level Fock spaces*

$$\bigoplus_{\substack{\vec{s} \in \mathbf{Z}^l \\ |\vec{s}|=s}} \Lambda_{\mathbf{q}}^{\vec{s}} \xleftarrow{\sim} \Lambda_{\mathbf{q}}^s \xrightarrow{\sim} \bigoplus_{\substack{\vec{r} \in \mathbf{Z}^m \\ |\vec{r}|=s}} \Lambda_{\mathbf{q}}^{\vec{r}}.$$

Above,  $\Lambda_{\mathbf{q}}^{\vec{s}} \simeq \bigoplus_N K_0(S_N \ltimes \mathbf{Z}_l^N) \otimes \mathbf{Q}(\mathbf{q})$ , *etc.*

*Level-rank duality* of Frenkel, Uglov, Chuang–Miyachi, Rouquier–Shan–Varagnolo–Vasserot. . .

Our conjectures generalize level-rank duality from  $\mathrm{GL}_n$  to arbitrary  $G$ .

Theorem (T–Xue) (1), (2) are compatible with block sizes for essentially all  $G, l, m$  with  $G$  exceptional.

Conjecture (T–Xue) In the preceding setup:

3 Via KZ functors, the bijection in (2) lifts to a derived equivalence between category-O blocks of appropriate rational Cherednik algebras.

Theorem (T–Xue) (1), (2), (3) hold for  $G = \mathrm{GL}_n$  when  $l, m$  are coprime.

Note that  $W_{L,\lambda}^{\mathrm{GL}_n} \simeq S_N \ltimes \mathbf{Z}_l^N$  for some  $N$ , *etc.*

$$\mathrm{Rep}(H_{L,\lambda}^{\mathrm{GL}_n}(\zeta_m)) \quad \text{and} \quad \mathrm{Rep}(H_{M\mu}^{\mathrm{GL}_n}(\zeta_l))$$

can be interpreted in terms of *higher-level Fock spaces*

$$\bigoplus_{\substack{\vec{s} \in \mathbf{Z}^l \\ |\vec{s}|=s}} \Lambda_{\mathbf{q}}^{\vec{s}} \xleftarrow{\sim} \Lambda_{\mathbf{q}}^s \xrightarrow{\sim} \bigoplus_{\substack{\vec{r} \in \mathbf{Z}^m \\ |\vec{r}|=s}} \Lambda_{\mathbf{q}}^{\vec{r}}.$$

Above,  $\Lambda_{\mathbf{q}}^{\vec{s}} \simeq \bigoplus_N K_0(S_N \ltimes \mathbf{Z}_l^N) \otimes \mathbf{Q}(\mathbf{q})$ , *etc.*

*Level-rank duality* of Frenkel, Uglov, Chuang–Miyachi, Rouquier–Shan–Varagnolo–Vasserot. . .

Our conjectures generalize level-rank duality from  $\mathrm{GL}_n$  to arbitrary  $G$ .

Note that  $W_{L,\lambda}^{\mathrm{GL}_n} \simeq S_N \ltimes \mathbf{Z}_l^N$  for some  $N$ , *etc.*

$$\mathrm{Rep}(H_{L,\lambda}^{\mathrm{GL}_n}(\zeta_m)) \quad \text{and} \quad \mathrm{Rep}(H_{M\mu}^{\mathrm{GL}_n}(\zeta_l))$$

can be interpreted in terms of *higher-level Fock spaces*

$$\bigoplus_{\substack{\vec{s} \in \mathbf{Z}^l \\ |\vec{s}|=s}} \Lambda_{\mathbf{q}}^{\vec{s}} \xleftarrow{\sim} \Lambda_{\mathbf{q}}^s \xrightarrow{\sim} \bigoplus_{\substack{\vec{r} \in \mathbf{Z}^m \\ |\vec{r}|=s}} \Lambda_{\mathbf{q}}^{\vec{r}}.$$

*Thank you for listening.*

Above,  $\Lambda_{\mathbf{q}}^{\vec{s}} \simeq \bigoplus_N K_0(S_N \ltimes \mathbf{Z}_l^N) \otimes \mathbf{Q}(\mathbf{q})$ , *etc.*

*Level-rank duality* of Frenkel, Uglov, Chuang–Miyachi,  
Rouquier–Shan–Varagnolo–Vasserot. . .

Our conjectures generalize level-rank duality from  
 $\mathrm{GL}_n$  to arbitrary  $G$ .