

MATH 251: TOPOLOGY II
SPRING 2026 PRACTICE PROBLEMS

CONTENTS

0. Review of Topology I	2
Exam 1 Topics	4
1. Fundamental Groups and Covering Spaces	5
Exam 2 Topics	9
2. Fixed Points and Separations in the Plane	10
3. Free Groups and Free Products	12
Exam 3 Topics	15
4. Simplicial Complexes and Surfaces	16

NOTE: All citations are to Munkres's textbook, *Topology*, 2nd Edition, unless indicated otherwise. When a problem statement has a proof in Munkres, try your best to find your own proof, before comparing with his.

For any integer $n > 0$, the topology on $\mathbf{R}^n$ is the analytic topology unless otherwise specified.

0. REVIEW OF TOPOLOGY I

Problem 1. Show that no two of the spaces

$$(0, 1), \quad (0, 1], \quad [0, 1]$$

are homeomorphic. *Hint:* What happens if you remove any point from $(0, 1)$?

Problem 2. Suppose that $A \subseteq X$. Recall that the *interior* of A in X is

$$\text{Int}_X(A) := \{x \in X \mid x \in U \text{ for some } U \text{ open in } X \text{ such that } U \subseteq A\}$$

and the *closure* of A in X is

$$\text{Cl}_X(A) := \{x \in X \mid x \in K \text{ for all } K \text{ closed in } X \text{ such that } K \supseteq A\}.$$

Show the identity $\text{Cl}_X(A) = X \setminus \text{Int}_X(X \setminus A)$.

Problem 3. Suppose that $A \subseteq Y \subseteq X$, where Y is a subspace of X .

- (1) Show that $\text{Int}_Y(A) \supseteq \text{Int}_X(A)$.
- (2) Give an example where $\text{Int}_Y(A) \neq \text{Int}_X(A)$. *Hint:* You can assume $Y = A$.

Problem 4. Let X be the real line $\mathbf{R}$ in its finite complement, or *cofinite*, topology. Show that every sequence of points in X converges to every point of X simultaneously. Deduce that X is not Hausdorff.

Problem 5. Show that any metric space is Hausdorff.

Problem 6. Show that for any integer $n \geq 1$, the analytic topology on $\mathbf{R}^n$ matches the product topology on $\mathbf{R} \times \cdots \times \mathbf{R}$, where there are n factors.

Problem 7. Let $p: \mathbf{R} \rightarrow S^1$ be the map

$$p(t) = (\cos(2\pi t), \sin(2\pi t)).$$

For any $a, b \in \mathbf{R}$, we define the *(open) arc* $J_{a,b} \subseteq S^1$ to be

$$J_{a,b} = \{p(t) \mid a < t < b\}.$$

Show that the collection of arcs $\{J_{a,b} \mid a, b \in \mathbf{R}\}$ satisfies the definition of a basis (Munkres page 78).

Problem 8. Show that the following topologies on S^1 are all the same:

- The topology generated by the basis $\{J_{a,b} \mid a, b \in \mathbf{R}\}$ in Problem 7.
- The subspace topology that S^1 inherits from its inclusion into $\mathbf{R}^2$.
- The quotient topology that S^1 inherits from the surjective map $p: \mathbf{R} \rightarrow S^1$.

Problem 9. Let $f_1, f_2, f_3: A \rightarrow X$ be continuous maps. Suppose that φ is a homotopy from f_1 to f_2 and ψ is a homotopy from f_2 to f_3 . Construct a homotopy from f_1 to f_3 explicitly in terms of φ and ψ .

Problem 10. In the setup of Problem 9, suppose that

$$\begin{aligned} A &= [0, 1], \\ f_1(0) &= f_2(0) = f_3(0), \\ f_1(1) &= f_2(1) = f_3(1). \end{aligned}$$

Show that in this case, if φ and ψ are path homotopies, then we can choose the solution of (1) to be a path homotopy as well.

Problem 11. Let $f, g: X \rightarrow Y$ and $F, G: Y \rightarrow Z$ be continuous. Show that

$$\text{if } f \sim g \text{ and } F \sim G, \quad \text{then } F \circ f \sim G \circ g.$$

Problem 12. Use the $f = g$ case of Problem 11 to show: If X, Y are nonempty and Y is contractible, then any continuous map from X into Y is *nulhomotopic*: that is, homotopic to a constant map.

Problem 13. Give an explicit homotopy equivalence between S^1 and $\mathbf{R}^2 \setminus \{(0, 0)\}$, and show that it is indeed a homotopy equivalence.

Problem 14. Suppose that:

- $f: X \rightarrow Y$ and $g: Y \rightarrow X$ define a homotopy equivalence between X and Y .
- $j: Y \rightarrow Z$ and $k: Z \rightarrow Y$ define a homotopy equivalence between Y and Z .

Show that $j \circ f: X \rightarrow Z$ and $g \circ k: Z \rightarrow X$ define a homotopy equivalence between X and Z .

Hint: Among other things, you will need to show that $g \circ k \circ j \circ f \sim \text{id}_X$. But we are given that $k \circ j \sim \text{id}_Y$. Use Problem 11 to show that $k \circ j \sim \text{id}_Y$ implies $g \circ k \circ j \circ f \sim \text{id}_X$.

Problem 15. Show that if the identity map of a space X is homotopic to a constant map with value $c \in X$, then X is homotopy equivalent to $\{c\}$. Deduce that all *contractible* spaces are homotopy equivalent to each other.

Problem 16. A subset $A \subseteq \mathbf{R}^n$ is *star convex* if and only if there is some point $c \in A$ such that the line segment between c and any other point of A is contained within A . That is, for any $a \in A$ and $t \in [0, 1]$, we have $(1 - t)c + ta \in A$.

- (1) Show that if A is star convex, then A is contractible.
- (2) Show that if A is star convex, then any loop in A based at a_0 is path homotopic to the constant loop.
- (3) Give a star convex subset of $\mathbf{R}^2$ that is not convex.

EXAM 1 TOPICS

Exam 1 will be a **closed-book** exam. It will be held in-class, 12:40–2:00 pm, on ~~January 26~~ ~~February 2~~ **February 4**.

You should know by heart the definitions of the following terms. You should be able to state multiple examples of each (with justification), and in some cases, non-examples.

- Munkres §13. Bases
- §18. Continuous maps
- §15. The product topology (only on finite direct products)
- §16. Subspaces
- §22. Quotient spaces
- §17. Interiors and closures
- §23–24. Separations, connectedness
- §23. Paths, path connectedness
- §26–27 Compactness
- §51. Homotopies, path homotopies
- §58. Homotopy equivalences

1. FUNDAMENTAL GROUPS AND COVERING SPACES

Problem 17. Show that if X is path connected, then X is connected.

Problem 18. We say that X is *totally disconnected* if and only if its only nonempty connected subspaces are one-point sets.

- (1) Show that if X is discrete, then X is totally disconnected.
- (2) Show that the set of rational numbers $\mathbf{Q}$, as a subspace of (analytic) $\mathbf{R}$, is totally disconnected, but not discrete.
- (3) Show that if X is totally disconnected, then $\pi_1(X, x)$ is trivial for all $x \in X$.

Hint: Use Problem 17.

Problem 19. Consider the points $p = (1, 0)$ and $q = (-1, 0)$ in $\mathbf{R}^2$.

- (1) Give two paths in S^1 from p to q that are not path homotopic.
- (2) Explain why the paths in (1) become path homotopic when viewed as paths in $\mathbf{R}^2$.

Problem 20. Let $x_0, x_1 \in X$. Let $\alpha: [0, 1] \rightarrow X$ be a path from x_0 to x_1 , and let $\bar{\alpha}$ be the reverse path defined by $\bar{\alpha}(s) = \alpha(1 - s)$.

- (1) Show that $\alpha * \bar{\alpha}$ is path homotopic to e_{x_0} .
- (2) Using part (1), show that the map

$$\hat{\alpha}: \pi_1(X, x_0) \rightarrow \pi_1(X, x_1) \quad \text{given by } \hat{\alpha}([\gamma]) = [\bar{\alpha} * \gamma * \alpha]$$

is a group homomorphism.

- (3) Show that $\hat{\hat{\alpha}}$ is a two-sided inverse of $\hat{\alpha}$. Deduce that $\hat{\alpha}$ is an isomorphism.

Problem 21. Let $f: X \rightarrow Y$ be a continuous map.

- (1) Show that if $\beta, \gamma: [0, 1] \rightarrow X$ are paths such that $\beta(1) = \gamma(0)$, then

$$f \circ (\beta * \gamma) = (f \circ \beta) * (f \circ \gamma).$$

- (2) Let $x \in X$. Using part (1), show that the map

$$f_*: \pi_1(X, x) \rightarrow \pi_1(Y, f(x)) \quad \text{given by } f_*([\gamma]) = [f \circ \gamma]$$

is a group homomorphism.

- (3) Let $g: Y \rightarrow Z$ be another continuous map. Show that $(g \circ f)_* = g_* \circ f_*$ as homomorphisms from $\pi_1(X, x)$ into $\pi_1(Z, g(f(x)))$.

Problem 22. Let X be a subspace of some space X' , and let $f: X \rightarrow Y$ be a continuous map. Suppose that

$$f = f'|_X \quad \text{for some continuous map } f': X' \rightarrow Y,$$

and that X' is *simply connected*, meaning X' is path connected and its fundamental group is trivial. Show that in this case, the homomorphism $f_*: \pi_1(X, x) \rightarrow \pi_1(Y, f(x))$ is *trivial* for all $x \in X$, meaning it sends every element of $\pi_1(X, x)$ to the identity element of $\pi_1(Y, f(x))$. *Hint:* Use part (3) of Problem 21.

Problem 23. Let X_1, X_2 be spaces with points $x_1 \in X_1$ and $x_2 \in X_2$. Let $X = X_1 \times X_2$ and $x = (x_1, x_2) \in X$. For $i = 1, 2$, let $\text{pr}_i: X \rightarrow X_i$ be the appropriate *projection map* (Munkres page 87).

(1) Show that the map

$$\varphi: \pi_1(X, x) \rightarrow \pi_1(X_1, x_1) \times \pi_1(X_2, x_2) \quad \text{given by } \varphi([\gamma]) = ([\text{pr}_1 \circ \gamma], [\text{pr}_2 \circ \gamma])$$

is a homomorphism.

(2) Show that the map

$$\psi: \pi_1(X_1, x_1) \times \pi_1(X_2, x_2) \rightarrow \pi_1(X, x) \quad \text{given by } \psi([\gamma_1], [\gamma_2]) = [\gamma_1 \times \gamma_2]$$

is a two-sided inverse to φ . Deduce that

$$\pi_1(X, x) \simeq \pi_1(X_1, x_1) \times \pi_1(X_2, x_2).$$

(3) Using the conclusion to part (2), show that $\mathbf{Z}^n$ is the fundamental group of an explicit (path-connected) topological space, for any integer $n > 0$.

Problem 24. By considering what happens on fundamental groups when a point is removed, show that $\mathbf{R}^2$ and $\mathbf{R}^n$ are not homeomorphic if $n > 2$.

Problem 25. Learn the definitions of a *free group* and of a *presentation of a group (by generators and relations)*. (They are covered in Munkres §69, but you may find other sources online helpful as well.)

Let F_2 denote a free group on generators a and b . Show that the homomorphism

$$\varphi: F_2 \rightarrow \mathbf{Z}^2 \quad \text{given by } \varphi(a) = (1, 0) \text{ and } \varphi(b) = (0, 1)$$

is surjective and that its kernel contains the *commutator* of a and b : that is, the element $aba^{-1}b^{-1}$.

(In fact, the kernel is exactly the smallest normal subgroup of F_2 containing the commutator. This shows that $\mathbf{Z}^2$ has a presentation with generators a, b and a single relation $aba^{-1}b^{-1}$.)

Problem 26. Suppose that A, U_1, U_2 are path-connected open subsets of X such that $X = U_1 \cup U_2$ and $A = U_1 \cap U_2$. For $k = 1, 2$, let $i_k: A \rightarrow U_k$ and $j_k: U_k \rightarrow X$ be the appropriate inclusion maps. Fix a point $x \in A$.

(1) Describe a homomorphism

$$\Psi: \pi_1(U_1, x) \star \pi_1(U_2, x) \rightarrow \pi_1(X, x)$$

such that $\Psi|_{\pi_1(U_k, x)} = (j_k)_*$ for $k = 1, 2$. (It is necessarily unique.)

(2) Show that for any $[\gamma] \in \pi_1(A, x)$, the element

$$(i_1)_*([\gamma]) * (i_2)_*([\gamma])^{-1} \in \pi_1(U_1, x) \star \pi_1(U_2, x)$$

belongs to the kernel of Ψ . *Hint:* Compare $j_1 \circ i_1$ and $j_2 \circ i_2$.

Problem 27. Recall that $X \sqcup Y$ denotes the disjoint union of X and Y . The *wedge sum* of X and Y at points $x \in X$ and $y \in Y$ is a quotient space

$$X \vee Y = (X \sqcup Y) / \sim,$$

where $\sim$ is an equivalence relation that only identifies two distinct points when those points are x and y . Use Seifert-van Kampen to show that $\pi_1(S^1 \vee S^1, p) \simeq \mathbf{Z} \star \mathbf{Z}$ for any basepoint p .

Problem 28. Keep the hypotheses of Problem 26. Thus, the Seifert–van Kampen theorem applies to the homomorphism Ψ in part (1) of that problem. Give examples where:

- (1) A is simply-connected, but X, U_1, U_2 are not. *Hint:* Problem 27.
- (2) X, U_1, U_2 are simply-connected, but A is not. *Hint:* Munkres §59.
- (3) X and U_1 are simply-connected, but U_2 is not.
- (4) X is simply-connected, but U_1 and U_2 are not.

Problem 29. For each of the following spaces, the fundamental group is either trivial, $\mathbf{Z}$, or the free group $F_2 \simeq \mathbf{Z} \star \mathbf{Z}$. Determine which option is the case. You do not need to give explicit homeomorphisms or homotopy equivalences, but give informal descriptions (or pictures) to support your reasoning.

- (1) $D \times S^1$, where $D = \{(x, y) \in \mathbf{R}^2 \mid x^2 + y^2 < 1\}$.
- (2) The *torus* $S^1 \times S^1$.
- (3) The bounded cylinder $S^1 \times [0, 1]$.
- (4) The unbounded cylinder $S^1 \times \mathbf{R}$.
- (5) The punctured plane $\mathbf{R}^2 - \{p\}$, where p is any point.
- (6) *Harder:* The punctured torus $S^1 \times S^1 - \{q\}$, where q is any point.

Problem 30. Determine the fundamental group of the punctured 3-dimensional space $\mathbf{R}^3 - \{(0, 0, 0)\}$. Note that it is path-connected, so different basepoints give isomorphic fundamental groups.

Hint: Is it homotopy equivalent to a space whose fundamental group you already know?

Problem 31. Let $p: E \rightarrow B$ be a covering map. Show that if B is connected and $p^{-1}(b_0)$ has k elements for some $b_0 \in B$, then $p^{-1}(b)$ has k elements for all $b \in B$. In this case, we say that p is a *k-fold covering*.

Problem 32. Draw every 2-fold covering space of $S^1 \vee S^1$ up to homeomorphism. You do not need to prove that your list is exhaustive.

Note 1: Some covering spaces could be disconnected! *Note 2:* $S^1 \vee S^1$ has a symmetry of order two. If two covering spaces differ by a lift of this symmetry, you do not need to draw both.

Problem 33. Let $\mathbf{R}_+$ be the set of positive numbers. Let $p: \mathbf{R} \times \mathbf{R}_+ \rightarrow \mathbf{R}^2 - \{(0, 0)\}$ be defined by $p(u, r) = (r \cos(2\pi u), r \sin(2\pi u))$. This is a covering map. Find liftings

along p of the following paths in $\mathbf{R}^2 - \{(0, 0)\}$:

$$f(t) = (2 - t, 0),$$

$$g(t) = ((1 + t) \cos(2\pi t), (1 + t) \sin(2\pi t)),$$

$$h = f * g.$$

Sketch (the images of) these paths and their liftings.

Problem 34. Let $p: \mathbf{R} \rightarrow S^1$ be defined by

$$p(t) = (\cos(2\pi t), \sin(2\pi t)).$$

Consider the path in $S^1 \times S^1$ given by

$$f(t) = ((\cos(2\pi t), \sin(2\pi t)), (\cos(4\pi t), \sin(4\pi t))).$$

Find an explicit lifting $\tilde{f}$ of f along $(p, p): \mathbf{R} \times \mathbf{R} \rightarrow S^1 \times S^1$, and sketch it.

Problem 35. Find all *connected* covering spaces of S^1 up to homeomorphism.

EXAM 2 TOPICS

Exam 2 will be an **open-book** exam. It will be held in-class, 12:40–2:00 pm, on March 2.

“Open book” means: You will be allowed to look at Munkres, *Topology*, 2nd Ed., and any notes on paper that you wrote prior to the exam, but you will not be allowed to use electronic devices, including phones, computers, tablets, *etc.*

Topics Included.

- Definition of the fundamental group $\pi_1(X, x)$
- Whether $\pi_1(X, x)$ and $\pi_1(X, x')$ are related, when x, x' belong to the same path component of X or to different path components of X
- Definition of $f_*: \pi_1(X, x) \rightarrow \pi_1(Y, f(x))$, for a (continuous) map $f: X \rightarrow Y$
- Relation between $(g \circ f)_*$ and $g_* \circ f_*$, for maps $f: X \rightarrow Y$ and $g: Y \rightarrow Z$
- Relation between $\pi_1(X, x)$ and $\pi_1(Y, f(x))$, when $f: X \rightarrow Y$ forms part of a homotopy equivalence
- $\pi_1(X, x)$ when $X = \mathbf{R}^N$ or S^N , for different values of N
- $\pi_1(X \times Y, (x, y))$ in terms of $\pi_1(X, x)$ and $\pi_1(Y, y)$
- Explicit loops representing the elements of $\pi_1(S^1, o)$, where $o = (1, 0)$
- Surjectivity statement in the Seifert–van Kampen theorem: relation between $\pi_1(X, x)$ and $\pi_1(U_1, x) \star \pi_1(U_2, x)$ when $X = U_1 \cup U_2$ for some open U_1, U_2 such that $U_1, U_2, U_1 \cap U_2$ are each path connected and $x \in U_1 \cap U_2$
- The formula for $\pi_1(S^1 \vee S^1)$ (via Seifert–van Kampen)
- Variants of Problems 28 and 29 above
- Definition of covering maps
- Why $p(t) = (\cos(2\pi t), \sin(2\pi t))$ defines a covering map $p: \mathbf{R} \rightarrow S^1$, but not if we replace $\mathbf{R}$ by $[0, 1)$ or $[0, 1]$ or $(0, 1)$, *etc.*
- Some examples of coverings of $S^1 \vee S^1$ and of $S^1 \times S^1$
- The path lifting and path-homotopy lifting properties of covering maps (*e.g.*, lifting a path in X along a covering $E \rightarrow X$ to a starting point in E)
- Relation between $\pi_1(X, p(e))$ and $\pi_1(E, e)$, when $p: E \rightarrow X$ is a covering

Topics Not Included.

- The precise definition of a free product of groups
- The kernel statement in the Seifert–van Kampen theorem
- The Jordan separation/curve theorems

2. FIXED POINTS AND SEPARATIONS IN THE PLANE

Problem 36. Recall that a *simple closed curve* in a space X is a subspace of X homeomorphic to S^1 . Find simple closed curves $C, C' \subseteq S^1 \times S^1$ such that:

- (1) $S^1 \times S^1 - C$ is path connected.
- (2) $S^1 \times S^1 - C'$ is not path connected.

Hint: It may help to view $S^1 \times S^1$ as a quotient space of the square $[0, 1] \times [0, 1]$ in which opposite edges have been identified.

Problem 37. Find a 2×2 matrix with strictly positive real entries that does not have any positive real eigenvalues.

Problem 38. Let $f: S^1 \rightarrow S^1$ be continuous and *nulhomotopic*. Show that:

- (1) There is some point $p \in S^1$ such that $f(p) = p$: that is, a *fixed point* of f .
- (2) There is some point $q \in S^1$ such that $f(q) = -q$.

Hint: Theorem 55.5, or the version proved in class.

Problem 39. Recall that if $s: A \rightarrow X$ and $r: X \rightarrow A$ satisfy $r \circ s = \text{id}_A$, then s is called a *section* and r is called a *retraction*.

- (1) Show that in this situation,

s is injective,

r is surjective,

$s_*: \pi_1(A, a) \rightarrow \pi_1(X, s(a))$ is injective for any $a \in A$,

$r_*: \pi_1(X, x) \rightarrow \pi_1(A, r(x))$ is surjective for any $x \in X$.

- (2) Deduce from (1) that there is no retraction from the closed disk

$$B^2 := \{(x, y) \in \mathbf{R}^2 \mid x^2 + y^2 \leq 1\}$$

onto its boundary circle S^1 .

- (3) Deduce from (1) that if $S \subseteq \mathbf{R}^2$ is nonempty, then there is no retraction from $\mathbf{R}^2$ onto $\mathbf{R}^2 - S$. *Hint:* Show by induction on the size of S that the fundamental group of $\mathbf{R}^2 - S$ must be nontrivial.

Problem 40. Fix an integer $n \geq 0$. Writing $\|-\|$ to mean Euclidean distance, let

$$S^n = \{v \in \mathbf{R}^{n+1} \mid \|v\| = 1\}, \quad \text{the } n\text{-sphere,}$$

$$B^{n+1} = \{v \in \mathbf{R}^{n+1} \mid \|v\| \leq 1\}, \quad \text{the closed } n\text{-ball.}$$

Let $\sim$ be the equivalence relation on $S^n \times [0, 1]$ that collapses the subset $S^n \times \{1\}$ to a point: that is, $(v, t) \sim (v', t)$ if and only if either $t = 1$ or $v = v'$.

Write down an explicit homeomorphism from $(S^n \times [0, 1])/\sim$ onto B^{n+1} . You should be able to show that your example is continuous and bijective, but you do not need to show that it is a homeomorphism. *Hint:* Use a generalization of the map in the “(1) $\Rightarrow$ (2)” step of Munkres’s proof of Lemma 55.3.

Problem 41. Using Problem 40, show that if a continuous map $f: S^n \rightarrow X$ is nulhomotopic, then $f = F|_{S^n}$ for some map $F: B^{n+1} \rightarrow X$. *Hint:* Again, look at the proof of Lemma 55.3.

Problem 42. Assume the fact that there is no retraction from B^{n+1} onto S^n , in the sense of Problem 39. Using Problem 41, show that the identity map of S^n cannot be nulhomotopic.

Problem 43. Suppose that $f: X \rightarrow X$ is continuous. Show that:

- (1) If X is the closed interval $[0, 1]$, then f has some fixed point. *Hint:* The Intermediate Value Theorem.
- (2) If X is the half-closed interval $[0, 1)$, then f need not have fixed points.

3. FREE GROUPS AND FREE PRODUCTS

Problem 44. Recall that for any set X , we write F_X to denote the *free group* generated by X . Show that X and F_X are never in bijection. *Hint:* Imitate Cantor's diagonalization argument.

Problem 45. Show that the following conditions on a subset S of a group G are equivalent:

- (1) The only subgroup of G containing S is G itself.
- (2) The homomorphism from F_S into G induced by the inclusion map from S into G is surjective.

In this case, we say that S *generates* G , or that S is a *generating set* for G . If $S = \{s_1, s_2, \dots\}$, then we also say that $s_1, s_2, \dots$ *generate* G .

Problem 46. Let a, m be arbitrary positive integers. Recall that “the *residue class* of $a \bmod m$ ” refers to the coset $a + m\mathbf{Z}$.

- (1) For which a and m does $a \bmod m$ generate $\mathbf{Z}/m\mathbf{Z}$?
- (2) In general, describe the subgroup generated by $a \bmod m$.

We say that a group is *cyclic* when it can be generated by a single element.

Problem 47. For any positive integer m , let

$$U_m = \{a \bmod m \mid ab \equiv 1 \bmod m \text{ for some } b\} \subseteq \mathbf{Z}/m\mathbf{Z}.$$

- (1) Show that U_m forms a group under the operation of multiplication mod m .
- (2) Compute U_2, U_3, U_4, U_5, U_6 .
- (3) Which of the groups in (2) are trivial? Which are cyclic?

Problem 48. Which subsets of $\mathbf{Z}$ are generating sets for $\mathbf{Z}$ as a group? *Hint:* Look up Bézout's Lemma.

Problem 49. Look up the definition of the *Cayley graph* of a group G equipped with a generating set S . Draw the Cayley graph in the case where:

- (1) $G = \mathbf{Z}$ (under addition) and $S = \{1\}$.
- (2) $G = \mathbf{Z}$ and $S = \{2, 3\}$.
- (3) $G = \mathbf{Z}/5\mathbf{Z}$ and $S = \{2\}$.
- (4) $G = F_S$ and $S = \{a, b, c\}$.

Problem 50. Let $(a, b), (c, d) \in \mathbf{Z}^2$ and

$$M = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

- (1) Show that M^{-1} exists and has only integer entries if and only if $\det M = \pm 1$.
- (2) Deduce from (1) that $(1, 0)$ and $(0, 1)$ are integer linear combinations of (a, b) and (c, d) if and only if $\det M = \pm 1$.
- (3) Deduce from (2) that $\{(a, b), (c, d)\}$ generates $\mathbf{Z}^2$ if and only if $\det M = \pm 1$.

Problem 51. Keep the setup of Problem 50. If $\det M = 0$, then what can you say about the subgroup of $\mathbf{Z}^2$ generated by (a, b) and (c, d) ?

Problem 52. Let a, b be arbitrary positive integers. Show that $\langle s \mid s^a, s^b \rangle$ is trivial if and only if a and b are relatively prime. *Hint:* Recall how we showed that $\langle s \mid s^b \rangle \simeq \mathbf{Z}/b\mathbf{Z}$.

Problem 53. Recall that the *order* of an element $g \in G$ is the smallest positive integer m such that $g^m = e$, if it exists, and infinity otherwise.

Show that if G is abelian, then its elements of finite order form a subgroup. It is called the *torsion subgroup* of G . If it is trivial, then we say that G is *torsion-free*.

Problem 54. Let G be a group in which $g^2 = e$ for all $g \in G$. Show that G is abelian. (Note that in this case, G is equal to its torsion subgroup.)

Problem 55. Let G be an arbitrary group. Show that:

- (1) As a set, G is in bijection with the set of homomorphisms from $\mathbf{Z}$ into G .
Hint: $\mathbf{Z}$ is isomorphic to a free group on one element.
- (2) The set $\{g \in G \mid g^2 = e\}$ is in bijection with the set of homomorphisms from $\mathbf{Z}/2\mathbf{Z}$ into G .

Note: If G is not abelian, then $\{g \in G \mid g^2 = e\}$ need not be a subgroup of G !

In Problems 56 and 57 below, let G_1, G_2 be groups with presentations

$$G_i = \langle s \in S_i \mid r \in R_i \rangle \quad \text{for } i = 1, 2.$$

Recall that their *free product* $G_1 * G_2$ has the presentation

$$G_1 * G_2 = \langle s \in S_1 \sqcup S_2 \mid r \in R_1 \sqcup R_2 \rangle.$$

You may assume the following fact: If $w \in G_1 * G_2$ is a reduced word that involves elements of both G_1 and G_2 , then $w \neq e$. For a reminder on the definition of a *reduced word*, see Munkres 415–416.

Problem 56. (1) Give a surjective homomorphism from $G_1 * G_2$ onto $G_1 \times G_2$.

Hint: In class, we discussed the case where $G_1, G_2 \simeq \mathbf{Z}$.

- (2) Show that if G_1 and G_2 are nontrivial, then the homomorphism in (1) is not injective.

Problem 57. For all $g \in G_1 * G_2$, we define the *length* of g to be its length as a reduced word in the elements of G_1 and G_2 .

- (1) Show that if x has even length at least 2, then it cannot have finite order.
- (2) Show that if x has odd length at least 3, then it is conjugate to some element of shorter length. Here, recall that the *conjugates* of x are the elements gxg^{-1} , where $g \in G$.
- (3) Deduce that any element of G of finite order is conjugate to an element of G_1 or G_2 of finite order.

Problem 58. Suppose that $X = X_1 \cup X_2$, where X_1, X_2 are closed, path-connected, and intersect in a single point p . (Recall that this is an example of a *wedge sum*.) Suppose that for $i = 1, 2$, the subspace X_i contains an open neighborhood of p , say W_i , such that $\{p\}$ is a deformation retract of W_i . Use Seifert–Van Kampen to show that in this situation,

$$\pi_1(X, p) \simeq \pi_1(X_1, p) * \pi_1(X_2, p),$$

Hint: In class, we discussed the case where $X_1, X_2 \simeq S^1$.

Hint: Set $U_1 = X_1 \cup W_2$ and $U_2 = X_2 \cup W_1$. Check that X_i is homotopy equivalent to U_i for $i = 1, 2$, and that $\{p\}$ is homotopy equivalent to $U_1 \cap U_2$. Then check the hypotheses needed for Seifert–Van Kampen. You may assume without proof that path-connectedness is preserved by homotopy equivalence.

Problems 59 and 60 below discuss the *real projective plane* P^2 from Munkres §60, page 372, and the *Klein bottle* K from Munkres §75, page 454.

Problem 59. Let X_1, X_2 be two copies of P^2 . Let X be the quotient space formed from $X_1 \sqcup X_2$ by excising small open disks from X_1 and X_2 , then gluing along the boundaries of the resulting holes.

- (1) Show that X is homeomorphic to K , using the descriptions of P^2 and K as quotients of $[0, 1]^2$.
- (2) Compute $\pi_1(X, x)$ by applying Seifert–van Kampen to open neighborhoods of X_1 and X_2 within X . Do not simply cite the formula for the fundamental group of a Klein bottle.

Problem 60. (1) Find a 2-fold covering $p: S^1 \times S^1 \rightarrow K$.

- (2) Describe the homomorphism of fundamental groups induced by p .

EXAM 3 TOPICS

Exam 3 will be an **open-book** exam. It will be held in-class, 12:40–2:00 pm, on ~~April 13~~ **April 17**.

“Open book” means: You will be allowed to look at Munkres, *Topology*, 2nd Ed., and any notes on paper that you wrote prior to the exam, but you will not be allowed to use electronic devices, including phones, computers, tablets, *etc.*

Topics Included.

- Sections, retractions, and their properties (Problem 39)
- Why there is no retraction from D^2 onto its subspace S^1
- Equivalent ways to describe when a map $S^1 \rightarrow X$ extends to a map $D^2 \rightarrow X$ (Munkres Lemma 55.3)
- Statement of the fixed-point theorem for D^2
- Statement of the Jordan separation theorem (for S^2 or $\mathbf{R}^2$)
- Examples of normal subgroups, homomorphisms, kernels, quotient groups
- Definition of the free group F_S on a set S , in terms of words
- How a map from a set S into a group G defines a homomorphism from F_S into G
- Examples and non-examples of generating sets for groups that we discussed, like $\mathbf{Z}$ or $\mathbf{Z}/m\mathbf{Z}$ or $\mathbf{Z}^2$
- Meaning of a group presentation $\langle s \in S \mid r \in R \rangle$, where $R \subseteq F_S$
- Statement of Seifert–van Kampen in terms of group presentations
- Presentations for the fundamental groups of the torus, Klein bottle, real projective plane, m -fold dunce cap, and how to describe elements in these groups using the presentations

Topics Not Included.

- The proofs of the Brouwer fixed-point theorem, Jordan separation theorem, or Seifert–van Kampen theorem
- The Jordan curve theorem
- Listing all possible generating sets of a given group
- Computing the precise smallest normal subgroup of F_S containing a given set R

4. SIMPLICIAL COMPLEXES AND SURFACES

Recall that if S is a finite set of points in general position in some real vector space, then we write $\Delta(S)$ for the simplex formed by its convex hull.

Problem 61. In $\mathbf{R}^3$, let $e_1 = (1, 0, 0)$ and $e_2 = (0, 1, 0)$ and $e_3 = (0, 0, 1)$.

- (1) Determine every possible simplicial complex $\{\Delta(S)\}_S$ in which each set S is a subset of $\{e_1, e_2, e_3\}$. There should be 12, including the empty simplicial complex.
- (2) Which simplicial complexes in (1) have a geometric realization that is itself a simplex?

Problem 62. For each space X below, draw or describe a geometric realization of a simplicial complex that is homeomorphic (not just homotopy equivalent) to X .

- (1) S^1
- (2) $S^1 \vee S^1 \vee S^1$
- (3) D^2 (the set $\{(x, y) \in \mathbf{R}^2 \mid x^2 + y^2 \leq 1\}$)
- (4) $S^1 \times [0, 1]$
- (5) S^2
- (6) $S^1 \times S^1$
- (7) $D^2 \times S^1$
- (8) $\{(x, y, z) \in \mathbf{R}^3 \mid x^2 + y^2 + z^2 \leq 1\}$

Hint: To handle $S^1 \times S^1$ and $D^2 \times S^1$, first describe them as quotient spaces of $[0, 1] \times [0, 1]$ and $D^2 \times [0, 1]$, respectively.

(Recall that a *triangulation* of X is a homeomorphism from X onto such a geometric realization. Each space above has a triangulation.)

Problem 63. For each space in Problem 62, compute its Euler characteristic using the triangulation you found.

Problem 64. Let $I = [0, 1]$. The (closed) *Möbius band* is the quotient space

$$M = (I \times I) / \sim,$$

where $(0, y) \sim (1, 1 - y)$ for all y , and no other pairs of distinct points of $I \times I$ get identified under $\sim$.

Recall that the circle S^1 is homeomorphic to a similar quotient space. Give an explicit homotopy equivalence between M and S^1 . (Hence, they have the same fundamental groups.)

Problem 65. Keep the setup of Problem 64.

- (1) What space do you get if you cut M along the image of the line segment in $I \times I$ between $(0, \frac{1}{2})$ and $(1, \frac{1}{2})$?
- (2) What space do you get if you glue two copies of M along their boundary?
Hint: Work with $I \times I$, keeping track of equivalence relations. The way that the Möbius boundaries are oriented relative to each other will not matter.

Problem 66. Keep the setup of Problems 64 and 65.

- (1) Find triangulations of $I \times I$ and M such that the quotient map from $I \times I$ onto M takes simplices to simplices.
- (2) Using the triangulations in (1), verify directly that the Euler characteristic of $I \times I$ is 1, while that of M is 0. Why are these values consistent with Problems 63 and 64?

Problem 67. Suppose that $\mathcal{K}$ is a simplicial complex whose geometric realization $|\mathcal{K}|$ is homeomorphic to $[0, 1]$. Show that there is a finite list of points

$$0 = a_0 < a_1 < \cdots < a_k = 1$$

such that under $|\mathcal{K}| \simeq [0, 1]$, the 1-simplices in $\mathcal{K}$ correspond to the subintervals of the form $[a_i, a_{i+1}]$.

Problem 68. Use a triangulation to compute the mod 2 simplicial homology of the closed interval $[0, 1]$. Using Problem 67, deduce that the answer is the same regardless of the triangulation.

Problem 69. Using the triangulations you found in Problem 62, compute the mod 2 simplicial homology of:

- (1) S^1 .
- (2) D^2 .
- (3) S^2 .

Problem 70. Recall the Klein bottle K from Munkres §75, page 454.

- (1) Find a triangulation of K . *Hint:* As with $S^1 \times S^1$, it helps to first describe K as a quotient space of $[0, 1] \times [0, 1]$.
- (2) *Hard:* Compute the mod 2 simplicial homology of K and $S^1 \times S^1$. Are they the same or different?
- (3) Read in Crossley §9.3 the integral simplicial homology of K and $S^1 \times S^1$. Are they the same or different?