

Nephron Nursing Coaching Center, Rajshahi
Comprehensive/Licensing Exam-2026
Course Name: Diploma in Midwifery

Time: 1:30 Hours

Full Marks: 100

Multiple Choice Questions (MCQ)

Direction: Fill up the "T" for true and "F" for false to left side of each option.

- 1. Depression is characterized by-**
 - a. Suicidal tendency
 - b. Impaired concentration
 - c. Social support
 - d. Hormonal factors
 - e. Insomnia
- 2. Common maternal mental health disorders-**
 - a. Depression and anxiety
 - b. Disharmony
 - c. Postpartum depression
 - d. Loss of appetite
 - e. Lack of support and family conflict
- 3. Factors of gender-based violence, except-**
 - a. Unemployment
 - b. Male dependence
 - c. Economic dependence
 - d. Child marriage
 - e. Dowry practices
- 4. Following are Sexual violence in Bangladesh-**
 - a. Intimate partner violence
 - b. Digital Sexual violence
 - c. Social security
 - d. Rape and gang rape
 - e. Physical assault
- 5. Drugs are safe to use during pregnancy -**
 - a. Iron and folate
 - b. Mebendazole
 - c. Aspirin
 - d. Multivitamin
 - e. Tetracycline
- 6. Anti-emetic drugs can help to --**
 - a. Induce vomiting
 - b. Relieve vomiting
 - c. Induce nausea
 - d. Relieve nausea
 - e. Reduce motion sickness
- 7. Common physiological changes during pregnancy, except-**
 - a. Enlarged breast
 - b. Big uterine enlargement
 - c. Increase height of the mother
 - d. Perineal edematous and more vascular
 - e. Weight gain
- 8. Danger signs in pregnancy, includes-**
 - a. Nausea and vomiting
 - b. Bleeding per vagina
 - c. Constipation
 - d. Severe abdominal pain
 - e. Severe headache and blurring of vision
- 9. Birth panning included**
 - a. Decided birthing place
 - b. Support person
 - c. Foods and dinks
 - d. Necessary drugs
 - e. Transportation facilities
- 10. Criteria of normal labor includes -**
 - a. Multiple fetus
 - b. Completed 37 weeks
 - c. Spontaneous onset
 - d. Transvers lie
 - e. Cephalic presentation
- 11. Pillars of safe motherhood includes -**
 - a. Pre-conception care
 - b. Cervical screening
 - c. Intrapartum care
 - d. Post abortion care
 - e. Vaccination
- 12. Function of placenta includes -**
 - a. Maintain fetal position
 - b. Supply oxygen
 - c. Maintain temperature
 - d. Excrete west product
 - e. Supply nutrition
- 13. Risk factors uterine rupture includes-**
 - a. Lithotomy position
 - b. Excess use of oxytocin
 - c. Exercise
 - d. Over distention of uterus
 - e. H/O previous uterine surgery
- 14. Complication of the vagina delivery -**
 - a. Water breakdown
 - b. Vaginal bleeding
 - c. Perineal tears
 - d. Prolong labor
 - e. Confusion
- 15. Pillars of midwifery are -**
 - a. Professionalism
 - b. Behavior
 - c. Language
 - d. Knowledge
 - e. Partnership
- 16. National Professional midwifery issues are -**
 - a. Midwifery education
 - b. Hospital management
 - c. Midwifery services
 - d. Achievement of SDG
 - e. Community services
- 17. Components of ICM includes -**
 - a. Education
 - b. Regulation
 - c. Collaboration
 - d. Decision
 - e. Association
- 18. Leadership syres includes -**
 - a. Democratic
 - b. Autocratic
 - c. Empathetic
 - d. Charismatic
 - e. Instructive
- 19. Problems of modern family are-**

- a. Lack of trust
- b. Increase moral
- c. Instability
- d. Increase family members
- e. Sexual mis conduct

20. Climate change effects in pregnancy are-

- a. Dehydration
- b. Miscarriage
- c. Preterm delivery
- d. Weight loses
- e. Low birth weight baby

21. Ethical principles are-

- a. Autonomy
- b. Behaviors
- c. Malfeasance
- d. Manner
- e. Justice

22. Loss and grief related factors are -

- a. Loss a Loved one
- b. Severe headache
- c. Relationship breakdown
- d. Gender inequality
- c. Child leaving home

23. Conflict resolution strategies are –

- a. Avoiding
- b. Fighting
- c. Collaborating
- d. Compromising
- e. Sharing

24. Complication of septic abortion includes-

- a. Infertility
- b. Septic shock
- c. Renal failure
- d. Irregular menstruation
- e. Chronic pelvic pain

25. Sexual transmitted diseases are-

- a. Malaria
- b. AIDS
- c. Skin infections
- d. Syphilis
- e. Gonorrhea

26. Step of active management of 3rd stage of labor includes-

- a. Lithotomy position
- b. Inj. Oxytocin 10 IU IM
- c. Fundal pressure
- d. CCT with CT
- e. Abdominal palpation

27. Characteristic of hypoglycemia includes

- a. Irregular heart rate
- b. Hot flashes
- c. Shakiness and trembling
- d. Pain
- e. Sweating, chills and Clamminess

28. Phases of ovulation includes -

- a. Proliferative Phase
- b. Follicular Phase
- c. Secretory Phase
- d. Luteal Phase
- e. Ovulation Phase

29. Factors related for growth and development of an embryo includes-

- a. Genetic factors
- b. Social factors
- c. Nutritional factors

- d. Socio-economic factors
- e. Hormonal factors

30. EPI vaccination includes-

- a. Vaccine of Rota virus
- b. Vaccine of Polio
- c. Vaccine of corona
- d. Vaccine of Diphtheria
- e. Vaccine of Tetanus

31. Features of missed abortion includes-

- a. Disappearance of pregnancy symptoms
- b. Closed cervical OS
- c. Raise of temperature
- d. Passage of mass through vagina
- e. Absence of fetal movement

32. P/V examination helps to identify -

- a. Uterine contraction
- b. Identifying the presenting part
- c. Dilatation and effacement of cervix
- d. Fetal movement
- e. Color of the Liquor

33. Drugs used for control of hypertension –

- a. Hydralazine
- b. Labetalol
- c. Nifedipine
- d. Methyldopa
- e. Diazepam

34. Preventive measures of pre-eclampsia –

- a. Regular ANC
- b. Early detection of PIH
- c. Control of hypertension
- d. Adequate rest
- e. No antenatal care

35. Nursing care in pre-eclampsia –

- a. Monitor BP
- b. Monitor urine output
- c. Administer antihypertensive drugs
- d. Provide emotional support
- e. Ignore patient

36. Differential diagnosis of pre-eclampsia –

- a. Chronic hypertension
- b. Renal disease
- c. Epilepsy
- d. Gestational hypertension
- e. Normal pregnancy

37. Signs of mild pre-eclampsia –

- a. BP $\geq$ 140/90 mmHg
- b. Proteinuria trace to +1
- c. Oedema mild
- d. Convulsion
- e. Normal BP

38. Signs of moderate pre-eclampsia –

- a. BP $\geq$ 150/100 mmHg
- b. Proteinuria +2
- c. Oedema moderate
- d. Severe headache
- e. Normal BP

39. Signs of severe pre-eclampsia –

- a. BP $\geq$ 160/110 mmHg
- b. Proteinuria $\geq$ 3+
- c. Oliguria
- d. Cerebral symptoms
- e. Normal pregnancy

40. Management of mild pre-eclampsia –

- a. Rest
- b. Monitor BP

- c. Monitor urine protein
 - d. Regular ANC
 - e. Immediate termination
- 41. Maternal complications –**
- a. Septicemia
 - b. Shock
 - c. Death
 - d. Infertility
 - e. Recurrent Pregnancy
- 42. Drugs used in management of sepsis –**
- a. Broad-spectrum antibiotics
 - b. IV fluids
 - c. Analgesics
 - d. Antipyretics
 - e. Diazepam
- 43. Causes of puerperal sepsis –**
- a. Retained placenta
 - b. Genital tract trauma
 - c. PROM
 - d. Prolonged labour
 - e. Normal delivery
- 44. Signs of subinvolution of uterus –**
- a. Uterus not contracted
 - b. Persistent lochia
 - c. Enlarged uterus
 - d. Foul-smelling discharge
 - e. Normal involution
- 45. Midwifery Care for sepsis includes –**
- a. Monitor vital signs
 - b. Administer antibiotics
 - c. Maintain hygiene
 - d. Provide emotional support
 - e. Advise for family planning
- 46. Role of midwife –**
- a. Early detection
 - b. Prevent infection
 - c. Referral to hospital
 - d. Health education
 - e. Prescribe antibiotics
- 47. Differential diagnosis of Puerperal Sepsis –**
- a. Malaria
 - b. Typhoid fever
 - c. UTI
 - d. Appendicitis
 - e. Eclampsia
- 48. Signs of septic shock –**
- a. Hypotension
 - b. Tachycardia
 - c. Cold extremities
 - d. Oliguria
 - e. Bradycardia
- 49. Preventive measures –**
- a. Aseptic delivery
 - b. Proper ANC
 - c. Correction of anemia
 - d. Skilled birth attendant
 - e. Delay management
- 50. Investigations –**
- a. CBC
 - b. Blood culture
 - c. Urine culture
 - d. Swab culture
 - e. Chest X-ray
- 51. An assessment of a baby just born finds: heart rate of 110/min, the baby is crying, her arms and legs slow flexed, respiratory rate is <30/min, and the hands are slightly blue. APGAR score is:**
- a. 6
 - b. 9
 - c. 7
 - d. 8

- 52. The meconium stain liquor is the sign of**
- a. Prolong labor
 - b. Fetal distress
 - c. Still birth
 - d. Shoulder presentation
- 53. The purpose assessing a pregnant woman for quality nutrition is to support:-**
- a. Treating PPH
 - b. Optimal fetal growth
 - c. Development of obesity
 - d. Malaria treatment
- 54. Maternal anemia effects in pregnancy**
- a. Miscarriages
 - b. Low birth weight baby
 - c. Fundal massage
 - d. Preterm delivery
- 55. Initial management of newborn includes all of the below except:-**
- a. Delayed cord clamping
 - b. Drying of the baby
 - c. Delayed breast feeding
 - d. Mother and baby skin to skin
- 56. Common congenital heart defect is:**
- a. Congenital heart disease
 - b. Pulmonary stenosis
 - c. Arterial septal defect (ASD)
 - d. Down syndrome.
- 57. Hormone secreted from ovaries include -**
- a. Testosterone
 - b. Oestrogen
 - c. Oxytocin
 - d. Insulin
- 58. Principles of drug administration are all of these except-**
- a. Right patient
 - b. Right drugs
 - c. Right choice
 - d. Right dose
- 59. A 37y old women G7 P6 presents at term in normal active labor. What is one thing you may be concerned about? -**
- a. Incidence of abortion
 - b. Malpresentation
 - c. More anxiety for childbirth
 - d. Congenital abnormalities
- 60. Most important antenatal advice should be provided in each visit –**
- a. Personal hygiene
 - b. Dietary advice
 - c. Birth planning
 - d. Family planning
- 61. Prior to doing a vacuum assisted vaginal birth it is recommended that:**
- a. The bladder be emptied
 - b. Contractions must be absent
 - c. The mother is starting second Stage
 - d. Membranes must be intact
- 62. A woman presents to the hospital in active labor. She has a history of PPH. What should be done to help prevent PPH with this birth?**
- a. Pitocin augmentation of labor
 - b. Give blood to prevent anemia
 - c. Active management of 3rd stage labor
 - d. A cesarean section
- 63. Management of cord prolapse includes everything below except:-**
- a. Prepare for emergency delivery
 - b. Relieve pressure on cord
 - c. Try to re-insert the cord
 - d. Assess fetal heart rate every 5 minutes interval
- 64. When the membrane rupture 18 or more hours before the onset of labor what condition may develop?**
- a. Peritonitis
 - b. Chorioamnionitis
 - c. Appendicitis
 - d. Pelvic abscess
- 65. Lower abdominal pain in early pregnancy can be caused by all of the following except:**
- a. Infections -STI or urinary tract Infection
 - b. Life threatening ectopic pregnancy
 - c. False labor pain
 - d. Placental abruption
- 66. Postnatal psychosis -**
- a. Pre-conception
 - b. Postpartum period
 - c. Antenatal period
 - d. During delivery

67. A 24-year-old G2 P1 at 9 weeks admitted Ganey ward H/o P/V bleeding, passage of tissue in her vagina and lower abdominal pain. The most likely diagnosis is:

- a. Threatened abortion b. incomplete abortion
- c. Multiple pregnancy d. ectopic pregnancy

68. Important investigation for detection of cervical cancer -

- a. Urine for R/E b. Urine for C/S
- c. Per vaginal examination d. VIA test

69. Which Hormone helps in breast milk ejection is -

- a. Oxytocin b. Prolactin
- c. Estrogen d. Progesterone

70. Vaginal examination is contraindicated in pregnancy during which situation?

- a. Carcinoma of cervix b. prolapsed cord
- c. placenta previa d. Gonorrhoea

71. After the membranes rupture what is the priority assessment for the midwife

- a. presentation b. Fetal heart rate
- c. Blood pressure d. Pelvic examination

72. A main cause of infertility is-

- a. Pelvic Inflammatory Disease (PID)
- b. Sexual Transmitted Infections (STIs)
- c. Urinary Tract Infection (UTI)
- d. Genetic

73. Inflammation of the breast as a result of a infection is called-

- a. Mastitis b. Shalpingitis
- c. Engorged breast d. Abcess

74. A primipara mother presents one week after birth, She is tearful, has crying spells, and lack of appetite. Her most likely diagnosis is:

- a. Anemia b. post natal depression
- c. schizophrenia d. Sepsis

75. The condition in which good uterine contraction are present but there is no decent of the presenting part of the fetus, is known as?

- a. Precipitate labour b. Prolonged labour
- c. Obstructed Labour d. Preterm Labour

76. A 25-year-old G2 P2 presents after delivering at home with heavy vaginal bleeding. An initial evaluation diagnoses her with uterine atony. What is the best medication to use:

- a. Salbutamol b. Hydralazine
- c. Magnesium sulphate d. Oxytocin

77. The placenta should be delivered normally within the minutes after the delivery of the baby

- a. 5 minutes b. 45 minutes
- c. 30 minutes d. 60 minutes

78. Which of the following is a primary concern if the membranes rupture more than 24 hours before delivery?

- a. Infection b. Stillbirth
- c. Cerebral palsy d. Respiratory distress

79. A 20-year age primiparas women has admitted in labor ward with 35 weeks pregnancy with history of blurred of vision, decrease urine output, generalized oedema. BP 180/105mm of Hg. What is your probable diagnosis?

- a. Gestational hypertension b. Pre- Eclampsia
- c. Severe Pre- Eclampsia d. Eclampsia

80. What are some of the symptoms associated with trauma to the perineum at birth from obstructed labor? (circle all that apply)

- a. Fecal incontinence b. Perineal pain
- c. Pain with sex d. Cystocele

81. Commonly used non-pharmacological method for comfort measures

- a. Hot compression b. Hot drink
- c. Back massage d. Bed rest

82. Safe drugs during pregnancy is -

- a. Tetracycline b. Oxytocin
- c. Methyldopa d. Ergometrin

83. Vaccine to be avoided in pregnancy is -

- a. MMR b. TT
- c. Influenza d. Hepatitis B

84. Drug recommended in hypertension -

- a. Aspirine b. Morphine
- c. Pethedine d. Amlodipine

85. Regulatory body of Midwifery is-

- a. DGHS b. DGNM
- c. BNMC d. UNFPA

86. BCG is a

- a. Killed vaccine b. Toxoid
- c. Live vaccine d. Combined vaccine

87. Spore can be destroyed by -

- a. Boiling b. Autoclaving
- c. Steaming d. Disinfectant

88. Which is not an importance of colostrum-

- a. Rich in high protein b. Rich in anti-infective factors
- c. Rich in ig A d. All above

89. Sign of the 3rd stage of labour-

- a. A gush of blood b. Crowning
- c. Soft Uterus d. Show

90. Universal causes of maternal death...

- a. Haemorrhage b. Eclampsia
- c. Puerperal sepsis d. All above

91. when we call it as latent phase....

- a. 0 to 3 cm. b. 4 to 6 cm.
- c. 8 to 10 cm. d. 4 to 7 cm.

92. First stage of labour-

- a. Full dilation of cervix to Baby delivered
- b. True labour contractions to full dilation of the cervix
- c. Baby delivered to expulsion of placenta
- d. Expulsion of placenta

93. Neonatal danger sign-

- a. History of convulsion b. feeding well.
- c. Normal temperature d. Movement well.

94. Exclusive Breastfeeding Means....

- a. Up to Six months b. First 5 months of life
- c. Breastfed with additional food. d. First weeks of life.

95. Neonatal normal Birth weight

- a. 2000-4000 gm b. 2500-5000 gm
- c. 2500-4000 gm d. 2600-4000 gm.

96. What is called 5th vital sign?

- a. Mean arterial pulse b. ABG analysis
- c. Oxygen saturation d. Pain

97. For taking oral temperature after drinking anything cold, health care provider should wait -

- a. 5-10 minutes b. 10-20 minutes
- c. 20-30 minutes d. 30-40 minutes

98. Electrocardiograph is used to measure-

- a. Blood count b. Heartbeat
- c. Electricity d. Temperature

99. CPR for a heart attack patient-

- a. 2 times/sec b. 3 times/sec
- c. 6 times/sec d. 10 times/sec

100. What can be helpful for a road accident?

- a. First aid b. hospitalization
- c. Legal aid. d. No attempt