

# Universal Specificity Investigation 6: Inducing the Cause of Total Time Dilation and Total Energy's Relation to Potential Energy

Daniel Harris  
Northrop Grumman  
Morrisville, USA  
daniel.harris2@ngc.com

Prior investigations into the theory of universal specificity found a proper conception of time missed in common practice; which led to the realization that a universally stationary frame must exist; which led to the discovery that the average effective speed of light,  $c_0$ , is identical in all directions for any inertial reference frame, and is a function of velocity  $v$ , as shown in Equation (1); which led to discovering the cause of kinetic time dilation, shown Equation (2); which led to revisiting the relativistic kinetic energy and total energy model, shown in Equation (3); which finally led to discovering the cause of gravitational time dilation, also shown Equation (2).

$$c_0 = c\sqrt{1 - \frac{v^2}{c^2}} \quad (1)$$

$$\frac{dt}{dt'} = \sqrt{1 - \frac{w}{e_T}} \quad (2a)$$

$$\frac{dt}{dt'} = \sqrt{1 - \frac{\Delta e_K}{e_T}} \quad (2b)$$

$$\frac{dt}{dt'} = \sqrt{1 - \frac{\Delta e_P}{e_T}} \quad (2c)$$

$$\Delta E_K = \frac{1}{2}mv^2 \quad (3a)$$

$$E_T = \frac{1}{2}mc^2 = \frac{1}{2}mc_0^2 + \frac{1}{2}mv^2 \quad (3b)$$

## 1. TIME DILATION EQUIVALENCE

I now relate changes in ITDs to changes in specific total external energy, as in ignoring internal energy,  $\frac{1}{2}mc_0^2$ , and then I will update the total energy model to include potential energy.

Changes in specific kinetic or specific potential energy are related to changes in ITDs, but this is only half of the picture because we tacitly assumed all else remained equal. Now we test what if all else does not remain equal to discover a more precise cause to changes in ITDs.

In reviewing Equation (2), simple analysis reveals that transferring some amount of specific kinetic energy to some amount of specific potential energy (or vice versa) would not

cause an overall change in the ITD. As an example, consider an object with some amount of specific potential energy that then enters a state with equal specific kinetic energy, but without the potential. Equation (4) shows that the two ITDs in each state are equivalent.

Let  $\Delta e_P > 0$ .

$$\text{Let } \frac{1}{\gamma} = \frac{dt}{dt'} \quad (4a)$$

$$\frac{1}{\gamma_P^2} = 1 - \frac{\Delta e_P}{e_T} \quad (4b)$$

$$1 - \frac{1}{\gamma_P^2} = \frac{\Delta e_P}{e_T} \quad (4c)$$

$$\left(1 - \frac{1}{\gamma_P^2}\right)e_T = \Delta e_P \quad (4d)$$

$$\Delta e_P \implies \Delta e_K \quad (4e)$$

$$\left(1 - \frac{1}{\gamma_P^2}\right)e_T = \Delta e_K \quad (4f)$$

$$1 - \frac{1}{\gamma_P^2} = \frac{\Delta e_K}{e_T} \quad (4g)$$

$$\frac{1}{\gamma_P^2} = 1 - \frac{\Delta e_K}{e_T} \quad (4h)$$

$$\frac{1}{\gamma_P^2} = \frac{1}{\gamma_K^2} \blacksquare \quad (4i)$$

Invoking the method of agreement: observing that changes in specific potential energy and changes in specific kinetic energy occurred, while no changes in ITD occurred, proves inductively that they are not the fundamental causes to changes in ITDs—they each play half a role.

The same change in specific total external energy,  $\Delta e_t$ , in each state caused the same change in ITDs. This proves inductively, via method of agreement, that changes in ITD are caused by,  $\Delta e_t$ , and vice versa.

## 2. THE CAUSE OF TOTAL TIME DILATION

I begin this causal derivation by trying to solve the ITD for an object stationary within a gravitational field. Then I proceed to determine how a change in kinetic energy, as measured from the stationary position within the gravitational field, affects the ITD. This situation is depicted in Figure 1.




Figure 1. Total effective time differential example.

Ultimately, the desire is to measure total effective ITD,  $\frac{dt_2}{dt'}$ , for the moving object within the gravitational field.  $\frac{dt_1}{dt'}$  and  $\frac{dt_2}{dt_1}$  can be measured, and both of these ITDs are relatable to the total effective ITD via the chain rule, as shown in Equation (5).

$$\frac{dt_1}{dt'} = \sqrt{1 - \frac{\Delta e_{P1}}{e_T}} \quad (5a)$$

$$\frac{dt_2}{dt_1} = \sqrt{1 - \frac{\Delta e_{K2/P1}}{e_T}} \quad (5b)$$

$$\frac{dt_2}{dt'} = \frac{dt_2}{dt_1} \frac{dt_1}{dt'} = \sqrt{1 - \frac{\Delta e_{K2/P1}}{e_T}} \sqrt{1 - \frac{\Delta e_{P1}}{e_T}} \quad (5c)$$

This can be generalized to any condition where an object within a gravity potential gains some kinetic energy as measured inside that potential, as shown in Equation (6).

$$\text{Let : } \frac{1}{\gamma_P} = \frac{dt_P}{dt'} = \sqrt{1 - \frac{\Delta e_P}{e_T}} \quad (6a)$$

$$\text{Let : } \frac{1}{\gamma_{K/P}} = \frac{dt_K}{dt_P} = \sqrt{1 - \frac{\Delta e_{K/P}}{e_T}} \quad (6b)$$

$$\frac{1}{\gamma_T} = \frac{1}{\gamma_{K/P}} \frac{1}{\gamma_P} = \sqrt{1 - \frac{\Delta e_{K/P}}{e_T}} \sqrt{1 - \frac{\Delta e_P}{e_T}} \quad (6c)$$

$$\frac{1}{\gamma_T} = \sqrt{1 - \frac{(1 - \frac{\Delta e_P}{e_T}) \Delta e_{K/P} + \Delta e_P}{e_T}} \quad (6d)$$

$$\frac{1}{\gamma_T} = \sqrt{1 - \frac{\frac{1}{\gamma_P^2} \Delta e_{K/P} + \Delta e_P}{e_T}} \quad (6e)$$

$$\frac{1}{\gamma_T} = \sqrt{1 - \frac{\Delta e_K + \Delta e_P}{e_T}} = \sqrt{1 - \frac{\Delta e_t}{e_T}} \blacksquare \quad (6f)$$

Of note, what the term,  $\frac{1}{\gamma_P^2} \Delta e_{K/P}$ , in Equation (6e) tells us is that object's speed slow down, by a factor of  $\frac{1}{\gamma_P}$ , due to gravitational time dilation.

### 3. UPDATING TOTAL ENERGY MODEL

I can now use these tools to update the total energy model in Equation (3) to include the potential energy term, along with the other terms—internal energy and kinetic energy—as shown in Equation (7).

$$\frac{1}{\gamma_T} = \sqrt{1 - \frac{\Delta e_t}{e_T}} \quad (7a)$$

$$\frac{1}{\gamma_T^2} = 1 - \frac{\Delta e_t}{e_T} \quad (7b)$$

$$e_T = \frac{1}{\gamma_T^2} e_T + \Delta e_t = e_I + \Delta e_K + \Delta e_P \quad (7c)$$

$$E_T = E_I + \Delta E_K + \Delta E_P \quad (7d)$$

$$\frac{1}{2} mc^2 = \frac{1}{2} mc_0^2 + \frac{1}{\gamma_P^2} \frac{1}{2} mv^2 + m \int g(r) dr \quad (7e)$$

$$\frac{1}{2} mc^2 = \frac{1}{2} mc_0^2 + \frac{1}{2} mv_0^2 + m \int g(r) dr \blacksquare \quad (7f)$$

Just as  $c_0$  is the calibrated speed of light for a given reference frame (see investigation 4)<sup>2</sup>, so  $v_0$  is the calibrated velocity of the reference frame (relative to the universally stationary frame, see Investigation 1, in empty space).

### 4. CONCLUSION

In conclusion, total time dilation was found to be ultimately caused by a change in specific total external energy, which led to two grand integrations: (1) kinetic and gravitational time dilation are the same phenomenon caused by the same thing,

<sup>2</sup>Note:  $c_0 = \frac{1}{\gamma_T} c$  instead of just  $c_0 = \frac{1}{\gamma_K} c$  as found in investigation 4.

namely changes in specific total external energy, and (2) total energy, to be complete, must include a potential energy term.

This investigative series into the theory of universal specificity has discovered enough that we can circle back to one of the original questions: is there a way to objectively determine the location of the USF? That is the focus of the next investigation.