

PATIENT: MUSTANSIR

AGE: 18 Years

CHIEF COMPLAINT & SYMPTOMS

neck ache (Duration: 1 week | Severity: Moderate)

DETAILED CLINICAL SUMMARY

Cervical strain occurs when the muscles or tendons in the neck are stretched or torn, often due to activities that require repetitive motion or awkward positioning. The pathophysiology involves micro-tears in the muscle fibers, leading to inflammation and pain. Common triggers include poor ergonomics, sudden jerking motions, or prolonged periods of poor posture. The natural progression of cervical strain typically involves acute pain that may subside with rest and conservative management. Treatment usually includes physical therapy, pain management with NSAIDs, and ergonomic adjustments. In severe cases, corticosteroid injections or surgical intervention may be considered if conservative measures fail.

Rx RECOMMENDED PHARMACOTHERAPY

Medication Name	Dosage	Frequency	Duration
Ibuprofen 400mg	1 tablet	Every 8 hours as needed for pain	5 days

ADVICE & LIFESTYLE MODIFICATIONS

- Apply a warm compress to the neck for 15-20 minutes several times a day to promote blood flow and reduce muscle stiffness.

- Engage
future stra

HABITS TO CHANGE / PREVENTIVE ADVICE

- Incorporate regular stretching and strengthening exercises into the daily routine to improve neck muscle resilience and posture, which can help prevent future injuries.

- Adjust w

POSSIBLE ALLERGIES / INTERACTIONS TO WATCH

- Be aware of potential allergens such as dust and pollen that may exacerbate respiratory symptoms, especially during outdoor activities. Monitor for any interactions with current medications, particularly if new treatments are introduced.

DR. AI CONSULTANT

Electronically Generated via MediScan AI 3D Core Engine

Disclaimer: This digital prescription is an AI-assisted synthesis based on user-reported symptoms. It is meant for informational support and does not replace regular doctor consultation, physical diagnostics, or clinical tests.