

PATIENT: MUSTANSIR

AGE: 18 Years

CHIEF COMPLAINT & SYMPTOMS

neck pain (Duration: 5 days | Severity: Moderate)

DETAILED CLINICAL SUMMARY

Cervical strain occurs when the neck muscles and ligaments are overstretched or torn, often due to sudden movements, poor posture, or repetitive strain. The pathophysiology involves micro-tears in the muscle fibers and connective tissues, leading to inflammation and pain. Common triggers include physical activities such as swimming, which may involve sudden neck movements. The condition typically progresses from acute pain to chronic discomfort if not managed properly. Management includes rest, physical therapy, and pain relief medications. In severe cases, further imaging may be required to rule out other conditions.

Rx RECOMMENDED PHARMACOTHERAPY

Medication Name	Dosage	Frequency	Duration
Ibuprofen 400mg	1 tablet	Every 8 hours as needed for pain	5 days

ADVICE & LIFESTYLE MODIFICATIONS

- Apply a cold compress to the neck for 15-20 minutes every few hours to reduce inflammation and numb the pain. This can help alleviate acute pain and swelling.

- Gentle neck stretches
avoid excessive head movements

HABITS TO CHANGE / PREVENTIVE ADVICE

- Incorporate regular stretching and strengthening exercises for the neck and upper back to improve muscle tone and prevent future strains. This is crucial for someone with a sedentary lifestyle.

- Maintain good posture
cervical spine alignment

POSSIBLE ALLERGIES / INTERACTIONS TO WATCH

- The patient has allergies to dust, pollen, and milk. It is important to monitor for any allergic reactions to medications prescribed, especially non-steroidal anti-inflammatory drugs (NSAIDs) like ibuprofen.

DR. AI CONSULTANT

Electronically Generated via MediScan AI 3D Core Engine

Disclaimer: This digital prescription is an AI-assisted synthesis based on user-reported symptoms. It is meant for informational support and does not replace regular doctor consultation, physical diagnostics, or clinical tests.