

PATIENT: MUSTANSIR

AGE: 18 Years

CHIEF COMPLAINT & SYMPTOMS

neck pain (Duration: 1 week | Severity: Severe)

DETAILED CLINICAL SUMMARY

Cervical strain occurs when the neck muscles or ligaments are overstretched or torn, often due to sudden movements, poor posture, or repetitive strain. This condition can lead to inflammation and pain in the neck region, which may radiate to the shoulders or upper back. The severity of symptoms can vary, with some patients experiencing acute pain that worsens with specific activities, such as swimming in this case. Management typically includes rest, ice application, physical therapy, and pain relief medications. In chronic cases, further evaluation may be necessary to rule out underlying conditions.

Rx RECOMMENDED PHARMACOTHERAPY

Medication Name	Dosage	Frequency	Duration
Poiz 30	As prescribed	As prescribed	As prescribed
Cyblex M80	As prescribed	As prescribed	As prescribed

ADVICE & LIFESTYLE MODIFICATIONS

- Apply ice packs to the neck for 15-20 minutes every hour to reduce inflammation and numb the area, which can help alleviate pain and swelling.

- Perform
recovery a

HABITS TO CHANGE / PREVENTIVE ADVICE

- Incorporate regular stretching and strengthening exercises into the daily routine to enhance neck muscle resilience and prevent future strains.

- Avoid ac
healing.

POSSIBLE ALLERGIES / INTERACTIONS TO WATCH

- The patient has allergies to dust, pollen, and milk, which should be considered when prescribing medications or recommending over-the-counter treatments to avoid potential allergic reactions.

DR. AI CONSULTANT

Electronically Generated via MediScan AI 3D Core Engine

Disclaimer: This digital prescription is an AI-assisted synthesis based on user-reported symptoms. It is meant for informational support and does not replace regular doctor consultation, physical diagnostics, or clinical tests.