

Patient: MUSTANSIR

Age: 18 yrs | Urgency: Moderate

CHIEF COMPLAINT

stomach pain , nausea , loose motions

Duration: 1 day | Severity: Moderate

CONFIRMED SYMPTOMS (from consultation)

- stomach pain

- nausea

DETAILED CLINICAL SUMMARY

In acute gastro-enteritis, the lining of the stomach and intestines becomes inflamed, often due to pathogens such as viruses (e.g., norovirus, rotavirus) or bacteria (e.g., E. coli, Salmonella). This inflammation disrupts normal digestive processes, resulting in symptoms like diarrhea (often watery), nausea, and abdominal discomfort. The condition can be triggered by the consumption of contaminated food or water, or by exposure to infected individuals. Management typically involves rehydration, dietary adjustments, and, in some cases, medications to alleviate symptoms. Most cases resolve wi...

Rx PRIMARY RECOMMENDED MEDICATIONS

Commonly available in India - consult a physician before use

Medication	Dosage	Frequency	Duration
Loperamide / Imodium 2mg	2 mg after the first loose stool, then 1 mg per day	As needed, up to 8 times per day	Until symptoms resolve, not exceeding 48 h

ALTERNATIVE MEDICATIONS (Available in India)

OTC substitutes or alternatives if primary is unavailable or not tolerated

Medication	Dosage	Frequency	Notes
Probiotics / Enterogermina	1 vial (5ml)	Once daily	Consider this alternative to restore gut f
Oral Rehydration Salts (ORS) / Electral	1 packet dissolved in 1 liter of clean water	As needed to prevent dehydration	Essential for rehydration, especially if d

HOME REMEDIES & SUPPORTIVE CARE

Safe at-home measures alongside medical treatment

- Ginger tea: Boil fresh ginger slices in water for 10 minutes, strain, and drink warm. Ginger has anti-inflammatory properties and can help soothe the stomach.

- Rice w
digestive tra

LIFESTYLE & PREVENTIVE ADVICE

- Increase fluid intake: Encourage the patient to drink plenty of fluids, especially ORS, to prevent dehydration and support recovery.

- Avoid dai

ALLERGIES & DRUG INTERACTIONS

- The patient has known allergies to milk, dust, and pollen. Care should be taken to avoid dairy products to prevent exacerbation of gastrointestinal symptoms.

RECOMMENDED NEXT STEPS

- Consider stool analysis to check for pathogens if symptoms persist beyond 48 hours.

- Consult a

DR. AI CONSULTANT

Electronically Generated via MediScan AI

Disclaimer: This AI-assisted report is based on user-reported symptoms for informational support only. It does not replace in-person doctor consultation, physical examination, or clinical tests.