

Formule di elettro- e magnetostatica

Tommaso Brusco

16 dicembre 2020

1 Equazioni di Maxwell

$$\begin{aligned}\operatorname{div} \vec{E} &= 4\pi\rho & \operatorname{div} \vec{E} &= 0 & \vec{E} &= -\nabla V \\ \operatorname{div} \vec{B} &= 0 & \operatorname{div} \vec{B} &= \frac{4\pi}{c} \vec{J} & \vec{B} &= \operatorname{div} \vec{A}\end{aligned}$$

$$\vec{F} = q \left(\vec{E} + \frac{\vec{v}}{c} \times \vec{B} \right)$$

2 Risolvere Poisson o Laplace

2.1 Funzione di Green

$$V(\vec{r}) = \int d\vec{r}'^3 \rho(\vec{r}') G(\vec{r}, \vec{r}') - \frac{1}{4\pi} \oint_S \left(V \frac{\partial G}{\partial u} - G \frac{\partial V}{\partial u} \right)$$

Per il piano:

$$G(\vec{r}, \vec{r}') = \frac{1}{|\vec{r} - \vec{r}'|} - \frac{1}{|\vec{r} + \vec{r}'|}$$

Per la sfera (raggio a):

$$G(\vec{r}, \vec{r}') = \frac{1}{|\vec{r} - \vec{r}'|} - \frac{a}{r'} \frac{1}{|\vec{r} - \frac{a^2}{r'} \vec{r}'|}$$

2.2 Polinomi di Legendre

$$V(r, \theta) = \sum_{l=0}^{\infty} \left(A_l r^l + \frac{B_l}{r^{l+1}} \right) P_l(\cos \theta)$$

$$P_0 = 1 \quad P_1 = \cos \theta \quad P_2 = \frac{1}{2} (3 \cos^2 \theta - 1) \quad P_l(-x) = (-1)^l P_l(x)$$

$$\text{Normalizzazione: } \int_{-1}^1 P_{l'}(x) P_l(x) dx = \frac{2}{2l+1} \delta_{l,l'}$$

$$\frac{1}{|\vec{r} - \vec{r}'|} = \sum_{l=0}^{\infty} \frac{r_{<}^l}{r_{>}^{l+1}} P_l(\cos \theta)$$

2.3 Armoniche sferiche

$$V(r, \theta, \phi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l \left(A_{l,m} r^l + B_{l,m} \frac{1}{r^{l+1}} \right) Y_{l,m}(\theta, \phi)$$

Lo sviluppo in multipoli di una distribuzione $\rho(\vec{r}')$ limitata a grande distanza dall'origine è:

$$V(\vec{r}) = \sum_{l=0}^{\infty} \sum_{m=-l}^l C_{l,m} \frac{Y_{l,m}(\theta, \phi)}{r^{l+1}}$$

con $C_{l,m} = \int dr'^3 r'^l \rho(\vec{r}') Y_{l,m}(\theta, \phi)$. Similmente vicino all'origine. Le prime armoniche sferiche sono:

	0	1	2	3
0	$\frac{1}{2}\sqrt{\frac{1}{\pi}}$	$\frac{1}{2}\sqrt{\frac{3}{\pi}} \cos \theta$	$\frac{1}{4}\sqrt{\frac{5}{\pi}} (3 \cos^2 \theta - 1)$	$\frac{1}{4}\sqrt{\frac{7}{\pi}} (5 \cos^3 \theta - 3 \cos \theta)$
1		$-\frac{1}{2}\sqrt{\frac{3}{2\pi}} e^{i\phi} \sin \theta$	$-\frac{1}{2}\sqrt{\frac{15}{2\pi}} e^{i\phi} \cos \theta \sin \theta$	$-\frac{1}{8}\sqrt{\frac{21}{2\pi}} e^{i\phi} \sin \theta (5 \cos^2 \theta - 1)$
2			$\frac{1}{4}\sqrt{\frac{15}{2\pi}} e^{2i\phi} \sin^2 \theta$	$\frac{1}{4}\sqrt{\frac{105}{2\pi}} e^{2i\phi} \sin^2 \theta \cos \theta$
3				$-\frac{1}{8}\sqrt{\frac{35}{\pi}} e^{3i\phi} \sin^3 \theta$

3 Biot-Savart e altro magnetismo

Se la distribuzione di corrente è localizzata

$$\vec{B} = \frac{1}{c} \int dr'^3 \frac{\vec{J} \times (\vec{r} - \vec{r}')}{|\vec{r} - \vec{r}'|^3} \quad \vec{A} = \frac{1}{c} \int dr'^3 \frac{\vec{J}(\vec{r}')}{|\vec{r} - \vec{r}'|}$$

3.1 Multipoli magnetici

$$\vec{A} = \frac{1}{rc} \int dr'^3 \vec{J}(\vec{r}') \left(1 + \frac{\vec{r} \cdot \vec{r}'}{r^2} \right) + \frac{1}{r^3 c} \int dr'^3 \vec{J}(\vec{r}') (xx' + yy' + zz')$$

4 Dipoli e altri cazzilli

Si indica con \vec{p} il dipolo elettrico, con \vec{m} quello magnetico e con \vec{s} uno qualsiasi dei due. Inoltre si indica con \vec{G} un campo qualsiasi (elettrico o magnetico).

$$\vec{G} = \frac{3\vec{r}(\vec{s} \cdot \vec{r}) - r^2 \vec{s}}{r^5}$$

$$\vec{F} = (\vec{p} \cdot \vec{\nabla}) \vec{E} \quad U = -\vec{p} \cdot \vec{E} \quad U = \frac{1}{8\pi} \int |\vec{E}|^2 dr^3$$

$$\vec{m} = \frac{1}{2c} \int dr^3 \vec{r} \times \vec{J} \quad \vec{A} = \frac{\vec{m} \times \vec{r}}{r^3} \quad \vec{\tau} = \vec{m} \times \vec{B} \quad \vec{F} = \vec{\nabla} (\vec{m} \cdot \vec{B})$$

5 Operatori differenziali

5.1 Gradiente

$$\left(\frac{\partial f}{\partial \rho}, \frac{1}{\rho} \frac{\partial f}{\partial \phi}, \frac{\partial f}{\partial z} \right)$$

$$\left(\frac{\partial f}{\partial \rho}, \frac{1}{\rho} \frac{\partial f}{\partial \theta}, \frac{1}{\rho \sin \theta} \frac{\partial f}{\partial \phi} \right)$$

5.2 Divergenza

$$\frac{1}{\rho} \frac{\partial (\rho F_\rho)}{\partial \rho} + \frac{1}{\rho} \frac{\partial F_\phi}{\partial \phi} + \frac{\partial F_z}{\partial z}$$

$$\frac{1}{\rho^2} \frac{\partial (\rho^2 F_\rho)}{\partial r} + \frac{1}{\rho \sin \theta} \frac{\partial (\sin \theta F_\theta)}{\partial \theta} + \frac{1}{\rho \sin \theta} \frac{\partial F_\phi}{\partial \phi}$$

5.3 Rotore

$$\left(\frac{1}{\rho} \frac{\partial F_z}{\partial \phi} - \frac{\partial F_\phi}{\partial z}, \frac{\partial F_\rho}{\partial z} - \frac{\partial F_z}{\partial \rho}, \frac{1}{\rho} \left(\frac{\partial (\rho F_\phi)}{\partial \rho} - \frac{\partial F_\rho}{\partial \phi} \right) \right)$$

$$\left(\frac{1}{\rho \sin \theta} \left(\frac{\partial (\sin \theta F_\phi)}{\partial \theta} - \frac{\partial F_\theta}{\partial \phi} \right), \frac{1}{\rho} \left(\frac{1}{\sin \theta} \frac{\partial F_\rho}{\partial \phi} - \frac{\partial (\rho F_\phi)}{\partial \rho} \right), \frac{1}{\rho} \left(\frac{\partial (\rho F_\theta)}{\partial \rho} - \frac{\partial F_\rho}{\partial \theta} \right) \right)$$

5.4 Laplaciano

$$\frac{1}{\rho} \frac{\partial}{\partial \rho} \left(\rho \frac{\partial f}{\partial \rho} \right) + \frac{1}{\rho^2} \frac{\partial^2 f}{\partial \phi^2} + \frac{\partial^2 f}{\partial z^2}$$

$$\frac{1}{\rho^2} \frac{\partial}{\partial \rho} \left(\rho^2 \frac{\partial f}{\partial \rho} \right) + \frac{1}{\rho^2 \sin \theta} \frac{\partial}{\partial \theta} \left(\sin \theta \frac{\partial f}{\partial \theta} \right) + \frac{1}{\rho^2 \sin^2 \theta} \frac{\partial^2 f}{\partial \phi^2}$$