

The ultimate guide to Maxwell layer potentials on the sphere

In this note, we consolidate analytic expressions for various maxwell layer potentials on the sphere. In the following, $P_{nm}(t)$ are the associated legendre polynomials of degree n and order m , $Y_{nm}(\theta, \phi)$, are the spherical harmonics of degree n and order m given by

$$Y_{nm}(\theta, \phi) = \sqrt{2n+1} \frac{(n-|m|)!}{(n+|m|)!} P_{n|m|}(\cos \theta) e^{im\phi}. \quad (0.1)$$

Let $\mathbf{Y}_{nm}(\theta, \phi)$, $\mathbf{\Psi}_{nm}(\theta, \phi)$, $\mathbf{\Phi}_{nm}(\theta, \phi)$ denote the vector spherical harmonics given by

$$\begin{aligned} \mathbf{Y}_{nm}(\theta, \phi) &= Y_{nm}(\theta, \phi) \begin{bmatrix} \sin \theta \cos \phi \\ \sin \theta \sin \phi \\ \cos \theta \end{bmatrix}, \\ \mathbf{\Psi}_{nm}(\theta, \phi) &= -P'_{n|m|}(\cos \theta) \sin \theta \begin{bmatrix} \cos \theta \cos \phi \\ \cos \theta \sin \phi \\ -\sin \theta \end{bmatrix} + \frac{imY_{nm}(\theta, \phi)}{\sin(\theta)} \begin{bmatrix} -\sin \phi \\ \cos \phi \\ 0 \end{bmatrix}, \\ \mathbf{\Phi}_{nm}(\theta, \phi) &= \hat{\mathbf{r}} \times \mathbf{\Psi}_{nm}(\theta, \phi). \end{aligned} \quad (0.2)$$

Let $j_n(z)$, $h_n(z)$ denote the spherical Bessel and spherical Hankel functions of degree n . Let Ω be the interior of the unit sphere $\partial\Omega$ denote it's boundary. Finally, let $\mathcal{S}_k[\sigma]$ denote the Helmholtz single layer potential given by

$$\mathcal{S}_k[\sigma](\mathbf{x}) = \int_{\partial\Omega} \frac{e^{ik|\mathbf{x}-\mathbf{y}|}}{|\mathbf{x}-\mathbf{y}|} \sigma(\mathbf{y}) dS_{\mathbf{y}}. \quad (0.3)$$

For any $\mathbf{x} \in \Omega$, let (r, θ, ϕ) , denote it's spherical coordinates. Then the following equalities hold:

$$\begin{aligned} \mathcal{S}_k[\mathbf{\Psi}_{nm}](\mathbf{x}) &= in(n+1) \left(\frac{j_n(kr)}{kr} (h_n(k) + kh'_n(k)) + j'_n(kr)h_n(k) \right) \mathbf{Y}_{nm}(\theta, \phi) + \\ &\quad i \left(\left(j'_n(kr) + \frac{j_n(kr)}{kr} \right) (h_n(k) + kh'_n(k)) + n(n+1)h_n(k) \frac{j_n(kr)}{kr} \right) \mathbf{\Psi}_{nm}(\theta, \phi), \\ \mathcal{S}_k[\mathbf{\Phi}_{nm}](\mathbf{x}) &= ikj_n(kr)h_n(k) \mathbf{\Phi}_{nm}(\theta, \phi), \\ \nabla \mathcal{S}_k[Y_{nm}](\mathbf{x}) &= ik^2 h_n(k) j'_n(kr) \mathbf{Y}_{nm}(\theta, \phi) + ikh_n(k) \frac{j_n(kr)}{r} \mathbf{\Psi}_{nm}(\theta, \phi), \\ \nabla \times \mathcal{S}_k[\mathbf{\Psi}_{nm}](\mathbf{x}) &= -ikj_n(kr) (h_n(k) + kh'_n(k)) \mathbf{\Phi}_{nm}(\theta, \phi), \\ \nabla \times \mathcal{S}_k[\mathbf{\Phi}_{nm}](\mathbf{x}) &= -in(n+1)kh_n(k) \frac{j_n(kr)}{r} \mathbf{Y}_{nm}(\theta, \phi) - ih_n(k)k^2 \left(j'_n(kr) + \frac{j_n(kr)}{kr} \right) \mathbf{\Psi}_{nm}(\theta, \phi), \\ \nabla \times \nabla \times \mathcal{S}_k[\mathbf{\Psi}_{nm}](\mathbf{x}) &= in(n+1)k \frac{j_n(kr)}{r} (h_n(k) + kh'_n(k)) \mathbf{Y}_{nm}(\theta, \phi) + \\ &\quad i \left(k^2 j'_n(kr) + k \frac{j_n(kr)}{r} \right) (h_n(k) + kh'_n(k)) \mathbf{\Psi}_{nm}(\theta, \phi), \\ \nabla \times \nabla \times \mathcal{S}_k[\mathbf{\Phi}_{nm}](\mathbf{x}) &= ik^3 j_n(kr)h_n(k) \mathbf{\Phi}_{nm}(\theta, \phi). \end{aligned} \quad (0.4)$$