

# Study of impurities influence on the hydrogen dissolution energy in the bcc iron

Maxim Rakitin


maxim.rakitin@gmail.com, rms85@physics.susu.ac.ru

General and Theoretical Physics Department, South Ural State University  
(National Research University), Chelyabinsk, Russia

# Outline

- 1 Introduction
- 2 Problem
- 3 Calculation method
- 4 Parameters of the simulation
- 5 Results
- 6 Conclusions

# Hydrogen embrittlement of metals



I.M. Robertson / Engineering Fracture Mechanics 68 (2001) 671-692

The following processes can take place during cooling:

- Sharp decreasing of hydrogen solubility
- Defects formation (cracks, blisters)
- Plasticity decreasing and brittleness increasing
- Material degradation

# Ways to avoid hydrogen embrittlement

- Vacuumizing during a melting process
  - Purging of a liquid metal by noble gases
- } Expensive!
- Doping by impurities which are able to trap hydrogen and increase its solubility in iron



Simple and accessible way to control hydrogen content in iron

Interesting impurities: Palladium [1], Titanium [2, 3], Chromium [4], Vanadium [3], etc.

[1] Arkharov, et al. / PhChMM (USSR) 12 (1976) 47–51

[2] Myers, et al. / Rev. Mod. Phys. 64 (1992) 559–617

[3] Sigworth et al. / Metal Science 8 (1974) 298–310

[4] Thomas et al. / Metal. Trans. A. 33 (2002) 1991–2004

# Problem

To understand influence of impurities on hydrogen dissolution we need to calculate **dissolution energy** of hydrogen in bcc iron with impurities and **binding energy** of hydrogen with the impurities.

To solve this problem we need:

- 1 Find optimal parameters of the simulation method.
- 2 Calculate dissolution energy and binding energy of hydrogen in bcc iron with the following impurities: Mg, Al, Si,  $3d$  (Sc, Ti, V, Cr, Mn, Co, Ni, Cu, Zn) and  $4d$  (Y, Zr, Nb, Mo, Pd, Cd).
- 3 Calculate elastic and electronic contributions to the dissolution energies of hydrogen in bcc iron with the impurities.

# Dissolution and binding energy of hydrogen

Dissolution energy of hydrogen in iron:

$$\underbrace{E_{\text{diss}}(\text{H})}_{\sim 0.1 \text{ eV}} = \underbrace{E(\text{Fe}_n\text{H})}_{\sim 10^4 \dots 10^6 \text{ eV}} - \underbrace{E(\text{Fe}_n)}_{\sim 10^4 \dots 10^6 \text{ eV}} - \frac{1}{2}E[\text{H}_{2(g)}]$$

$E_{\text{diss}}(\text{H})$  — dissolution energy of H in Fe

$E(\text{Fe}_n\text{H})$  —  $\text{Fe}_n\text{H}$  supercell total energy

$E(\text{Fe}_n)$  —  $\text{Fe}_n$  supercell total energy

$E[\text{H}_{2(g)}]$  — total energy of the hydrogen molecule

$n$  — number of atoms in the ideal supercell

Binding energy of hydrogen with impurity metal **Me**:

$$E_{\text{b}}(\text{H-Me}) = \underbrace{(E(\text{Fe}_n\text{H}) - E(\text{Fe}_n))}_{E_{\text{diss}}(\text{H in Fe}_n\text{H})} - \underbrace{(E(\text{Fe}_{n-1}\text{MeH}) - E(\text{Fe}_{n-1}\text{Me}))}_{E_{\text{diss}}(\text{H in Fe}_{n-1}\text{MeH})}$$

# Calculation method




www.wien2k.at


We used WIEN2k simulation package based on **Density Functional Theory (DFT)** with **Full-Potential Linearized Augmented Planewave Method (FP-LAPW)**.

## Input:

Lattice constants (a,b,c)  
Supercell  
Lattice type (BCC, FCC, HCP, ...)  
NM, FM, AFM  
RK<sub>max</sub>  
k-points  
...



single mode



parallel mode

## Output

Total energy  
Forces on atoms  
Optimized structure  
DOS  
Electronic density  
...

# Parameters of the simulation

## Main parameters:

Supercell with bcc structure:

(53 atoms of Fe) + 1 impurity atom (Mg, Al, Si, Sc, Ti, V, Cr, Mn, Co, Ni, Cu, Zn, Y, Zr, Nb, Mo, Pd, Cd) + (1 atom of H)

$R_{mt}(\text{Fe})$ : 1.06 Å (2.0 a.u.)

$R_{mt}(\text{impurities})$ : 1.16 Å (2.2 a.u.)

$R_{mt}(\text{H})$ : 0.37 Å (0.7 a.u.)

$V_{xc}$ : GGA-PBE96

$E_{cut-off}$ : 340 eV

Number of  $k$ -points: 24



## Convergence criteria

energy: 1.0 meV

charge: 0.001  $e^-$



forces: 0.01 eV/Å

(volopt + min\_lapw on each step)





# Dissolution energy of H in Fe with impurities



# Contributions in the dissolution energy

$$E_{\text{diss}} = E_{\text{Fe}_{53}\text{MeH}}^{\text{relax}} - E_{\text{Fe}_{53}\text{Me}}^{\text{relax}} - \frac{1}{2}E_{\text{H}_2} = -E_{\text{elastic}} + E_{\text{electronic}}$$


$$E_{\text{elastic}} = -E_{\text{Fe}_{53}\text{MeH}}^{\text{relax}} + E_{\text{Fe}_{53}\text{MeH}}^{\text{nonrelax}}$$

$$E_{\text{electronic}} = E_{\text{Fe}_{53}\text{MeH}}^{\text{nonrelax}} - E_{\text{Fe}_{53}\text{Me}}^{\text{relax}} - \frac{1}{2}E_{\text{H}_2} = E_{\text{electronic}}^{\text{local}} + E_{\text{electronic}}^{\text{nonloc}}$$


$$E_{\text{electronic}}^{\text{local}} = E_{\text{Fe}_{53}\text{MeH}}^{\text{nonrelax}} - E_{\text{Fe}_{53}\text{Me}}^{\text{relax}} - E_{\text{Fe}_{54}\text{H}}^{\text{nonrelax}} + E_{\text{Fe}_{54}}^{\text{relax}}$$

$$E_{\text{electronic}}^{\text{nonloc}} = E_{\text{Fe}_{54}\text{H}}^{\text{nonrelax}} - E_{\text{Fe}_{54}}^{\text{relax}} - \frac{1}{2}E_{\text{H}_2}$$

# Contributions in dissolution energy of H in Fe with 3d and 4d metals




# Comparison of local electronic contributions with hydrides formation energies




Smithson et al. / Phys. Rev. B 66, 144107 (2002)

# Comparison of elastic energies with atomic radii of 3d and 4d metals



# Binding energy with $3d$ and $4d$ metals



# Conclusions

- Influence of Mg, Al, Si, 3d (Sc, Ti, V, Cr, Mn, Co, Ni, Cu, Zn) and 4d (Y, Zr, Nb, Mo, Pd, Cd) metallic impurities on dissolution energy of hydrogen in bcc iron was examined using first principles calculations in WIEN2k *ab initio* simulation package.
- Elastic and electronic contributions to the dissolution energies of hydrogen in bcc iron with impurities were calculated. We found direct connection of the elastic contributions with the atomic radii of the metallic impurities, and electronic contributions with the hydrides formation energies of the corresponding metals.
- Binding energies of hydrogen with the impurities were computed and the highest binding energies were observed for hydrogen and Sc, Ti, Y, Zr, Pd, Cd. The data are in good agreement with the recent work of Counts et al. It is worth to note, that in the literature Zr and Pd are not present, and our investigation fills up the gap.

## Main publications:

- Theory of hydrogen solubility in binary iron alloys based on *ab initio* calculation results / D. A. Mirzaev, A. A. Mirzoev, K. Yu. Okishev, M. S. Rakitin // *Molecular Physics*. — 2012. — V.110. — P. 1299–1304.
- Theory of Hydrogen Solubility in Binary Iron Alloys Based on First-Principles Calculation Results / A. A. Mirzoev, M. S. Rakitin, D. A. Mirzaev, K. Yu. Okishev // *Thermodynamics 2011: Book of abstracts*. — Athens, Greece. — 2011. — P. 524.
- *Rakitin, M. S.* Effect of palladium and titanium impurities on hydrogen solubility in bcc iron / M. S. Rakitin, A. A. Mirzoev // *Summer School: Computational Materials Science: Contributions*. — San Sebastian, Spain. — 2010. — P. 54–55.

