
BUDGIE

Melopsittacus undulatus



Beginner

Complete Care Package

Quick Profile · Care Guide · Health · Quick Reference · Owner Tools

BeastlyFacts

Practical care guides for curious animal lovers

Contents

How to Use This Package

How to read this manual and which pages to print first

1. Quick Profile

Species overview, fact table, fun facts, and cost range

2. Full Care Guide

Cage, diet, handling, enrichment, common mistakes

3. Health & Common Issues

Fatty liver, mites, respiratory, red flags

4. Quick Reference & Checklist

Setup checklist and targets

Emergency & Quick Targets

One-page emergency card to post near the cage

5. Owner Tools & Decision Guides

Budget, shopping list, first 30 days, symptoms, daily routine

Owner Log

Weight and daily tracking tables

Enrichment Checklist

Setup and monthly enrichment review

Glossary

Key terms in plain language

How to Use This Package

This is a practical manual, not a textbook. Use the Quick Profile for orientation, the Care Guide for day-to-day husbandry, and Health when something looks wrong. Owner Tools are designed to print and post near the cage.

This package is for educational purposes and is not a substitute for veterinary care. Birds hide illness until late - when in doubt, contact an avian veterinarian.

Print these first

- Symptom quick reference (Health) - page 7
- Setup checklist and targets (Quick Reference) - page 8
- Emergency card - page 9
- Shopping list and budget tiers (Owner Tools) - page 10
- First 30 days checklist - page 10
- Daily and weekly routine - page 11

1. Quick Profile

Budgies (*Melopsittacus undulatus*), also called parakeets, are small, social Australian parrots from open grassland and scrub. Wild flocks are highly mobile seed-eaters. In captivity they are popular first birds, but they still need flight space, a pellet-based diet, foraging work, and either a companion or substantial daily interaction.

Size: About 7 in (18 cm); 1-1.4 oz

Lifespan: 5-8 years typical; 12-15 possible with excellent care

Difficulty: Beginner (with correct diet and social plan)

Temperament: Social, vocal, active; best with a companion or lots of human time

Cost overview

Expect a solid starter cage and accessories to run roughly \$100-250 depending on size and brand. Monthly food (pellets, fresh produce, limited seed) is modest, but budget for an avian vet wellness exam and emergency fund - bird clinics are specialized and not cheap. A second bird adds ongoing food cost but often reduces behavioral problems from loneliness.

2. Full Care Guide

Housing & Cage

Minimum often cited for a single budgie is about 18x18x24 inches, with wider preferred so the bird can fly side-to-side. Bar spacing must be 1/2 inch or less - wider bars risk head entrapment. Place the cage at roughly eye level, away from kitchens (Teflon/PTFE fumes from overheated nonstick cookware are lethal), drafts, and direct blazing sun. Provide multiple perches of different diameters and textures (natural wood preferred over uniform dowels).



Example budgie cage layout (BeastlyFacts tank setup guide).

Diet & Feeding

Aim for about 60-70% high-quality small-parrot pellets (brands such as Harrison's Fine, Roudybush, or similar). Offer fresh vegetables daily (leafy greens, broccoli, carrot, bell pepper, cucumber). Seeds should be no more than about 10-20% of the diet and are best delivered inside foraging toys rather than an open dish. All-seed diets drive fatty liver disease, obesity, and shortened lifespan. Millet sprays are high-fat treats - small pieces a few times per week at most. Fresh water daily. Cuttlebone or a mineral block is useful especially if any seed is offered.

Handling & Social Needs

Budgies are flock animals. A bonded pair is often easier behaviorally than a single bird that must get all social needs from humans. Single birds need substantial daily interaction. Provide at least 30-60 minutes of supervised out-of-cage time in a bird-safe room (no open water, ceiling fans off, windows secured, toxic plants removed). Train step-up gently with positive reinforcement.

Common Mistakes to Avoid

- **Never** keep an all-seed diet as the main menu - fatty liver and deficiencies follow.
 - **Never** use cages with bar spacing wider than 1/2 inch.
 - **Never** place the cage in the kitchen where nonstick cookware is used.
 - **Never** leave a single budgie without either a companion or hours of daily interaction.
 - **Never** ignore tail-bobbing, open-mouth breathing, or floor-sitting - these are emergencies.
 - **Never** skip toy rotation - static toys become furniture.
-

Enrichment

Wild budgies spend a large share of the day flying, foraging, and interacting with flock members. In a cage with an open seed dish those hours collapse into minutes. Enrichment restores work, chewing, climbing, and social outlets so the bird stays mentally busy.

Foraging Enrichment

Start simple: hide seeds or millet under crumpled plain paper, inside a paper cup, or in a shallow tray of shredded paper. Progress to beginner foraging toys (treat balls, soft shreddable pockets), then gradually increase difficulty. Scatter a portion of the daily pellet ration. Clip leafy greens or herb sprigs through the bars. Never jump straight to hard acrylic puzzles for a bird that has never foraged - frustration leads to abandonment.

Shredding, Chewing & Beak Work

Budgies have continuously growing beaks and a strong drive to destroy soft materials. Offer soft wood, palm leaf, sola, seagrass, and plain cardboard or paper. Hanging shreddables and foot toys the bird can hold and demolish work well. Treat them as consumables and replace as destroyed. Avoid glitter, foil, sharp wire, toxic dyes, and parts small enough to swallow.

Perches, Climbing & Cage Layout

Provide perches of varied diameter and texture (natural wood preferred over uniform dowels). Add a swing, ladder, or rope perch for movement between levels. Zone the cage: top for swings and hanging foragers, middle for chew and puzzle toys near favorite perches, bottom for foot toys or a foraging tray. Leave open stretch/flight space - do not pack the cage full of toys.

Social Contact & Toy Rotation

Budgies are flock animals. A compatible pair provides built-in enrichment; a single bird needs substantial daily interaction. Rotate 3-5 toys in the cage at a time from a larger collection every few days to weekly. Supervised out-of-cage time on a play gym adds flight and exploration. Positive-reinforcement training (step-up, target, simple tricks) is high-value mental work and strengthens the human-bird bond.

3. Health & Common Issues

Birds hide illness until they are very sick. The most common preventable problems in budgies are diet-driven (fatty liver, obesity, nutritional deficiencies) and under-stimulation. Respiratory disease and mites also appear regularly. Know your bird's normal droppings, weight, and activity so changes stand out.

Red Flags - Contact an Avian Vet Promptly

- Fluffed feathers combined with daytime sleeping
- Tail-bobbing with each breath or open-mouth breathing
- Sitting on the cage floor instead of perching
- Sudden appetite loss or voice change
- Crusty honeycomb buildup on cere, beak, or legs (scaly mites)
- Persistent sneezing, nasal discharge, or lethargy

Tell the vet diet history (especially seed-only), recent weight, and when signs started. Psittacosis is zoonotic - mention respiratory signs, diarrhea, and weakness. Find an avian vet before an emergency.

Symptom Quick Reference

Sign	Possible concern	Action
Tail-bob / open-mouth breath	Respiratory distress	Avian vet ASAP
Floor-sitting, fluffed, lethargy	Serious illness	Avian vet within 24h
Crusty cere / legs	Scaly mites	Avian vet
Obesity, poor stamina	Seed-heavy diet / fatty liver	Diet fix + vet

Fatty Liver & Seed-Only Diets

All-seed diets are a leading driver of hepatic lipidosis, obesity, lipomas, and shortened lifespan. Transition to a pellet base with daily vegetables; keep seeds as foraging enrichment only.

Scaly Face / Leg Mites

Knemidocoptes mites cause crusty, honeycomb-like buildup on the cere, beak, and legs. Veterinary diagnosis and treatment are required - do not pick at crusts at home.

Respiratory Disease

Tail-bobbing, open-mouth breathing, and floor-sitting are late signs. Air quality, stress, and infectious agents (including zoonotic psittacosis) all matter. Prompt avian care is appropriate.

4. Quick Reference & Checklist

Essential setup checklist

- Cage at least ~18x18x24 in (wider better); bar spacing 1/2 in or less
 - Multiple natural-wood perches of varied diameter
 - Pellet-based diet + daily vegetables
 - Seeds only as minority / foraging portion
 - Foraging and shredding toys with a rotation plan
 - Cuttlebone or mineral block
 - Bird-safe room for daily out-of-cage time
 - Companion bird plan OR scheduled multi-hour interaction
 - Avian veterinarian identified
-

Quick targets

Target	Value
Cage minimum (single)	18x18x24 in - wider preferred
Bar spacing	1/2 inch or less
Diet base	60-70% quality small-parrot pellets
Seeds	Max ~10-20%, preferably in foraging toys
Out-of-cage time	30-60+ min daily supervised

Emergency & Quick Targets - Budgie

Print this page and keep it near the enclosure. When something looks wrong, start here.

Quick Targets

Parameter	Target
Cage minimum (single)	18x18x24 in - wider preferred
Bar spacing	1/2 inch or less
Diet base	60-70% quality small-parrot pellets
Seeds	Max 10-20% - foraging only, not open dish
Out-of-cage time	30-60+ min daily supervised

Red Flags - Act Early

- Fluffed feathers + daytime sleeping
 - Tail-bobbing with each breath or open-mouth breathing
 - Sitting on cage floor instead of perching
 - Sudden appetite loss or voice change
 - Crusty honeycomb buildup on cere/beak/legs (mites)
 - Persistent sneezing, nasal discharge, lethargy
-

Do This Now

- Any tail-bobbing or floor-sitting - avian vet within 24 hours
 - Note diet (especially seed-only history) and recent changes
 - Isolate from other birds if infectious disease suspected
-

Full symptom triage table (sign → concern → action) is in **3. Health & Common Issues**.

5. Owner Tools & Decision Guides

Budget tiers (approximate, single bird)

Tier	Upfront	What you get
Functional	\$100-150	Minimum-size cage, basic perches, dishes, starter pellets/toys
Recommended	\$150-250	Wider cage, natural perches, foraging + shred toys, scale, carrier
Premium	\$250-400+	Large flight-friendly cage, full toy rotation set, play gym, first avian visit

Monthly food is modest (\$15-40). Budget separately for an avian wellness exam and emergency care - specialized bird clinics cost more than general practices.

Shopping List

- Cage at least ~18x18x24 in (wider strongly preferred); bar spacing 1/2 in or less
 - Multiple natural-wood perches of varied diameter and texture
 - Food and water dishes (ceramic or stainless preferred)
 - High-quality small-parrot pellets (main diet)
 - Limited seed mix for foraging toys only (not open-dish staple)
 - Cuttlebone or mineral block
 - Foraging toys (paper wraps, cups, beginner puzzles)
 - Shreddable toys (palm, sola, soft wood, plain cardboard)
 - Swing and/or ladder
 - Play gym or out-of-cage stand
 - Travel carrier
 - Digital scale (grams) for weight checks
 - Budget for initial avian wellness exam
-

First 30 Days

- Days 1-3: Quiet settling. Offer pellets and familiar food; begin gentle vegetable introductions. Minimal forced handling.
- Days 4-7: Short training sessions (target or step-up). Confirm droppings and appetite are normal. Start simple foraging.
- Week 2: Establish out-of-cage routine in a bird-safe room. Rotate one toy. Weigh and log baseline.
- Week 3-4: Full daily routine. Book or confirm avian wellness visit. Assess whether a companion is planned.

Daily & Weekly Routine

Print this page and post it near the cage. Check boxes as you go. Use the date lines to track consistency.

Daily

- Fresh water; refresh pellets and vegetables
 - Quick visual check (posture, droppings, breathing, activity)
 - Foraging opportunity (paper wrap, cup, or scatter)
 - Social time / training or pair interaction
 - Supervised out-of-cage time when schedule allows
-

Weekly

- Wash dishes and wipe cage bars/perches
 - Replace destroyed shreddables; rotate 1-2 toys
 - Weigh bird and log the number
 - Change cage paper / substrate
 - Inspect feet, cere, and vent
-

Monthly / ongoing

- Deep-clean cage and accessories
- Review weight trend and diet balance
- Confirm avian vet contact is current
- Reassess social plan (companion vs human time)

Week of: _____ Initials: _____

Week of: _____ Initials: _____

Week of: _____ Initials: _____

Owner Log - Budgie

Track weight and basic daily notes. Weight change is often the earliest warning sign. Print extra copies as needed.

Weight & Daily Log

Date	Weight (g)	Ate?	Droppings	Notes

Seed Diet & Respiratory Notes

- All-seed diets drive fatty liver disease, obesity, and shortened lifespan. Transition to pellets with vegetables; use seeds only inside foraging toys.
- Tail-bobbing, open-mouth breathing, or floor-sitting are emergencies - birds hide illness until late.
- Psittacosis is zoonotic - mention respiratory signs, diarrhea, and weakness to the vet.
- Find an avian vet before an emergency; many general practices will not see birds.

Enrichment Checklist - Budgie

Enrichment is not optional decoration. Use this as a setup and monthly review checklist.

Setup & Ongoing Checklist

- Daily foraging - paper wraps, cups, treat balls, scatter feed
- Shreddable toys (paper, palm, sola, soft wood) rotated weekly
- Varied natural-wood perch diameters and textures
- Swing + ladder + open stretch space (do not overcrowd)
- Rotate 3-5 toys from a larger collection every few days
- Supervised out-of-cage time on a play gym
- Compatible companion preferred; single birds need hours of interaction
- Positive reinforcement training (step-up, target)

Monthly review notes (write in):

Date: _____ What changed: _____

Date: _____ What changed: _____

Date: _____ What changed: _____

Rule: stability for core hides/heat/water systems; novelty only in non-critical items.

Glossary

Cere. Fleshy area above the beak; color can help sex adult budgies and show mite damage.

Fatty liver (hepatic lipidosis). Serious liver disease often driven by all-seed diets and obesity.

Foraging. Working to find and extract food - a core wild behavior that should be restored in captivity.

Psittacosis. Disease caused by *Chlamydia psittaci*; can infect birds and people.

Scaly face mites. *Knemidocoptes* infestation causing crusty buildup on cere, beak, and legs.

Tail-bobbing. Visible up-down tail motion with each breath - a red-flag respiratory sign.

This care package was compiled from BeastlyFacts research and practical husbandry standards.

Version 1.0 · August 2026

© BeastlyFacts. All rights reserved.

Disclaimer: This package is for educational purposes and is not a substitute for veterinary care. Husbandry standards evolve; always verify critical health decisions with a qualified avian veterinarian.

No external links are included in this document.
