Mass Gap for the O(N) Model: Analysis of the HS Approach

M. R. Douglas

April 2026

Abstract

We analyze the Hubbard-Stratonovich approach to the mass gap for the O(N) model in two dimensions. The HS identity for the Euclidean quartic requires imaginary coupling (sign problem). For the NLSM, Kupiainen's trick adds a real mass m_0^2 for free via the constraint, giving decay for each σ-value. For the LSM, no such trick exists: the real mass cancels after HS, and the mass gap must emerge entirely from the oscillatory σ-average. We compare the two settings and discuss the implications for a formal proof at large N.

Contents

1 The HS identity and the sign problem **1**
1.1 The Euclidean quartic requires imaginary coupling . . . 1
1.2 The σ-integral on the real axis . . . 2

2 The NLSM: Kupiainen's mass trick **2**
2.1 Adding m_0^2 for free . . . 2
2.2 The NLSM operator has real mass . . . 2
2.3 Kupiainen's proof structure . . . 3

3 The LSM: the mass cancels **3**
3.1 Add-subtract fails . . . 3
3.2 Consequences . . . 3
3.3 Comparison . . . 4

4 The saddle-point structure **4**
4.1 The gap equation . . . 4
4.2 The saddle-point Gaussian . . . 4
4.3 The oscillatory corrections . . . 4

5 Proof strategies **5**
5.1 Strategy A: NLSM (Kupiainen) 5
5.2 Strategy B: LSM saddle-point 5
5.3 Strategy C: LSM → NLSM equivalence 5

6 The $N = 2$ obstruction **5**

7 Current status **5**
7.1 What is proved (in Lean) 5
7.2 What is axiomatized 6
7.3 Open questions 6

1 The HS identity and the sign problem

1.1 The Euclidean quartic requires imaginary coupling

The O(N) LSM action on $\Lambda \subset \mathbb{Z}^2$:

$$S[\varphi] = \tfrac{1}{2} \sum_i \langle \varphi^i, (-\Delta)\varphi^i \rangle + N \sum_x \lambda \big(\tfrac{|\varphi(x)|^2}{N} - v^2 \big)^2. \tag{1}$$

The Boltzmann weight is e^{-S}. The quartic appears as $e^{-\lambda a^2}$ with $a = |\varphi|^2/N - v^2$. The HS identity:

$$e^{-\lambda a^2} = \frac{1}{\sqrt{4\pi\lambda}} \int_{-\infty}^{\infty} e^{-\sigma^2/(4\lambda) \, + \, i\sigma a} \, d\sigma. \tag{2}$$

Here $\sigma \in \mathbb{R}$ (real integration variable), $e^{-\sigma^2/(4\lambda)}$ ensures convergence, and the coupling $i\sigma a$ is **imaginary**. This is a Fourier transform (proved from Mathlib's `fourierIntegral_gaussian`).

The imaginary coupling is forced by the *minus* sign in $e^{-\lambda a^2}$ (repulsive/confining quartic in Euclidean signature). A real coupling would give $e^{+\lambda a^2}$ (attractive, wrong sign).

1.2 The σ-integral on the real axis

After applying (2) at each site and integrating out φ (Gaussian with complex mass), the partition function becomes:

$$Z = c \int_{\mathbb{R}^{|\Lambda|}} \det\big(-\Delta + 2i\sigma z\big)^{-N/2} \, e^{-\sum_x [\sigma(x)^2/(4\lambda) + i\sigma(x) v^2]} \, d^{|\Lambda|}\sigma, \tag{3}$$

where $z = N^{-1/2}$ and the det comes from the φ-integral.

This integral **converges** on the real σ-axis:

- $e^{-\sigma^2/(4\lambda)}$ provides Gaussian decay in σ.
- $|\det(-\Delta + 2i\sigma z)^{-N/2}| \le \det(-\Delta)^{-N/2}$ (bounded, since $|1 + i\theta| \ge 1$ for real θ).

But the integrand is **complex** (the det and the $e^{i\sigma v^2}$ factor oscillate). This is the sign problem.

2 The NLSM: Kupiainen's mass trick

2.1 Adding m_0^2 for free

For the NLSM with constraint $|\varphi|^2 = N\beta$: any function of $|\varphi|^2$ is constant on the constraint surface. In particular, $\frac{1}{2}m_0^2|\varphi|^2 = \frac{1}{2}m_0^2 N\beta = \text{const}$.

So we can **freely add** $\frac{1}{2}m_0^2|\varphi|^2$ to the action, giving the massive GFF:

$$S_{\text{NLSM}} = \tfrac{1}{2}\langle\varphi, (-\Delta + m_0^2)\varphi\rangle + \text{const}, \qquad |\varphi|^2 = N\beta. \tag{4}$$

The parameter $m_0^2 > 0$ is arbitrary; we choose it via the gap equation (Kupiainen eq. 9b): $(-\Delta + m_0^2)^{-1}_{ii} = \beta$.

2.2 The NLSM operator has real mass

After HS (introducing the Lagrange multiplier α for the constraint), the φ-propagator at fixed α is:

$$\left(-\Delta + m_0^2 - 2i\alpha z\right)^{-1}(x, 0). \tag{5}$$

The operator has **real part** $-\Delta + m_0^2 \geq m_0^2 > 0$. By the triangle inequality (FK representation):

$$\left|(-\Delta + m_0^2 - 2i\alpha z)^{-1}(x, 0)\right| \leq (-\Delta + m_0^2)^{-1}(x, 0) \sim e^{-m_0|x|}. \tag{6}$$

Every α-value gives exponential decay with mass m_0. The α-average is a correction, not the leading effect.

2.3 Kupiainen's proof structure

1. Gap equation: $m_0 > 0$ (always in 2d).
2. Bound (6): each α-value gives decay.
3. Cluster expansion + RP: control the α-average (prefactor, not the mass).
4. Mass gap: $m = m_0(1 + O(1/N))$.

The hard step is (3), but the mass m_0 comes for free from (2).

3 The LSM: the mass cancels

3.1 Add-subtract fails

For the LSM (no constraint), we try the same trick: add $\frac{1}{2}m_0^2|\varphi|^2$ to the action and subtract it from the interaction:

$$S = \left[\tfrac{1}{2}\langle\varphi, (-\Delta + m_0^2)\varphi\rangle\right] + \left[\lambda(u - v^2)^2 - \tfrac{1}{2}m_0^2 u\right], \qquad u = |\varphi|^2/N. \tag{7}$$

Apply HS to $\lambda(u-v^2)^2$ only. The subtracted mass $-\frac{1}{2}m_0^2 u$ remains outside the HS. Computing the φ-dependence:

$$\begin{aligned} &-\tfrac{1}{2}\langle\varphi,(-\Delta+m_0^2)\varphi\rangle + i\sigma|\varphi|^2 + \tfrac{1}{2}m_0^2|\varphi|^2 \\ &\quad = -\tfrac{1}{2}\langle\varphi,(-\Delta)\varphi\rangle + i\sigma|\varphi|^2 \\ &\quad = -\tfrac{1}{2}\langle\varphi,(-\Delta-2i\sigma)\varphi\rangle. \end{aligned} \tag{8}$$

The m_0^2 from the GFF and the $-m_0^2$ from the subtraction **cancel exactly**. The LSM operator is:

$$-\Delta - 2i\sigma \qquad (\textbf{no real mass}). \tag{9}$$

3.2 Consequences

The operator (9) has eigenvalues $k^2 - 2i\sigma$ with real part $k^2 \geq 0$ (from $-\Delta$) and imaginary part -2σ. For $k=0$: the eigenvalue is $-2i\sigma$ (purely imaginary).

- **No single-σ decay**: $|(-\Delta-2i\sigma)^{-1}(x,0)|$ does NOT decay exponentially for each σ-value. At $k=0$: $|1/(-2i\sigma)| = 1/(2|\sigma|)$, independent of $|x|$.
- **Mass from σ-average only**: The exponential decay of $\langle\varphi(x)\varphi(0)\rangle$ must come *entirely* from the oscillatory cancellations in the σ-integral (3).
- **The FK bound fails**: The NLSM bound (6) relies on the real mass m_0^2. Without it, $|G_\sigma| \leq G_0$ gives $G_0 = (-\Delta)^{-1}$ (massless propagator, no decay).

3.3 Comparison

	NLSM	LSM
Mass in operator	m_0^2 (free, from constraint)	**none** (cancels)
Operator after HS	$-\Delta + m_0^2 - 2i\alpha z$	$-\Delta - 2i\sigma$
Single-σ decay	$e^{-m_0\|x\|}$ (yes)	**none**
Mass gap source	Real mass + corrections	**σ-average only**
FK bound	Works ($\|G\| \leq G_{m_0}$)	**Fails**

4 The saddle-point structure

Despite the difference in operator structure, both models share the same saddle-point analysis.

4.1 The gap equation

On the real σ-axis, the saddle of the exponent in (3) is at $\sigma = 0$ (after absorbing the saddle value into m_0^2 for the NLSM, or into the gap equation for the LSM). The gap equation in both cases:

$$(-\Delta + m_0^2)^{-1}_{xx} = \beta_{\text{eff}} \tag{10}$$

where β_{eff} encodes the coupling. In 2d: $m_0 \sim \Lambda\, e^{-c/g^2}$ (exponentially small).

4.2 The saddle-point Gaussian

Near $\sigma = 0$: the exponent is approximately $-\frac{1}{2}(\sigma, H\sigma)$ where $H > 0$ (positive definite Hessian, including both the bare $1/(2\lambda)$ and the $\mathrm{Tr}\log$ contribution). The saddle-point Gaussian IS log-concave on the real axis.

4.3 The oscillatory corrections

Away from $\sigma = 0$: the exponent develops an imaginary part $\mathrm{Im}\, f(\sigma) \neq 0$, causing oscillations. The sign problem is the difficulty of controlling these oscillations.

Key scales (at the saddle):

- Width of the Gaussian: $\sigma \sim 1/\sqrt{N}$ per site.
- Oscillation onset: $\mathrm{Im}\, f \sim O(\sigma^3/\sqrt{N})$ (cubic, since $\mathrm{Im}\, f'' = 0$ at the saddle).
- Total oscillation at the saddle width: $\sim |\Lambda|/N^2$.

For $N \gg \sqrt{|\Lambda|}$: oscillations are negligible within the saddle-point width. The saddle-point approximation is accurate.

5 Proof strategies

5.1 Strategy A: NLSM (Kupiainen)

Work with the NLSM directly. The constraint gives real mass m_0^2 for free. The FK bound gives $|G| \leq G_{m_0}$ (single-σ decay). Cluster expansion + RP control the σ-average.

Advantage: Real mass in the operator, clean FK bound.
Disadvantage: Cluster expansion needed (technically heavy).
Threshold: $N > C\, m_0^{-a_2}$ with $a_2 = O(d^2)$ (Kupiainen Thm. 1).

5.2 Strategy B: LSM saddle-point

Work with the LSM. No real mass in the operator, but the saddle-point Gaussian on the real σ-axis IS log-concave.

Idea: Apply BL to the saddle-point Gaussian (log-concave, real, positive) and control the oscillatory corrections as a perturbation.

Challenge: The mass gap comes entirely from the σ-average (no single-σ decay). Need to show the oscillatory corrections don't destroy the saddle-point mass.

Advantage: Avoids cluster expansion (BL for the Gaussian).
Disadvantage: No FK bound; need new estimates for the oscillatory σ-average.

5.3 Strategy C: LSM $\to$ NLSM equivalence

Prove that the LSM and NLSM have the same IR physics (they differ only at the σ-mass scale $M_\sigma \sim \sqrt{\lambda}\, v$), then use Strategy A.

Advantage: Gets the NLSM mass gap (with the real mass trick) from the LSM formulation.
Disadvantage: Proving IR equivalence rigorously is itself a major result.

6 The $N = 2$ obstruction

For $N = 2$: the BKT transition occurs at finite λ (Dario-Garban [4]), driven by vortices ($\pi_1(S^1) = \mathbb{Z}$). Any proof must exclude $N = 2$. For $N \geq 3$: $\pi_1(S^{N-1}) = 0$, no BKT.

7 Current status

7.1 What is proved (in Lean)

- HS identity (2): from Mathlib's `fourierIntegral_gaussian`.
- Convergence on the real axis: $|e^{i\sigma a}| = 1$, $|e^{-\sigma^2/(4\lambda)}| \leq 1$.
- HS equivalence (one-site): $\int d\sigma\, e^{-S_{\text{HS}}} = c \cdot e^{-S_{\text{original}}}$ (1 sorry for complex arithmetic).
- Contour rotation algebra: $(i\sigma')^2 = -\sigma'^2$, $i \cdot a \cdot (i\sigma') = -a\sigma'$, etc.
- Continuum limit: OS0–OS2 for the O(N) LSM on T_L^2.

7.2 What is axiomatized

- Contour shift (Cauchy's theorem for rectangles): provable from Mathlib + PNT project.
- Matrix calculus (det C^∞, Jacobi's formula).
- FK sub-Gaussian bound (placeholder for the mass gap step).
- Continuum limit technicalities (4 axioms, porting from pphi2/gaussian-field).

7.3 Open questions

1. Can Strategy B work? Is BL on the saddle-point Gaussian sufficient to control the oscillatory corrections?
2. What is the threshold N_0 for each strategy?
3. Can the N=1 test case (P(φ)$_2$) validate Strategy B?

References

[1] H. J. Brascamp and E. H. Lieb, J. Funct. Anal. **22** (1976) 366–389.

[2] A. Kupiainen, Comm. Math. Phys. **73** (1980) 273–294.

[3] A. Kupiainen, Comm. Math. Phys. **74** (1980) 199–222.

[4] P. Dario and C. Garban, Comm. Math. Phys. (2025); arXiv:2311.16546.

[5] J. Glimm, A. Jaffe, and T. Spencer, Ann. Math. **100** (1974) 585–632.