


# Response Theory

Mike Reppert

September 28, 2022

## Previously on CHM676...

In homogeneous dielectric materials, the dynamics of  $\mathbf{E}$  and  $\mathbf{B}$  are determined by the polarization density  $\mathbf{P}$ .



**Today:** How does  $\mathbf{P}$  respond to the field?

# Outline for Today:

- 1 Physical Guidelines
- 2 Mathematical Framework
- 3 Symmetry and Invariance of Response Tensors

# Physical Guidelines

# Physical Guidelines

We study  $P$  as a *functional* of  $E$  and  $B$ :

$$P(\boldsymbol{x}, t) = P[\boldsymbol{E}(\boldsymbol{x}', t'), \boldsymbol{B}(\boldsymbol{x}', t')].$$

# Physical Guidelines

We study  $P$  as a *functional* of  $E$  and  $B$ :

$$P(\boldsymbol{x}, t) = P[\boldsymbol{E}(\boldsymbol{x}', t'), \boldsymbol{B}(\boldsymbol{x}', t')].$$

Physically, we expect:

- Response to  $B$  is negligible

# Physical Guidelines

We study  $P$  as a *functional* of  $E$  and  $B$ :

$$P(\boldsymbol{x}, t) = P[E(\boldsymbol{x}', t'), B(\boldsymbol{x}', t')].$$

Physically, we expect:

- Response to  $B$  is negligible
- Response is *local*:  $P(\boldsymbol{x})$  depends only on  $E(\boldsymbol{x})$ .

# Physical Guidelines

We study  $P$  as a *functional* of  $E$  and  $B$ :

$$P(\mathbf{x}, t) = P[E(\mathbf{x}', t'), B(\mathbf{x}', t')].$$

Physically, we expect:

- Response to  $B$  is negligible
- Response is *local*:  $P(\mathbf{x})$  depends only on  $E(\mathbf{x})$ .
- Response is *causal*:  $P(t)$  depends only on  $E(t' \leq t)$ .



# Physical Guidelines

We study  $P$  as a *functional* of  $E$  and  $B$ :

$$P(\boldsymbol{x}, t) = P[E(\boldsymbol{x}', t'), B(\boldsymbol{x}', t')].$$

Physically, we expect:

- Response to  $B$  is negligible
- Response is *local*:  $P(\boldsymbol{x})$  depends only on  $E(\boldsymbol{x})$ .
- Response is *causal*:  $P(t)$  depends only on  $E(t' \leq t)$ .
- Response is *stable*:

# Physical Guidelines

We study  $P$  as a *functional* of  $E$  and  $B$ :

$$P(\mathbf{x}, t) = P[E(\mathbf{x}', t'), B(\mathbf{x}', t')].$$

Physically, we expect:

- Response to  $B$  is negligible
- Response is *local*:  $P(\mathbf{x})$  depends only on  $E(\mathbf{x})$ .
- Response is *causal*:  $P(t)$  depends only on  $E(t' \leq t)$ .
- Response is *stable*:
  - Must exist a time scale  $\delta t$  below which  $P$  no longer cares about variations in  $E(t + \delta t)$

# Physical Guidelines

We study  $P$  as a *functional* of  $E$  and  $B$ :

$$P(x, t) = P[E(x', t'), B(x', t')].$$

Physically, we expect:

- Response to  $B$  is negligible
- Response is *local*:  $P(x)$  depends only on  $E(x)$ .
- Response is *causal*:  $P(t)$  depends only on  $E(t' \leq t)$ .
- Response is *stable*:
  - Must exist a time scale  $\delta t$  *below* which  $P$  no longer cares about variations in  $E(t + \delta t)$
  - Must exist a time scale  $T$  *beyond* which  $P$  doesn't remember  $E(t - T)$ .

# Take-Home Point

## **Physical** constraints:

- locality
- causality
- stability

strongly limit the possible forms for the **mathematical** dependence of  $P$  on  $E$ .


# Mathematical Framework

# Mathematical Framework

The *response theory* framework is essentially a Taylor series expansion for functionals.


# Mathematical Framework

The *response theory* framework is essentially a Taylor series expansion for functionals.



# Mathematical Framework

The *response theory* framework is essentially a Taylor series expansion for functionals.




$$f(x) \approx f(0)$$



# Mathematical Framework


The *response theory* framework is essentially a Taylor series expansion for functionals.



$$f(x) \approx f(0) + \left. \frac{df}{dx} \right|_{x=0} x$$

# Mathematical Framework


The *response theory* framework is essentially a Taylor series expansion for functionals.



$$f(x) \approx f(0) + \left. \frac{df}{dx} \right|_{x=0} x + \frac{1}{2} \left. \frac{d^2 f}{dx^2} \right|_{x=0} x^2$$

# Mathematical Framework

The *response theory* framework is essentially a Taylor series expansion for functionals.



$$f(x) \approx f(0) + \left. \frac{df}{dx} \right|_{x=0} x + \frac{1}{2} \left. \frac{d^2 f}{dx^2} \right|_{x=0} x^2 + \frac{1}{6} \left. \frac{d^3 f}{dx^3} \right|_{x=0} x^3 \dots$$

# Mathematical Framework

The Taylor series of a multi-variable function  $g(x_1, \dots, x_N)$  looks like:

$$\begin{aligned}
 g(x_1, \dots, x_N) = & g(0, \dots, 0) \\
 & + \frac{\partial g}{\partial x_1} \Big|_{\mathbf{x}=\mathbf{0}} x_1 + \frac{\partial g}{\partial x_2} \Big|_{\mathbf{x}=\mathbf{0}} x_2 + \dots + \frac{\partial g}{\partial x_N} \Big|_{\mathbf{x}=\mathbf{0}} x_N \\
 & + \frac{1}{2!} \frac{\partial^2 g}{\partial x_1^2} \Big|_{\mathbf{x}=\mathbf{0}} x_1^2 + \frac{\partial g}{\partial x_1 \partial x_2} \Big|_{\mathbf{x}=\mathbf{0}} x_1 x_2 + \dots + \frac{1}{2!} \frac{\partial^2 g}{\partial x_N^2} \Big|_{\mathbf{x}=\mathbf{0}} x_N^2 \\
 & + \dots
 \end{aligned}$$

What is the corresponding expansion for a functional like  $P[\mathbf{E}]$ ?

# Mathematical Framework

Since the response is stable, we can sample  $\mathbf{E}$  at a finite number of time points:

$$P_I(t) \approx f_I(E_x(t_0), E_y(t_0), E_z(t_0), E_x(t_1), \dots, E_z(t_N); t, \delta t, T).$$

Expanding in a Taylor series:

$$\begin{aligned} P_I(t) &\approx f_I(0, \dots, 0; t, \delta t, T) \\ &+ \frac{\partial f_I}{\partial E_x(t_0)} \Big|_{\mathbf{E}=0} E_x(t_0) + \dots + \frac{\partial f_I}{\partial E_z(t_N)} \Big|_{\mathbf{E}=0} E_z(t_N) \\ &+ \frac{1}{2!} \frac{\partial^2 f_I}{\partial [E_x(t_0)]^2} \Big|_{\mathbf{E}=0} [E_x(t_0)]^2 + \frac{\partial^2 f_I}{\partial E_x(t_0) \partial E_y(t_0)} \Big|_{\mathbf{E}=0} E_x(t_0) E_y(t_0) + \dots \end{aligned}$$

# Mathematical Framework

As our sampling points get closer together, the sums converge to integrals:

$$\begin{aligned}
 P(t) = & \sum_{n=0}^{\infty} \sum_{\alpha_1, \dots, \alpha_n} \int_{-\infty}^t dt_n \int_{-\infty}^{t_n} dt_{n-1} \dots \int_{-\infty}^{t_2} dt_1 \\
 & \times E_{\alpha_1}(t_1) E_{\alpha_2}(t_2) \dots E_{\alpha_n}(t_n) \\
 & \times R_{\alpha_1 \dots \alpha_n \alpha}^{(n)}(t, t_n, t_{n-1}, \dots, t_1)
 \end{aligned}$$

where  $R_{\alpha_1 \dots \alpha_n \alpha}^{(n)}(t, t_n, t_{n-1}, \dots, t_1)$  is the  $n^{\text{th}}$ -order *response function*\* – the target of  $n^{\text{th}}$ -order spectroscopies.

\*Almost. Actually  $R^{(n)}$  depends only on time *differences*. Stay tuned!

# Symmetry and Invariance of Response Tensors

# Time-translation Invariance

All systems we study will satisfy **time-translation invariance**: Only time *differences* matter!

$$R_{\alpha_1 \dots \alpha_n \alpha}^{(n)}(t, t_n, t_{n-1}, \dots, t_1) \Rightarrow R_{\alpha_1 \dots \alpha_n \alpha}^{(n)}(t - t_n, t_n - t_{n-1}, \dots, t_2 - t_1)$$



# Time-translation Invariance

All systems we study will satisfy **time-translation invariance**: Only time *differences* matter!

$$R_{\alpha_1 \dots \alpha_n \alpha}^{(n)}(t, t_n, t_{n-1}, \dots, t_1) \Rightarrow R_{\alpha_1 \dots \alpha_n \alpha}^{(n)}(t - t_n, t_n - t_{n-1}, \dots, t_2 - t_1)$$

Rearranging:

$$P_{\alpha}^{(n)}(t) = \sum_{\alpha_1, \dots, \alpha_n} \int_{-\infty}^{\infty} d\tau_n \dots \int_{-\infty}^{\infty} d\tau_1 R_{\alpha_1 \dots \alpha_n \alpha}^{(n)}(\tau_1, \dots, \tau_n) \\ \times E_{\alpha_1}(t - \tau_1 - \dots - \tau_n) E_{\alpha_2}(t - \tau_2 - \dots - \tau_n) \dots E_{\alpha_n}(t - \tau_n).$$

# Time-translation Invariance

All systems we study will satisfy **time-translation invariance**: Only time *differences* matter!

$$R_{\alpha_1 \dots \alpha_n \alpha}^{(n)}(t, t_n, t_{n-1}, \dots, t_1) \Rightarrow R_{\alpha_1 \dots \alpha_n \alpha}^{(n)}(t - t_n, t_n - t_{n-1}, \dots, t_2 - t_1)$$

Rearranging:

$$P_{\alpha}^{(n)}(t) = \sum_{\alpha_1, \dots, \alpha_n} \int_{-\infty}^{\infty} d\tau_n \dots \int_{-\infty}^{\infty} d\tau_1 R_{\alpha_1 \dots \alpha_n \alpha}^{(n)}(\tau_1, \dots, \tau_n) \\ \times E_{\alpha_1}(t - \tau_1 - \dots - \tau_n) E_{\alpha_2}(t - \tau_2 - \dots - \tau_n) \dots E_{\alpha_n}(t - \tau_n).$$

**Causality** dictates that  $R_{\alpha_1, \dots, \alpha_n, \alpha}^{(n)}$  is non-zero only for *positive time delays*.

# Spatial Symmetries

**Neumann's Principle:** Spatial symmetries of the material *must* be reflected in the response tensor.


This **dramatically** simplifies the analysis of nonlinear experiments!

# Spatial Symmetries

**Neumann's Principle:** Spatial symmetries of the material *must* be reflected in the response tensor.

This **dramatically** simplifies the analysis of nonlinear experiments!

**NB:** Only *macroscopic* symmetry is relevant!




[https://commons.wikimedia.org/wiki/File:  
A\\_Molecular\\_Dynamics\\_Simulation\\_of\\_Liquid\\_Water\\_at\\_298\\_K.webm](https://commons.wikimedia.org/wiki/File:A_Molecular_Dynamics_Simulation_of_Liquid_Water_at_298_K.webm)

# Spatial Symmetries

**Example:**  $R_{xy}^{(1)}$  in an isotropic sample


Suppose  $\mathbf{E}$  is polarized along the  $y$ -axis. What happens to  $P_x^{(1)}$  when we invert the  $y$ -axis?



# Spatial Symmetries

**Example:**  $R_{xy}^{(1)}$  in an isotropic sample

Suppose  $\mathbf{E}$  is polarized along the  $y$ -axis. What happens to  $P_x^{(1)}$  when we invert the  $y$ -axis?



**Nothing!**

# Spatial Symmetries

**Under  $y$ -axis inversion:**

- $y \rightarrow -y$
- $P_x^{(1)} \rightarrow P_x^{(1)}$
- $E_y \rightarrow -E_y$
- $R_{xy} \rightarrow R_{xy}$

# Spatial Symmetries

**Under  $y$ -axis inversion:**

- $y \rightarrow -y$
- $P_x^{(1)} \rightarrow P_x^{(1)}$
- $E_y \rightarrow -E_y$
- $R_{xy} \rightarrow R_{xy} \Leftarrow$  Neumann's Principle



# Spatial Symmetries

**Under  $y$ -axis inversion:**

- $y \rightarrow -y$
- $P_x^{(1)} \rightarrow P_x^{(1)}$
- $E_y \rightarrow -E_y$
- $R_{xy} \rightarrow R_{xy} \Leftarrow$  **Neumann's Principle**

But response theory says:

$$P_x^{(1)}(t) = \int_{-\infty}^{\infty} d\tau_1 R_{xy}^{(1)}(\tau_1) E_y(t - \tau_1) = -P_x^{(1)}(t).$$

# Spatial Symmetries

**Under  $y$ -axis inversion:**

- $y \rightarrow -y$
- $P_x^{(1)} \rightarrow P_x^{(1)}$
- $E_y \rightarrow -E_y$
- $R_{xy} \rightarrow R_{xy} \Leftarrow$  **Neumann's Principle**

But response theory says:

$$P_x^{(1)}(t) = \int_{-\infty}^{\infty} d\tau_1 R_{xy}^{(1)}(\tau_1) E_y(t - \tau_1) = -P_x^{(1)}(t).$$

The only possible conclusion is that  $R_{xy}^{(1)} = 0$ !

# Spatial Symmetries

**More generally:** In isotropic media

- All tensor elements with an odd number *of any index* vanish (e.g.,  $R_{xxxy}^{(3)} = 0$ )

# Spatial Symmetries

**More generally:** In isotropic media

- All tensor elements with an odd number *of any index* vanish (e.g.,  $R_{xxxy}^{(3)} = 0$ )
- Corollary: all even-order response functions vanish(!)

# Spatial Symmetries

**More generally:** In isotropic media

- All tensor elements with an odd number *of any index* vanish (e.g.,  $R_{xxxy}^{(3)} = 0$ )
- Corollary: all even-order response functions vanish(!)
- Response tensor elements are symmetry-related (e.g.,  $R_{xxyy}^{(3)} = R_{yyxx}^{(1)}$ )

# Even-Order Spectroscopies

**But remember: Not all materials are isotropic**

# Even-Order Spectroscopies

**But remember: Not all materials are isotropic**

Even-order spectroscopies are *specifically sensitive* to material boundaries

# Even-Order Spectroscopies

**But remember: Not all materials are isotropic**

Even-order spectroscopies are *specifically sensitive* to material boundaries  
⇒ Imaging!





# Even-Order Spectroscopies

**But remember: Not all materials are isotropic**

Even-order spectroscopies are *specifically sensitive* to material boundaries  
 $\Rightarrow$  Imaging!



**Garth Simpson**



# Take-Home Points

**Time-translation invariance** and **causality** dictate that response functions depend only on *positive time delays* between interactions.

**Spatial symmetries** in the material must be reflected in the response tensors.

In **isotropic media**:

- Response elements with unpaired axes vanish
- Surviving elements are symmetry-related
- Even-order spectroscopies are forbidden – hence useful for detecting defects