

# LOGIC AND THEORETICAL FOUNDATION OF COMPUTER SCIENCE

LATFoCS

---

Pamela Fleischmann

fpa@informatik.uni-kiel.de

Winter Semester 2019

Kiel University  
Dependable Systems Group



# SEMANTICS OF PROPOSITIONAL LOGIC

---

# Motivation

Let's think about the natural language and our understanding of it:



# Motivation

Let's think about the natural language and our understanding of it:

- I like bananas and chocolate.



Let's think about the natural language and our understanding of it:

- I like bananas and chocolate.
  - Decomposition into atomic propositions tells us that I like bananas and I like chocolate.



Let's think about the natural language and our understanding of it:

- I like bananas and chocolate.
  - Decomposition into atomic propositions tells us that I like bananas and I like chocolate.
- I don't like rain and being on the bike.



Let's think about the natural language and our understanding of it:

- I like bananas and chocolate.
  - Decomposition into atomic propositions tells us that I like bananas and I like chocolate.
- I don't like rain and being on the bike.
  - intuitively: being on a bike **during** rain is nothing what I like



Let's think about the natural language and our understanding of it:

- I like bananas and chocolate.
  - Decomposition into atomic propositions tells us that I like bananas and I like chocolate.
- I don't like rain and being on the bike.
  - intuitively: being on a bike **during** rain is nothing what I like
  - logic:  $\neg (\text{rain and bike})$  equals  $\neg \text{rain or } \neg \text{bike}$





Let's think about the natural language and our understanding of it:

- I like bananas and chocolate.
  - Decomposition into atomic propositions tells us that I like bananas and I like chocolate.
- I don't like rain and being on the bike.
  - intuitively: being on a bike **during** rain is nothing what I like
  - logic:  $\neg (\text{rain and bike})$  equals  $\neg \text{rain or } \neg \text{bike}$
  - if I just don't like rain at all, the formula is true



Let's think about the natural language and our understanding of it:

- I like bananas and chocolate.
  - Decomposition into atomic propositions tells us that I like bananas and I like chocolate.
- I don't like rain and being on the bike.
  - intuitively: being on a bike **during** rain is nothing what I like
  - logic:  $\neg (\text{rain and bike})$  equals  $\neg \text{rain or } \neg \text{bike}$
  - if I just don't like rain at all, the formula is true
- we have to formalise this!



## Definition

- The set of **truth values** is defined as  $\mathcal{T} = \{\text{true}, \text{false}\}$ .



# Truth values and Models

## Definition

- The set of **truth values** is defined as  $\mathcal{T} = \{\text{true}, \text{false}\}$ .
- A **valuation**, **interpretation**, or **model** of a formula  $\varphi$  is a total mapping  $\beta : A \rightarrow \mathcal{T}$



# Truth values and Models

## Definition

- The set of **truth values** is defined as  $\mathcal{T} = \{\text{true}, \text{false}\}$ .
- A **valuation**, **interpretation**, or **model** of a formula  $\varphi$  is a total mapping  $\beta : A \rightarrow \mathcal{T}$

other notations for true and false: 1 and 0, t and f



# Truth value of a Formula

## Definition

Let  $\varphi \in \Phi$  and  $\beta$  an interpretation. Then  $\hat{\beta}(\varphi)$  is inductively defined by

$$\hat{\beta}(p) = \beta(p) \text{ for atoms } p \in A$$

$$\hat{\beta}(\neg\psi) = \text{true iff } \hat{\beta}(\psi) = \text{false}$$

$$\hat{\beta}(\psi \wedge \chi) = \begin{cases} \text{true} & \text{if } \hat{\beta}(\psi) = \text{true and } \hat{\beta}(\chi) = \text{true} \\ \text{false} & \text{otherwise} \end{cases}$$

$$\hat{\beta}(\psi \vee \chi) = \begin{cases} \text{false} & \text{if } \hat{\beta}(\psi) = \text{false and } \hat{\beta}(\chi) = \text{false} \\ \text{true} & \text{otherwise} \end{cases}$$

$$\hat{\beta}(\psi \rightarrow \chi) = \begin{cases} \text{false} & \text{if } \hat{\beta}(\psi) = \text{true and } \hat{\beta}(\chi) = \text{false} \\ \text{true} & \text{otherwise} \end{cases}$$



# Truth value of a formula (Cont.)

## Definition

$$\hat{\beta}(\psi \leftrightarrow \chi) = \text{true} \text{ iff } \hat{\beta}(\psi) = \hat{\beta}(\chi)$$

$$\hat{\beta}(\psi \dot{\vee} \chi) = \text{true} \text{ iff } \hat{\beta}(\psi) \neq \hat{\beta}(\chi)$$

$$\hat{\beta}(\psi \downarrow \chi) = \text{true} \text{ iff } \hat{\beta}(\psi \vee \chi) = \text{false}$$

$$\hat{\beta}(\psi \uparrow \chi) = \text{true} \text{ iff } \hat{\beta}(\psi \wedge \chi) = \text{false}$$



## Definition

Let  $\varphi \in \Phi$  be a propositional formula with the atoms  $A_\varphi \subseteq A$ . A function  $\bar{\beta}$  is a partial interpretation of  $\varphi$  if its domain is a subset of  $A_\varphi$ .





# Partial Interpretation

## Definition

Let  $\varphi \in \Phi$  be a propositional formula with the atoms  $A_\varphi \subseteq A$ . A function  $\bar{\beta}$  is a partial interpretation of  $\varphi$  if its domain is a subset of  $A_\varphi$ .

For  $\varphi = p \vee q$  ( $p, q \in A$ ) we have  $\hat{\beta}(\varphi) = \text{true}$  if the interpretation of at least one atom is true. Thus with  $\bar{\beta}(p) = \text{true}$  we have  $\bar{\beta}(\varphi) = \text{true}$  although  $\bar{\beta}$  is not total on  $A_\varphi$ .



# Truth Tables - a convenient way to deal with truth

## Definition

A **truth table** for  $\varphi \in \Phi$  with the atoms  $A_\varphi \subseteq A$  and  $|A_\varphi| = n \in \mathbb{N}$  is a  $2^n \times (n+1)$ -matrix where the first  $n$  columns indicate one atom (assuming an (unimportant) order) and the last column indicates  $\varphi$ . Each row is associated with an interpretation of the atoms; the interpretation of the formula is stated in the last entry of a row.



# Some truth tables

$p$	$\neg p$	$p$	$q$	$p \vee q$	$p$	$q$	$p \dot{\vee} q$
0	1	0	0	0	0	0	0
0	1	0	1	1	0	1	1
1	0	1	0	1	1	0	1
		1	1	1	1	1	0

$p$	$q$	$p \wedge q$	$p$	$q$	$p \rightarrow q$	$p$	$q$	$p \leftrightarrow q$
0	0	0	0	0	1	0	0	1
0	1	0	0	1	1	0	1	0
1	0	0	1	0	0	1	0	0
1	1	1	1	1	1	1	1	1



# Example

$p$	$q$	$p \rightarrow q$	$\neg p$	$\neg p \vee q$	$(p \rightarrow q) \leftrightarrow (\neg p \vee q)$
0	0				
0	1				
1	0				
1	1				



# Example

$p$	$q$	$p \rightarrow q$	$\neg p$	$\neg p \vee q$	$(p \rightarrow q) \leftrightarrow (\neg p \vee q)$
0	0	1			
0	1	1			
1	0	0			
1	1	1			



# Example

$p$	$q$	$p \rightarrow q$	$\neg p$	$\neg p \vee q$	$(p \rightarrow q) \leftrightarrow (\neg p \vee q)$
0	0	1	1	1	
0	1	1	1	1	
1	0	0	0	0	
1	1	1	0	1	



# Example

$p$	$q$	$p \rightarrow q$	$\neg p$	$\neg p \vee q$	$(p \rightarrow q) \leftrightarrow (\neg p \vee q)$
0	0	1	1	1	1
0	1	1	1	1	1
1	0	0	0	0	1
1	1	1	0	1	1



# Example

$p$	$q$	$p \rightarrow q$	$\neg p$	$\neg p \vee q$	$(p \rightarrow q) \leftrightarrow (\neg p \vee q)$
0	0	1	1	1	1
0	1	1	1	1	1
1	0	0	0	0	1
1	1	1	0	1	1





# Example

$p$	$q$	$p \rightarrow q$	$\neg p$	$\neg p \vee q$	$(p \rightarrow q) \leftrightarrow (\neg p \vee q)$
0	0	1	1	1	1
0	1	1	1	1	1
1	0	0	0	0	1
1	1	1	0	1	1

- Each row stands for a different interpretation.
- Is the procedure for determining the interpretation of a formula efficient?



# Let's think about the operators: Disjunction

I buy milk or I buy bread.

- Do I lie if I buy only milk or only bread?
- Do I lie if I buy both?
- Do I lie if I buy nothing?



# Let's think about the operators: Disjunction

I buy milk or I buy bread.

- Do I lie if I buy only milk or only bread? No!
- Do I lie if I buy both? No!
- Do I lie if I buy nothing? Yes!



# Let's think about the operators: Disjunction

I buy milk or I buy bread.

- Do I lie if I buy only milk or only bread? No!
- Do I lie if I buy both? No!
- Do I lie if I buy nothing? Yes!

I buy either milk or I buy bread.

- Do I lie if I buy only milk or only bread? No!
- Do I lie if I buy both? Yes!
- Do I lie if I buy nothing? Yes!



# Let's think about the operators: Implication

If I buy milk, then I buy bread.

- Do I lie if I buy only milk?
- Do I lie if I buy only bread?
- Do I lie if I buy both?
- Do I lie if I buy nothing?



# Let's think about the operators: Implication

If I buy milk, then I buy bread.

- Do I lie if I buy only milk? **Yes!**
- Do I lie if I buy only bread? **No!**
- Do I lie if I buy both? **No!**
- Do I lie if I buy nothing? **No!**



# Let's think about the operators: Implication

If I buy milk, then I buy bread.

- Do I lie if I buy only milk?    **Yes!**
- Do I lie if I buy only bread?    **No!**
- Do I lie if I buy both?    **No!**
- Do I lie if I buy nothing?    **No!**

We defined **material implication**:

the **premise/antecedent** (LHS of  $\rightarrow$ ) does not cause necessarily  
the **conclusion** (RHS of  $\rightarrow$ )



# LOGICAL EQUIVALENCE

---



# And now?

We have syntax and semantic of propositional formulae - what have we gained?



# And now?

We have syntax and semantic of propositional formulae - what have we gained? let's have a look what we do in other artificial languages



# And now?

We have syntax and semantic of propositional formulae - what have we gained? let's have a look what we do in other artificial languages

```
f(a,b){  
  d=1  
  for i=2 to min{a,b}  
    if i|a and i|b  
      then d=i  
  return d  
}
```

```
g(a,b){  
  if b=0  
    then return a  
  else return g(b, a mod b)  
}
```



# And now?

We have syntax and semantic of propositional formulae - what have we gained? let's have a look what we do in other artificial languages

```
f(a,b){  
  d=1  
  for i=2 to min{a,b}  
    if i|a and i|b  
      then d=i  
  return d  
}  
  
g(a,b){  
  if b=0  
    then return a  
  else return g(b, a mod b)  
}
```

what are the programmes doing?



# And now?

We have syntax and semantic of propositional formulae - what have we gained? let's have a look what we do in other artificial languages

<pre>f(a,b){   d=1   for i=2 to min{a,b}     if i a and i b       then d=i   return d }</pre>	<pre>g(a,b){   if b=0     then return a     else return g(b, a mod b) }</pre>
---	---

what are the programmes doing?      the same



# Revisiting an Example

What can we deduce from this truth table?

$p$	$q$	$p \rightarrow q$	$\neg p$	$\neg p \vee q$	$(p \rightarrow q) \leftrightarrow (\neg p \vee q)$
0	0	1	1	1	1
0	1	1	1	1	1
1	0	0	0	0	1
1	1	1	0	1	1



## Definition

Two formula  $\varphi_1, \varphi_2 \in \Phi$  are called **logically equivalent** ( $\varphi_1 \equiv \varphi_2$ ) iff  $\hat{\beta}(\varphi_1) = \hat{\beta}(\varphi_2)$  for all interpretations  $\beta$ .



# How to prove logical equivalence?

For checking if  $\varphi_1, \varphi_2$  are logical equivalent, we have to check  $\hat{\beta}(\varphi_1) = \hat{\beta}(\varphi_2)$  for all interpretations  $\beta$ .





# How to prove logical equivalence?

For checking if  $\varphi_1, \varphi_2$  are logical equivalent, we have to check  $\hat{\beta}(\varphi_1) = \hat{\beta}(\varphi_2)$  for all interpretations  $\beta$ .

- Do we have to take every single interpretation and check the property? (exponentially many!)



# How to prove logical equivalence?

For checking if  $\varphi_1, \varphi_2$  are logical equivalent, we have to check  $\hat{\beta}(\varphi_1) = \hat{\beta}(\varphi_2)$  for all interpretations  $\beta$ .

- Do we have to take every single interpretation and check the property? (exponentially many!)
- What if we take an arbitrary one, having only the properties each interpretation has, and checking the equality?



# General Statements

For proving general statements, we use the method of  
representatives.



# General Statements

For proving general statements, we use the method of **representatives**.

- by the name representatives represent something



# General Statements

For proving general statements, we use the method of **representatives**.

- by the name representatives represent something
- notice: politicians are a bad example!



# General Statements

For proving general statements, we use the method of **representatives**.

- by the name representatives represent something
- notice: politicians are a bad example!
- better: among a group of objects or living beings find out all properties they have in common



# General Statements

For proving general statements, we use the method of **representatives**.

- by the name representatives represent something
- notice: politicians are a bad example!
- better: among a group of objects or living beings find out all properties they have in common
- then define a new object or living being (representative, prototype) having only these properties



# General Statements

For proving general statements, we use the method of **representatives**.

- by the name representatives represent something
- notice: politicians are a bad example!
- better: among a group of objects or living beings find out all properties they have in common
- then define a new object or living being (representative, prototype) having only these properties
- whatever statement you are claiming about the group, if the new object has it, the group will have it





# Commutativity of the Disjunction

## Lemma

*The operator  $\vee$  is commutative.*



# Commutativity of the Disjunction

## Lemma

*The operator  $\vee$  is commutative.*

### Proof.

- we have to show  $\varphi \vee \psi \equiv \psi \vee \varphi$  for all  $\varphi, \psi \in \Phi$ .



# Commutativity of the Disjunction

## Lemma

*The operator  $\vee$  is commutative.*

### Proof.

- we have to show  $\varphi \vee \psi \equiv \psi \vee \varphi$  for all  $\varphi, \psi \in \Phi$ .
- let  $\varphi, \psi \in \Phi$ ,  $\beta$  an arbitrary interpretation



# Commutativity of the Disjunction

## Lemma

*The operator  $\vee$  is commutative.*

### Proof.

- we have to show  $\varphi \vee \psi \equiv \psi \vee \varphi$  for all  $\varphi, \psi \in \Phi$ .
- let  $\varphi, \psi \in \Phi$ ,  $\beta$  an arbitrary interpretation
- by definition  $\hat{\beta}(\varphi \vee \psi) = \text{false}$  iff  $\hat{\beta}(\varphi) = \text{false}$  and  $\hat{\beta}(\psi) = \text{false}$



# Commutativity of the Disjunction

## Lemma

*The operator  $\vee$  is commutative.*

### Proof.

- we have to show  $\varphi \vee \psi \equiv \psi \vee \varphi$  for all  $\varphi, \psi \in \Phi$ .
- let  $\varphi, \psi \in \Phi$ ,  $\beta$  an arbitrary interpretation
- by definition  $\hat{\beta}(\varphi \vee \psi) = \text{false}$  iff  $\hat{\beta}(\varphi) = \text{false}$  and  $\hat{\beta}(\psi) = \text{false}$
- this is equivalent to  $\hat{\beta}(\psi) = \text{false}$  and  $\hat{\beta}(\varphi) = \text{false}$



# Commutativity of the Disjunction

## Lemma

*The operator  $\vee$  is commutative.*

### Proof.

- we have to show  $\varphi \vee \psi \equiv \psi \vee \varphi$  for all  $\varphi, \psi \in \Phi$ .
- let  $\varphi, \psi \in \Phi$ ,  $\beta$  an arbitrary interpretation
- by definition  $\hat{\beta}(\varphi \vee \psi) = \text{false}$  iff  $\hat{\beta}(\varphi) = \text{false}$  and  $\hat{\beta}(\psi) = \text{false}$
- this is equivalent to  $\hat{\beta}(\psi) = \text{false}$  and  $\hat{\beta}(\varphi) = \text{false}$
- by definition this is equivalent to  $\hat{\beta}(\psi \vee \varphi) = \text{false}$



# Commutativity of the Disjunction

## Lemma

*The operator  $\vee$  is commutative.*

### Proof.

- we have to show  $\varphi \vee \psi \equiv \psi \vee \varphi$  for all  $\varphi, \psi \in \Phi$ .
- let  $\varphi, \psi \in \Phi$ ,  $\beta$  an arbitrary interpretation
- by definition  $\hat{\beta}(\varphi \vee \psi) = \text{false}$  iff  $\hat{\beta}(\varphi) = \text{false}$  and  $\hat{\beta}(\psi) = \text{false}$
- this is equivalent to  $\hat{\beta}(\psi) = \text{false}$  and  $\hat{\beta}(\varphi) = \text{false}$
- by definition this is equivalent to  $\hat{\beta}(\psi \vee \varphi) = \text{false}$
- since all steps are equivalent we have proven  $\hat{\beta}(\varphi \vee \psi) = \hat{\beta}(\psi \vee \varphi)$



# The confusion with metalanguages

- in the last proof we used the word *equivalent* for stating that two claims imply each other

$$\begin{aligned} &(\hat{\beta}(\varphi) = \text{false and } \hat{\beta}(\psi) = \text{false}) \text{ is equivalent to} \\ &(\hat{\beta}(\psi) = \text{false and } \hat{\beta}(\varphi) = \text{false}) \end{aligned}$$





# The confusion with metalanguages

- in the last proof we used the word *equivalent* for stating that two claims imply each other

$$(\hat{\beta}(\varphi) = \text{false and } \hat{\beta}(\psi) = \text{false}) \text{ is equivalent to } \\ (\hat{\beta}(\psi) = \text{false and } \hat{\beta}(\varphi) = \text{false})$$

- don't confuse this with the logical operator  $\leftrightarrow$



# The confusion with metalanguages

- in the last proof we used the word *equivalent* for stating that two claims imply each other
$$(\hat{\beta}(\varphi) = \text{false and } \hat{\beta}(\psi) = \text{false}) \text{ is equivalent to } (\hat{\beta}(\psi) = \text{false and } \hat{\beta}(\varphi) = \text{false})$$
- don't confuse this with the logical operator  $\leftrightarrow$
- a metalanguage is a language  $L_2$  for talking about a language  $L_1$ : verb, noun, format-free etc.



# The confusion with metalanguages

- in the last proof we used the word *equivalent* for stating that two claims imply each other

$$(\hat{\beta}(\varphi) = \text{false and } \hat{\beta}(\psi) = \text{false}) \text{ is equivalent to } \\ (\hat{\beta}(\psi) = \text{false and } \hat{\beta}(\varphi) = \text{false})$$

- don't confuse this with the logical operator  $\leftrightarrow$
- a metalanguage is a language  $L_2$  for talking about a language  $L_1$ : verb, noun, format-free etc.
- if we want to talk about formulae (like comparing) we need a language for that



# The confusion with metalanguages

- in the last proof we used the word *equivalent* for stating that two claims imply each other
$$(\hat{\beta}(\varphi) = \text{false and } \hat{\beta}(\psi) = \text{false}) \text{ is equivalent to } (\hat{\beta}(\psi) = \text{false and } \hat{\beta}(\varphi) = \text{false})$$
- don't confuse this with the logical operator  $\leftrightarrow$
- a metalanguage is a language  $L_2$  for talking about a language  $L_1$ : verb, noun, format-free etc.
- if we want to talk about formulae (like comparing) we need a language for that
- we don't have necessarily different words (bad habit but reality)



# The confusion with metalanguages

- in the last proof we used the word *equivalent* for stating that two claims imply each other

$$(\hat{\beta}(\varphi) = \text{false and } \hat{\beta}(\psi) = \text{false}) \text{ is equivalent to } (\hat{\beta}(\psi) = \text{false and } \hat{\beta}(\varphi) = \text{false})$$

- don't confuse this with the logical operator  $\leftrightarrow$
- a metalanguage is a language  $L_2$  for talking about a language  $L_1$ : verb, noun, format-free etc.
- if we want to talk about formulae (like comparing) we need a language for that
- we don't have necessarily different words (bad habit but reality)
- equivalence in our logic language is  $\leftrightarrow$  (in  $\Phi$ )



# The confusion with metalanguages

- in the last proof we used the word *equivalent* for stating that two claims imply each other

$$(\hat{\beta}(\varphi) = \text{false and } \hat{\beta}(\psi) = \text{false}) \text{ is equivalent to } (\hat{\beta}(\psi) = \text{false and } \hat{\beta}(\varphi) = \text{false})$$

- don't confuse this with the logical operator  $\leftrightarrow$
- a metalanguage is a language  $L_2$  for talking about a language  $L_1$ : verb, noun, format-free etc.
- if we want to talk about formulae (like comparing) we need a language for that
- we don't have necessarily different words (bad habit but reality)
- equivalence in our logic language is  $\leftrightarrow$  (in  $\Phi$ )
- equivalence in the metalanguage is  $\equiv$  (talking about  $\Phi$ )



# Why do we use twice the same word?

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*



# Why do we use twice the same word?

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*

Before we can prove the theorem, we need a little bit of work and two definitions for convenience.





# Why do we use twice the same word?

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*

Before we can prove the theorem, we need a little bit of work and two definitions for convenience.

## Definition

A formula  $\varphi$  that is true for all interpretations  $\beta$  is called a **tautology**; if it is false for all interpretations  $\beta$  it is called a **contradiction**.



# How to prove *iff* statements?

**iff** means *if and only if* and contains **two** statements



# How to prove *iff* statements?

**iff** means *if and only if* and contains **two** statements

I buy milk if and only if I buy bread.



# How to prove *iff* statements?

**iff** means *if and only if* and contains **two** statements

I buy milk if and only if I buy bread.

- *I buy milk if I buy bread* is only true if I buy always milk when I buy bread.



# How to prove *iff* statements?

**iff** means *if and only if* and contains **two** statements

I buy milk if and only if I buy bread.

- *I buy milk if I buy bread* is only true if I buy always milk when I buy bread.
- but we know that the statement is also true if I don't buy bread, but milk.



# How to prove *iff* statements?

**iff** means *if and only if* and contains **two** statements

I buy milk if and only if I buy bread.

- *I buy milk if I buy bread* is only true if I buy always milk when I buy bread.
- but we know that the statement is also true if I don't buy bread, but milk.
- the *and only if* excludes these other options!



# How to prove *iff* statements?

**iff** means *if and only if* and contains **two** statements

I buy milk if and only if I buy bread.

- *I buy milk if I buy bread* is only true if I buy always milk when I buy bread.
- but we know that the statement is also true if I don't buy bread, but milk.
- the *and only if* excludes these other options!
- I buy milk if I buy bread, and I do so exactly in the cases when I by bread - never else!



# How to prove *iff* statements?

**iff** means *if and only if* and contains **two** statements

I buy milk if and only if I buy bread.

- *I buy milk if I buy bread* is only true if I buy always milk when I buy bread.
- but we know that the statement is also true if I don't buy bread, but milk.
- the *and only if* excludes these other options!
- I buy milk if I buy bread, and I do so exactly in the cases when I buy bread - never else!
- thus I buy milk if I buy bread, and if I buy milk, I buy bread





# Proving an Implication

- we have reduced the proof of an equivalence to two proofs of implications



# Proving an Implication

- we have reduced the proof of an equivalence to two proofs of implications
- how to prove an implication?



# Proving an Implication

- we have reduced the proof of an equivalence to two proofs of implications
- how to prove an implication?
- by definition: an implication is true if the conclusion is true in the case that the premise is true



# Proving an Implication

- we have reduced the proof of an equivalence to two proofs of implications
- how to prove an implication?
- by definition: an implication is true if the conclusion is true in the case that the premise is true
- we do not have to consider the case when the premise is false!



# Proving an Implication

- we have reduced the proof of an equivalence to two proofs of implications
- how to prove an implication?
- by definition: an implication is true if the conclusion is true in the case that the premise is true
- we do not have to consider the case when the premise is false!
- we can assume that the premise is true and then we prove that the conclusion is also true



# Proof of the Theorem

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*



# Proof of the Theorem

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*

**Proof:**



# Proof of the Theorem

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*

## Proof:

- we have to prove the statements
  1.  $\varphi_1 \equiv \varphi_2$  if  $\varphi_1 \leftrightarrow \varphi_2$
  2.  $\varphi_1 \leftrightarrow \varphi_2$  if  $\varphi_1 \equiv \varphi_2$





# Proof of the Theorem

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*

## Proof:

- we have to prove the statements
  1.  $\varphi_1 \equiv \varphi_2$  if  $\varphi_1 \leftrightarrow \varphi_2$
  2.  $\varphi_1 \leftrightarrow \varphi_2$  if  $\varphi_1 \equiv \varphi_2$
- ad 1. Assume  $\varphi_1 \leftrightarrow \varphi_2$ .



# Proof of the Theorem

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*

## Proof:

- we have to prove the statements
  1.  $\varphi_1 \equiv \varphi_2$  if  $\varphi_1 \leftrightarrow \varphi_2$
  2.  $\varphi_1 \leftrightarrow \varphi_2$  if  $\varphi_1 \equiv \varphi_2$
- ad 1. Assume  $\varphi_1 \leftrightarrow \varphi_2$ .
  - by definition of  $\leftrightarrow$ :  $\hat{\beta}(\varphi_1) = \hat{\beta}(\varphi_2)$  for all interpretations  $\beta$ .



# Proof of the Theorem

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*

### Proof:

- we have to prove the statements
  1.  $\varphi_1 \equiv \varphi_2$  if  $\varphi_1 \leftrightarrow \varphi_2$
  2.  $\varphi_1 \leftrightarrow \varphi_2$  if  $\varphi_1 \equiv \varphi_2$
- ad 1. Assume  $\varphi_1 \leftrightarrow \varphi_2$ .
  - by definition of  $\leftrightarrow$ :  $\hat{\beta}(\varphi_1) = \hat{\beta}(\varphi_2)$  for all interpretations  $\beta$ .
  - by definition of  $\equiv$ :  $\hat{\beta}(\varphi_1) = \hat{\beta}(\varphi_2)$  for all interpretations  $\beta$  implies  $\varphi_1 \equiv \varphi_2$



# Proof of the Theorem

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*

## Proof:

- we have to prove the statements
  1.  $\varphi_1 \equiv \varphi_2$  if  $\varphi_1 \leftrightarrow \varphi_2$
  2.  $\varphi_1 \leftrightarrow \varphi_2$  if  $\varphi_1 \equiv \varphi_2$
- ad 2. Assume  $\varphi_1 \equiv \varphi_2$ .



# Proof of the Theorem

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*

## Proof:

- we have to prove the statements
  1.  $\varphi_1 \equiv \varphi_2$  if  $\varphi_1 \leftrightarrow \varphi_2$
  2.  $\varphi_1 \leftrightarrow \varphi_2$  if  $\varphi_1 \equiv \varphi_2$
- ad 2. Assume  $\varphi_1 \equiv \varphi_2$ .
- by definition of  $\equiv$ :  $\hat{\beta}(\varphi_1) = \hat{\beta}(\varphi_2)$  for all interpretations  $\beta$



# Proof of the Theorem

## Theorem

*Let  $\varphi_1, \varphi_2 \in \Phi$ . Then  $\varphi_1 \equiv \varphi_2$  iff  $\varphi_1 \leftrightarrow \varphi_2$  is true for all interpretations  $\beta$ .*

## Proof:

- we have to prove the statements
  1.  $\varphi_1 \equiv \varphi_2$  if  $\varphi_1 \leftrightarrow \varphi_2$
  2.  $\varphi_1 \leftrightarrow \varphi_2$  if  $\varphi_1 \equiv \varphi_2$
- ad 2. Assume  $\varphi_1 \equiv \varphi_2$ .
- by definition of  $\equiv$ :  $\hat{\beta}(\varphi_1) = \hat{\beta}(\varphi_2)$  for all interpretations  $\beta$
- by definition of  $\leftrightarrow$ :  $\hat{\beta}(\varphi_1) = \hat{\beta}(\varphi_2)$  for all interpretations  $\beta$  implies  $\varphi_1 \leftrightarrow \varphi_2$ .

