Maryland Department of Health
Larry Hogan, Governor – Boyd Rutherford, Lt. Governor – Dennis R. Schrader, Secretary

June 2021

Unintentional Drug- and Alcohol-Related Intoxication Deaths in Maryland, 2020




Table of Contents

I. Methods **1**

II. **Summary of trends in deaths** **6**

III. **Figures** 10

A. Total intoxication deaths 11

B. Deaths by selected substances 15

C. Opioid-related deaths 16

1. Fentanyl-related 18

2. Heroin-related 21

3. Prescription Opioid-related 24

4. Carfentanil 29

D. Cocaine-related deaths 30

E. Benzodiazepine-related deaths 34

F. Phencyclidine-related deaths 38

G. Methamphetamine-related deaths 42

H. Alcohol-related deaths 46

I. Drug combinations 50

J. Age-adjusted death rates 56

IV. Tables **57**

Table 1. Total Number of Drug- and Alcohol-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 58

Table 2. Number of Opioid-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 59

Table 3. Number of Heroin-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 60

Table 4. Number of Prescription Opioid-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 61

Table 5. Number of Oxycodone-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 62

Table 6. Number of Methadone-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 63

Table 7. Number of Fentanyl-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 64

Table 8. Number of Cocaine-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 65

Table 9. Number of Benzodiazepine-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 66

Table 10. Number of Phencyclidine-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 67

Table 11. Number of Methamphetamine-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020.....68

Table 12. Number of Alcohol-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020 69

METHODS

Introduction

The purpose of this report is to describe trends in the number of unintentional drug- and alcohol-related intoxication deaths occurring in Maryland during the period 2011-2020. Trends are examined by age at time of death, race/ethnicity, gender, place of death, and substances related to death.

This report was prepared using drug and alcohol intoxication data housed in a registry developed and maintained by the Vital Statistics Administration (VSA) of the Maryland Department of Health (MDH). The methodology for reporting on drug-related intoxication deaths in Maryland was developed by VSA with assistance from the MDH Behavioral Health Administration, the Office of the Chief Medical Examiner (OCME) and the Maryland Poison Control Center. Assistance was also provided by authors of a Baltimore City Health Department report on intoxication deaths.[1]

Sources of data

The data included in this report were obtained mainly from the OCME. Maryland law requires the OCME to investigate all drug deaths occurring in the State, as well as non-natural and unattended deaths. In these instances, information compiled during an investigation is used to determine the cause or causes of death. Depending on the circumstances, an investigation may involve a combination of scene examination, review of witness reports, review of medical and police reports, autopsy, and toxicological analysis of autopsy specimens. Toxicological analysis is routinely performed when there is suspicion that a death was the result of drug or alcohol intoxication.

A small number of death records involving intoxication deaths were filed by sources other than OCME and were identified through death records maintained by VSA. This included records filed by medical facilities rather than OCME, and records filed by federal investigators following deaths involving U.S. military personnel. Information available on these cases was included in the registry.

Information on place of death and race/ethnicity was missing for a small number of records provided by OCME and was obtained through death certificate data. Death certificate data were also used to update demographic information on records that were amended after the records were filed with the Division of Vital Records.

[1] Office of Epidemiology and Planning, Baltimore City Health Department. Intoxication Deaths Associated with Drugs of Abuse or Alcohol. Baltimore City, Maryland: Baltimore City Health Department. January 2007.

Identification of drug-related intoxication deaths

For the purpose of this report, an intoxication death was defined as a death that was the result of recent ingestion or exposure to alcohol or another type of drug, including heroin, fentanyl, cocaine, prescription opioids, benzodiazepines, phencyclidine (PCP), methamphetamines, and other prescribed and unprescribed drugs. OCME provided all records to VSA for which the text of the cause of death included one or more of the following terms: poisoning, intoxication, toxicity, inhalation, ingestion, overdose, exposure, chemical, effects, or use. Any records provided by OCME that were not unintentional drug-related intoxication deaths, such as deaths due to smoke inhalation, carbon monoxide intoxication, cold exposure, and chronic use of alcohol or other drugs, were excluded in the registry. Also excluded from the registry were deaths for which the manner of death was determined to be natural, suicide, or homicide.

Analyses

Trends in the number of unintentional drug- and alcohol-related intoxication deaths occurring in Maryland during the years 2011-2020 were analyzed by age group, race/ethnicity, gender, place of occurrence of death, and substances related to the death. Changes over time were examined for deaths related to the following substances:

1. Opioids
    a. Heroin
    b. Prescription opioids
    c. Fentanyl (prescribed and illicit)
2. Cocaine
3. Benzodiazepines and related drugs
4. Phencyclidine
5. Methamphetamine
6. Alcohol

The number of deaths by place of occurrence was computed by jurisdiction and by region, categorized as follows:

Northwest Area	Baltimore Metro Area	National Capital Area	Southern Area	Eastern Shore Area
Garrett Co. Allegany Co. Washington Co. Frederick Co.	Baltimore City Baltimore Co. Anne Arundel Co. Carroll Co. Howard Co. Harford Co.	Montgomery Co. Prince George's Co.	Calvert Co. Charles Co. St. Mary's Co.	Cecil Co. Kent Co. Queen Anne's Co. Caroline Co. Talbot Co. Dorchester Co. Wicomico Co. Somerset Co. Worcester Co.

Trends in deaths for the period 2011-2020 are shown in Figures 1 through 42. Data on intoxication deaths related to a combination of substances are shown in Figures 43 through 49. Counts of the number of total deaths and deaths related to classes of substances or specific substances by place of occurrence are shown in Tables 1 through 12.

Age-adjusted death rates

Age-adjusted death rates by place of residence are shown in Figure 50. Age-adjusted death rates were calculated in order to allow for the comparison of drug death rates among Maryland jurisdictions. Unlike all other data included in this report, these rates are based on place of residence of the decedent rather than place where the drug-related incident occurred. Since out-of-state data are generally not available until approximately six months after the close of a calendar year, only data through 2019 were available at the time this report was prepared. Therefore, age-adjusted rates cover the period 2017 through 2019. Since the number of drug deaths is relatively small in many Maryland jurisdictions, it was necessary to calculate rates for a three year period in order to obtain counts that were large enough to be used to calculate stable rates.

Drug death information received from other states is far less detailed than the data available from OCME and often does not include information on the substances involved in a death. For that reason, rates could only be calculated for total deaths and not deaths related to individual substances.

Since an intoxication death may involve more than one substance, counts of deaths related to specific substances do not sum to the total number of deaths in this report.*

Opioid-related deaths

Opioids include heroin and prescription opioid drugs such as oxycodone, hydrocodone, hydromorphone, methadone, tramadol and codeine, and prescribed and illicit fentanyl. In this report, an opioid was considered to be associated with a death if a specific opioid drug was indicated in the cause of death. If the cause of death did not identify a specific drug (e.g., the cause of death indicated “Narcotic Intoxication”), OCME toxicology results were reviewed to determine whether the presence of any opioid drug was detected. If so, the cause of death was considered to be opioid-related, regardless of the level of the drug. Scene investigation notes were also reviewed in an attempt to better categorize death records with non-specific causes of death.

Since heroin is rapidly metabolized into morphine, the records of many deaths that are likely to be heroin-related do not list “heroin” as a cause of death, and therefore cannot be identified using only information listed in the cause of death. Therefore, a combination of information contained in the cause of death field, toxicology results, and scene investigation

notes is used to identify heroin-related deaths. In this report, a death was considered to be heroin-related if:

1. "Heroin" was mentioned in the cause of death; or
2. The toxicology screen showed a positive result for 6-monacetylmorphine; or
3. The toxicology screen showed positive results for both morphine and quinine; or
4. The cause of death was nonspecific and the scene investigation notes indicated that heroin was likely to have been involved in the death; or
5. The death was associated with morphine through either cause of death information or toxicology results, unless information contained in the investigation notes did not support this assumption.

A record was not coded as heroin-related, despite the presence of morphine, if OCME determined that another substance caused the death.

Prescription opioid-related deaths were defined as deaths that involve one or more prescription opioids, as identified through cause of death information when a specific drug was indicated and through toxicology results when the cause of death was nonspecific. Prescription opioids include buprenorphine, codeine, hydrocodone, hydromorphone, meperidine, methadone, morphine, oxycodone, pentazocine, propoxyphene, tramadol and prescribed fentanyl. Prescribed fentanyl is an opioid analgesic approved for patient use to manage severe or chronic pain. There are also forms of fentanyl that are produced illicitly in clandestine laboratories and mixed with (or substituted for) heroin or other illicit drugs. Although in some cases it was difficult to determine whether a prescribed or illicit form of fentanyl was related to a death, the count of prescription opioid-related drugs in this report includes only fentanyl deaths in which a prescription form of the drug was clearly involved.

Fentanyl-related deaths began increasing in late 2013 as a result of overdoses involving nonpharmaceutical fentanyl, that is, nonprescription fentanyl produced in clandestine laboratories and mixed with, or substituted for, heroin or other illicit substances. Nearly all fentanyl-related deaths occurring in recent years have involved the use of nonpharmaceutical fentanyl. Fentanyl is many times more potent than heroin, and greatly increases the risk of an overdose death. Carfentanil, an extremely potent analog of fentanyl, was first detected in Maryland drug intoxication death cases in 2017, and is reported separately in Figures 21 and 22.

Cocaine-related deaths

Cocaine is a highly addictive stimulant drug derived from coca leaves. It is frequently mixed with other non-psychoactive substances, such as cornstarch or talcum powder, to dilute its potency, however in the last few years, it has been mixed with fentanyl.

Benzodiazepine-related deaths

Benzodiazepines are a class of depressants that include drugs such as alprazolam, clonazepam, diazepam, and multiple related drugs. The category of benzodiazepine-related drugs in this report includes both benzodiazepines and related drugs, such as zolpidem, which have similar sedative effects.

Phencyclidine-related deaths

Phencyclidine, or phenylcyclohexyl piperidine (PCP), is an illicit hallucinogenic drug that can induce acute psychosis and aggressive behaviors. In the last few years it has been mixed with fentanyl.

Methamphetamine-related deaths

Methamphetamine is another highly addictive stimulant drug. Illicit forms of methamphetamine have also been found to be mixed with fentanyl or other opioids.

SUMMARY OF TRENDS IN DRUG INTOXICATION DEATHS — 2011 TO 2020

Total drug and alcohol intoxication deaths

- The number of drug- and alcohol-related intoxication deaths occurring in Maryland increased in 2020, after a slight decrease seen in 2019. The total number of deaths in 2020 was 2,799, which represented an 18% increase from the number of deaths (2,379) in 2019.
- Between the years 2011 through 2016, intoxication deaths increased among all age groups, and were highest among those aged 45-54 years old. The number of deaths among those aged <25 years and those 25-34 started to decline through 2019. In 2020, the older age groups saw the steepest increases in deaths, with those aged 55 years an older having the highest number of deaths each year since 2018. Between 2019 and 2020, deaths increased among those aged 45-54 years by 23% and increased by 20% among those aged 55 years and over.
- The number of deaths among non-Hispanic Whites decreased in 2019 but increased in 2020 by 16%. Deaths have increased steadily among non-Hispanic Blacks since 2012, and increased 16% between 2019 and 2020. Among Hispanics, deaths increased from 75 in 2019 to 126 in 2020.
- Deaths increased by 17% among men between 2019 and 2020, after slight decrease between 2018 and 2019. Deaths among women increased 19% between 2019 and 2020 after a three-year plateau. Intoxication deaths were 2.6 times higher among men than women in 2020.
- Two counties had no change in the number of deaths occurring between 2019 and 2020: St. Mary's and Frederick. Although there were decreases in the number of deaths occurring in five jurisdictions of the state between 2019 and 2020: Calvert, Carroll, Garrett, Harford, and Kent, the remainder saw increases: Allegany, Anne Arundel, Baltimore City, Baltimore County, Caroline, Cecil, Charles, Dorchester, Howard, Montgomery, Prince George's, Queen Anne's, Somerset, Talbot, Washington, Wicomico, and Worcester.

Opioid-related deaths

- Ninety percent of all intoxication deaths that occurred in Maryland in 2020 were **opioid**-related. **Opioid**-related deaths include deaths related to **heroin**, **prescription opioids**, and nonpharmaceutical **fentanyl**.
- The number of **opioid**-related deaths increased by 20% between 2019 and 2020, following a 2% decrease the previous year. Non opioid-related drug deaths increased slightly in 2020.
- **Fentanyl**-related deaths continued to drive opioid-related deaths. Between 2019 and 2020 the number of **fentanyl**-related deaths increased by 22% (from 1927 to 2342). The number of **heroin**-related deaths declined for the fourth straight year, decreasing by 55% between 2016 and 2020 to 548 deaths. The number of **prescription opioid**-related deaths increased by 23% between 2019 and 2020, following a three-year decrease.

Fentanyl

- **Fentanyl**-related deaths have increased rapidly since 2013, with a 229% increase between 2015 and 2016. Deaths related to **fentanyl** increased sharply again in 2020, rising 22% to a 10 year high of 2,342 deaths.
- In 2020, **Fentanyl**-related deaths rose among all age groups, with the highest increases among those 25-34 years (25%) and among those 55 and older (28%).
- **Fentanyl**-related deaths increased among non-Hispanic Whites (19%) and non-Hispanic Blacks (20%) between 2019 and 2020. The number of deaths among Hispanics nearly doubled, increasing 96% from 2019.
- **Fentanyl**-related deaths increased by 20% among men and by 26% among women in 2020.
- In 2020, **fentanyl** deaths increased in 18 jurisdictions, declined in 4 jurisdictions, and remained the same in 2 jurisdictions.
- Thirty-six percent of **fentanyl**-related deaths in 2020 occurred in combination with cocaine, 22% in combination with **heroin**, 18% in combination with **alcohol,** and 13% in combination with **prescription opioids**.
- Deaths related to **carfentanil** (a **fentanyl** analog) were first identified in 2017 (testing began in 2016). There were 60 **carfentanil**-related deaths in 2017, however this number dropped to 2 deaths in 2018 and 2019. There were no carfentanil deaths in 2020.

Heroin

- **Heroin**-related deaths continued to decrease in 2020, declining by 25% since 2019.
- There was a decrease in **Heroin**-related deaths in 2020 among all age groups.
- Deaths among both sexes continued to decline for the fourth consecutive year, falling 24% among men and 26% among women.
- **Heroin**-related deaths declined among non-Hispanic Whites (27%), non-Hispanic Blacks (21%) and Hispanics (20%) in 2020.
- In 2019, **heroin** deaths declined in 16 jurisdictions, remained the same in 3 jurisdictions, and increased in 5 jurisdictions.
- Ninety-four percent of **heroin**-related deaths in 2020 occurred in combination with **fentanyl,** 38% in combination with **cocaine**, 19% in combination with **prescription opioids**, and 12% in combination with **alcohol**.

Prescription Opioids

- The number of **prescription opioid**-related deaths had been rising since 2013, but started to decline 2017. After a three-year decrease, deaths increased sharply in 2020 rising by 23% compared with 2019.
- In 2020, the number of **prescription opioid**-related deaths rose in all age groups.
- **Prescription opioid**-related deaths increased among non-Hispanic Whites (18%), non-Hispanic Blacks (31%) from 2019 to 2020. The number of deaths nearly tripled for Hispanics.
- Deaths related to **prescription opioids** increased by 20% among men and by 28% among women in 2020.

Cocaine-related deaths

- The number of **cocaine**-related deaths increased 6% in 2020 following a 2% decline in 2019.
- **Cocaine**-related deaths increased in 2020 among those 25-34 years (12%), those 45-54 (3%), and those 55 years and older (18%), but decreased among those under 25 (9%) and those 35-44 years (6%).
- **Cocaine**-related deaths increased by 6% among non-Hispanic Whites and 4% among non-Hispanic Blacks in 2020. There was a 59% increase among Hispanics.
- **Cocaine**-related deaths among women increased by 20% in 2020, while the number of deaths among men was similar to 2019.
- The overall increase in **cocaine**-related deaths is largely the result of deaths occurring in combination with opioids. Ninety-one percent of **cocaine**-related deaths in 2020 occurred in combination with **fentanyl**, and 23% in combination with **heroin**.

Benzodiazepine-related deaths

- The number of **benzodiazepine**-related deaths increased by 7% between 2019 and 2020, reversing a two-year decline.
- **Benzodiazepine**-related deaths rose in 2020 among those aged under 25 years (60%), those 35-44 years (27%), and those 45-54 years (47%). Deaths decreased among those aged 25-34 years (23%) and those 55 years and over (26%).
- Deaths increased among non-Hispanic Whites by 11% in 2020, but remained stable among non-Hispanic Blacks and Hispanics.
- A thirteen percent increase was seen among men, while deaths remained stable among women in 2020.
- Ninety-two percent of **benzodiazepine**-related deaths in 2020 were in combination with **opioids**. Seventy-one percent of all **benzodiazepine**-related deaths occurred in combination with **fentanyl**, 38% in combination with **prescription opioids,** and 24% in combination with **heroin**.

Phencyclidine-related deaths (PCP)

- The number of **phencyclidine**-related deaths has been rising since 2018. These deaths increased by 29% between 2019 and 2020.
- **Phencyclidine**-related deaths increased among all age groups between 2019 and 2020 except those under 25 years, who had no deaths in 2020.
- Deaths increased among non-Hispanic Blacks (55%) and decreased 20% in non-Hispanic Whites between 2019 and 2020. After three years of no deaths among Hispanics, there were four deaths in 2020.
- Deaths increased by 30% among men in 2020, and increased by 29% among women.
- Seventy-six percent of **phencyclidine**-related deaths in 2020 were in combination with **opioids**.
- The number of **phencyclidine**-related deaths occurring in Prince George's County was three times higher than the next highest jurisdiction.

Methamphetamine-related deaths

- The number of **methamphetamine**-related deaths has been rising since 2015. These deaths increased by 85% between 2019 and 2020.
- **Methamphetamine**-related deaths increased among all age groups.
- Deaths increased among non-Hispanic Whites (75%) and non-Hispanic Blacks (138%). There was one death among Hispanics, the first death since 2015.
- Deaths increased among both sexes, 90% among males and 70% among females.
- Eighty-three percent of **methamphetamine**-related deaths in 2020 were in combination with **opioids**. Eighty percent of all **methamphetamine**-related deaths occurred in combination with **fentanyl,** 16% in combination with **heroin,** and 11% in combination with **prescription opioids**. Twenty-four percent of **methamphetamine**-related deaths occurred in combination with **cocaine**.
- The number of **methamphetamine**-related deaths occurring in Cecil County was twice as high as the next highest jurisdiction.

Alcohol-related deaths

- The number of **alcohol**-related deaths had declined steadily since 2017. However, deaths sharply increased by 34% between 2019 and 2020.
- **Alcohol**-related deaths in 2020 rose in every age group.
- Deaths increased in 2020 by 41% among non-Hispanic Whites, by 19% among non-Hispanic Blacks, and by 57% among Hispanics.
- Deaths increased in 2020 among both men and women, 34% among males and 33% among females.
- Eighty-one percent of acute **alcohol**-related deaths in 2020 occurred in combination with opioids. Seventy-five percent occurred in combination with **fentanyl**, 27% occurred in combination with **cocaine**, and 11% occurred in combination with **heroin**.

Age-adjusted death rates

- Age-adjusted death rates for the period 2017-2019 ranged from lows of 9.7 and 11.7 per 100,000 population in Montgomery and Prince George's Counties, respectively, to a high of 95.5 per 100,000 population in Baltimore City. The Maryland state age-adjusted mortality rate for deaths related to unintentional drug and alcohol-related intoxication was 33.5 deaths per 100,000 population over the three-year period.

TOTAL INTOXICATION DEATHS

Figure 1. Total Number of Drug- and Alcohol-Related Intoxication Deaths Occurring in Maryland, 2011-2020.




Figure 2. Total Number of Intoxication Deaths Occurring in Maryland by Place of Occurrence, 2020.






Figure 3. Total Number of Drug- and Alcohol-Related Intoxication Deaths Occurring in Maryland by Age Group, Race/Ethnicity and Gender, 2011-2020.

AGE (years)




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<25 years	76	87	79	97	99	161	154	131	118	143
25-34 years	167	169	216	234	297	525	588	561	530	647
35-44 years	128	171	175	217	281	428	461	513	557	589
45-54 years	209	243	246	290	339	550	560	582	529	651
55+ years	91	129	141	203	241	424	517	618	643	769

RACE/ETHNICITY




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
NH White	505	590	611	702	868	1349	1505	1479	1342	1556
NH Black	132	183	214	298	350	654	687	823	928	1076
Hispanic	23	18	21	30	21	53	59	67	75	126

GENDER




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Male	449	548	584	743	893	1513	1626	1766	1723	2019
Female	222	251	274	298	365	574	652	640	654	780

Figure 4. Total Number of Drug- and Alcohol-Related Intoxication Deaths by Place of Occurrence, Maryland, 2011-2020.



DRUG- AND ALCOHOL-RELATED INTOXICATION DEATHS BY SUBSTANCE

Figure 5. Total Number of Drug- and Alcohol-Related Intoxication Deaths by Selected Substances[1], Maryland, 2011-2020.




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Heroin	247	392	464	578	748	1212	1078	830	726	548
Prescription opioids	342	311	316	330	351	418	413	379	369	453
Alcohol	161	195	239	270	309	582	517	472	423	566
Benzodiazepines	68	73	69	103	91	126	146	127	107	114
Cocaine	148	153	154	198	221	464	691	891	869	921
Fentanyl	26	29	58	186	340	1119	1594	1888	1927	2342

[1]Since an intoxication death may involve more than one substance, counts of deaths related to specific substances do not sum to the total number of deaths.

[2]Includes deaths caused by benzodiazepines and related drugs with similar sedative effects.

OPIOID-RELATED DEATHS


Figure 6. Total Number of Opioid* and Non-Opioid- Related Deaths Occurring in Maryland, 2011-2020.



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Opioid-related	529	648	729	888	1089	1856	2009	2143	2106	2518
Non-opioid-related	142	151	129	153	170	233	273	263	273	281

*Total opioids include heroin, prescription opioids, and illicit forms of fentanyl.

Figure 7. Number of Opioid-Related Deaths Occurring in Maryland by Substance, 2011-2020.



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Heroin	247	392	464	578	748	1212	1078	830	726	548
Prescription opioids	342	311	316	330	351	418	413	379	369	453
Fentanyl	26	29	58	186	340	1119	1594	1888	1927	2342

Figure 8. Number of Fentanyl-Related Deaths Occurring in Maryland, 2011-2020.




Figure 9. Number of Fentanyl-Related Deaths Occurring in Maryland by Place of Occurrence, 2020.






Figure 10. Number of Fentanyl-Related Deaths Occurring in Maryland by Age Group, Race/Ethnicity and Gender, 2011-2020.

AGE (years)




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<25 years	1	1	7	28	27	93	118	115	103	120
25-34 years	6	6	18	48	93	313	454	467	452	567
35-44 years	9	10	10	38	72	221	333	414	453	501
45-54 years	9	5	13	49	91	292	380	454	432	534
55+ years	1	7	10	23	57	200	307	437	485	620

RACE/ETHNICITY




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
NH White	25	27	43	103	234	729	1061	1143	1080	1283
NH Black	0	2	14	75	102	350	483	670	776	934
Hispanic	1	0	0	6	2	24	29	43	48	94

GENDER



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Male	15	18	42	148	259	836	1184	1413	1434	1725
Female	11	11	16	38	81	282	410	475	491	617

Figure 11. Number of Fentanyl-Related Deaths by Place of Occurrence, Maryland, 2011-2020.






Figure 12. Number of Heroin-Related Deaths Occurring in Maryland, 2011-2020.




Figure 13. Number of Heroin-Related Deaths Occurring in Maryland by Place of Occurrence, 2020.







Figure 14. Number of Heroin-Related Deaths Occurring in Maryland by Age Group, Race/Ethnicity and Gender, 2011-2020.



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<25 years	28	53	54	61	69	100	86	55	32	19
25-34 years	67	87	129	148	202	312	278	200	186	132
35-44 years	52	81	83	118	165	229	226	196	165	116
45-54 years	67	124	133	157	178	335	277	205	160	138
55+ years	33	47	65	94	133	235	210	174	183	143





	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
NH White	178	285	321	379	520	772	742	536	424	308
NH Black	62	100	131	188	204	397	309	269	273	217
Hispanic	6	2	8	8	8	25	17	18	20	16



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Male	178	294	344	454	588	897	821	615	543	413
Female	69	98	120	124	159	314	257	215	183	135


Figure 15. Number of Heroin-Related Deaths by Place of Occurrence, Maryland, 2011-2020.





PRESCRIPTION OPIOID-RELATED DEATHS

Figure 16. Number of Deaths Occurring in Maryland by Selected Prescription Opioids, 2011-2020.



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Total	342	311	316	330	351	418	413	379	369	453
Methadone	172	170	138	152	183	197	246	196	201	279
Oxycodone	118	99	86	120	104	157	122	103	124	108
Tramadol	24	25	30	33	30	35	42	57	47	76

Figure 17. Number of Prescription Opioid-Related Deaths Occurring in Maryland, 2011-2020.




Figure 18. Number of Prescription Opioid-Related Deaths Occurring in Maryland by Place of Occurrence, 2020.






Figure 19. Number of Prescription Opioid-Related Deaths Occurring in Maryland by Age Group, Race/Ethnicity and Gender, 2011-2020.

AGE (years)




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<25 years	50	29	26	21	16	23	20	14	9	19
25-34 years	88	83	84	69	71	92	90	65	70	78
35-44 years	58	67	68	66	83	98	95	82	90	109
45-54 years	111	80	89	91	107	114	113	101	85	120
55+ years	35	52	49	83	74	91	95	116	115	127

RACE/ETHNICITY




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
NH White	278	258	247	256	265	308	306	263	245	290
NH Black	52	46	63	66	80	105	95	108	118	154
Hispanic	9	4	3	6	3	2	7	5	3	8

GENDER



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Male	218	187	178	199	206	258	238	239	239	286
Female	124	124	138	131	145	160	175	140	130	167

Figure 20. Number of Prescription Opioid-Related Deaths by Place of Occurrence, Maryland, 2011-2020.






Figure 21. Number of Carfentanil-Related Deaths Occurring in Maryland, 2011-2020.





Figure 22. Number of Carfentanil-Related Deaths Occurring in Maryland by Age Group, Race/Ethnicity, and Gender, 2017-2020.



COCAINE-RELATED DEATHS

Figure 23. Number of Cocaine-Related Deaths Occurring in Maryland, 2011-2020.




Figure 24. Number of Cocaine-Related Deaths Occurring in Maryland by Place of Occurrence, 2020.






Figure 25. Number of Cocaine-Related Deaths Occurring in Maryland by Age Group, Race/Ethnicity and Gender, 2011-2020.

AGE (years)




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<25 years	9	8	7	8	9	20	38	36	35	32
25-34 years	34	34	29	33	46	113	175	208	188	211
35-44 years	39	30	35	49	54	97	137	193	219	205
45-54 years	45	57	54	69	76	146	202	239	213	220
55+ years	21	24	29	39	36	88	138	214	214	253

RACE/ETHNICITY




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
NH White	101	92	97	112	134	274	429	532	464	493
NH Black	43	55	51	79	81	172	244	326	368	383
Hispanic	4	3	3	4	3	12	9	21	22	35


GENDER



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Male	106	103	110	149	170	347	485	656	654	663
Female	42	50	44	49	51	117	206	235	215	258

Figure 26. Number of Cocaine-Related Deaths by Place of Occurrence, Maryland, 2011-2020.





BENZODIAZEPINE-RELATED DEATHS

Figure 27. Number of Benzodiazepine-Related Deaths Occurring in Maryland, 2011-2020.




Figure 28. Number of Benzodiazepine-Related Deaths Occurring in Maryland by Place of Occurrence, 2020.







Figure 29. Number of Benzodiazepine-Related Deaths Occurring in Maryland by Age Group, Race/Ethnicity and Gender, 2011-2020.



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<25 years	7	8	3	8	5	12	15	12	5	8
25-34 years	21	18	21	30	23	33	42	33	30	23
35-44 years	13	15	16	18	25	28	31	26	30	38
45-54 years	19	24	18	25	25	28	27	23	19	28
55+ years	8	8	11	22	13	25	31	33	23	17





	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
NH White	60	67	59	82	79	106	130	102	81	90
NH Black	3	6	8	16	9	17	15	20	20	21
Hispanic	2	0	1	2	1	2	0	3	3	3



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Male	37	44	39	67	50	77	82	78	63	71
Female	31	29	30	36	41	49	64	49	44	43

Figure 30. Number of Benzodiazepine-Related Deaths by Place of Occurrence, Maryland, 2011-2020.





PHENCYCLIDINE-RELATED DEATHS

Figure 31. Number of Phencyclidine-Related Deaths Occurring in Maryland, 2011-2020.




Figure 32. Number of Phencyclidine-Related Deaths Occurring in Maryland by Place of Occurrence, 2020.






Figure 33. Number of Phencyclidine-Related Deaths Occurring in Maryland by Age Group, Race/Ethnicity and Gender, 2011-2020.

AGE (years)




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<25 years	0	1	2	1	1	3	0	1	2	0
25-34 years	2	2	0	3	5	6	10	10	11	25
35-44 years	2	5	2	9	4	11	7	6	18	20
45-54 years	6	6	2	1	4	5	7	11	12	14
55+ years	0	2	2	1	0	7	4	9	15	16

RACE/ETHNICITY




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
NH White	6	12	6	7	3	15	8	18	25	20
NH Black	4	4	2	8	11	16	20	18	33	51
Hispanic	0	0	0	0	0	1	0	0	0	4


GENDER



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Male	7	13	5	10	11	27	21	32	44	57
Female	3	3	3	5	3	5	7	5	14	18

Figure 34. Number of Phencyclidine-Related Deaths by Place of Occurrence, Maryland, 2011-2020.





METHAMPHETAMINE-RELATED DEATHS

Figure 35. Number of Methamphetamine-Related Deaths Occurring in Maryland, 2011-2020.




Figure 36. Number of Methamphetamine-Related Deaths Occurring in Maryland by Place of Occurrence, 2020.






Figure 37. Number of Methamphetamine-Related Deaths Occurring in Maryland by Age Group, Race/Ethnicity and Gender, 2011-2020.

AGE (years)




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<25 years	0	1	1	0	1	0	4	4	2	3
25-34 years	0	0	2	1	4	10	9	14	14	31
35-44 years	0	1	0	1	3	4	8	8	13	20
45-54 years	0	0	2	1	2	3	6	4	6	13
55+ years	0	0	1	0	0	1	1	2	6	9

RACE/ETHNICITY




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
NH White	0	2	3	3	6	11	22	28	32	56
NH Black	0	0	3	0	3	6	5	4	8	19
Hispanic	0	0	0	0	1	0	0	0	0	1


GENDER



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Male	0	2	3	3	8	13	24	27	31	59
Female	0	0	3	0	2	5	4	5	10	17

Figure 38. Number of Methamphetamine-Related Deaths by Place of Occurrence, Maryland, 2011-2020.





ALCOHOL-RELATED DEATHS

Figure 39. Number of Alcohol-Related Deaths Occurring in Maryland, 2011-2020.




Figure 40. Number of Alcohol-Related Deaths Occurring in Maryland by Place of Occurrence, 2020.






Figure 41. Number of Alcohol-Related Deaths Occurring in Maryland by Age Group, Race/Ethnicity and Gender, 2011-2020.

AGE (years)




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
<25 years	18	10	9	12	15	25	23	13	13	23
25-34 years	37	26	35	51	52	115	97	85	75	119
35-44 years	31	46	56	55	73	123	102	110	112	132
45-54 years	56	77	88	104	109	195	154	123	97	145
55+ years	19	36	50	48	61	124	141	141	126	147

RACE/ETHNICITY




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
NH White	116	128	146	161	201	348	298	240	215	304
NH Black	32	54	77	87	95	203	174	193	166	198
Hispanic	11	11	13	18	12	26	36	32	35	55


GENDER



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Male	127	150	181	219	256	452	399	372	331	444
Female	34	45	58	51	54	130	118	100	92	122


Figure 42. Number of Alcohol-Related Deaths by Place of Occurrence, Maryland, 2011-2020.






DRUG COMBINATIONS

Figure 43. Number of Drug- and Alcohol-Related Intoxication Deaths Involving Opioids, 2011-2020.




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Total deaths	671	799	858	1041	1259	2089	2282	2406	2379	2799
Opioid-related	529	648	729	888	1089	1856	2009	2143	2106	2518
Not opioid-related	142	151	129	153	170	233	273	263	273	281

Figure 44. Number of Intoxication Deaths by Presence of Heroin and/or Fentanyl, 2011-2020.




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Total deaths	671	799	858	1041	1259	2089	2282	2406	2379	2799
Both Heroin and fentanyl	5	2	24	101	220	705	844	718	662	517
Either Heroin or Fentanyl	268	419	498	663	868	1626	1828	2000	1991	2373
Neither Heroin or Fentanyl	403	380	360	378	391	463	454	406	388	426

Figure 45. Number of Prescription Opioid-Related Intoxication Deaths Involving Heroin or Fentanyl, 2011-2020.




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Total deaths	342	311	316	330	351	418	413	379	369	453
In combination with heroin or fentanyl	81	82	85	105	130	213	244	246	261	317
Not in combination with heroin or fentanyl	261	229	231	225	221	205	169	133	108	136

Figure 46. Number of Cocaine-Related Intoxication Deaths Involving Heroin or Fentanyl, 2011-2020.




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Total deaths	148	153	154	198	221	464	691	891	869	921
In combination w/heroin or fentanyl	64	82	95	143	170	365	554	770	759	843
Not in combination w/heroin or fentanyl	84	71	59	55	51	99	137	121	110	78

Figure 47. Number of Benzodiazepine-Related Intoxication Deaths Involving Heroin or Fentanyl, 2011-2020.



	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Total deaths	68	73	69	103	91	126	146	127	107	114
In combination with heroin or fentanyl	13	20	16	36	45	77	91	83	83	82
Not in combination with heroin or fentanyl	55	53	53	67	46	49	55	44	24	32

Figure 48. Number of Alcohol-Related Intoxication Deaths Involving Heroin or Fentanyl, 2011-2020.




	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020
Total deaths	161	195	239	270	310	582	517	472	423	566
In combination with heroin or fentanyl	46	95	137	162	207	438	364	356	316	430
Not in combination with heroin or fentanyl	115	100	102	108	103	144	153	116	107	136

Figure 49. Combinations of Substances Related to Unintentional Drug-and Alcohol-Related Intoxication Deaths, Maryland, 2020		
	Number	Percent
Fentanyl		
Total	2342	
In combination		
With cocaine	835	35.7
With heroin	517	22.1
With alcohol	425	18.1
With prescription opioids	309	13.2
With benzodiazepines	81	3.5
With methamphetamine	61	2.6
With phencyclidine	54	2.3
Cocaine		
Total	921	
In combination		
With fentanyl	835	90.7
With heroin	208	22.6
With alcohol	152	16.5
With prescription opioids	110	11.9
With benzodiazepines	25	2.7
With phencyclidine	21	2.3
With methamphetamine	18	2.0
Heroin		
Total	548	
In combination		
With fentanyl	517	94.3
With cocaine	208	38.0
With prescription opioids	106	19.3
With alcohol	64	11.7
With benzodiazepines	27	4.9
With phencyclidine	15	2.7
With methamphetamine	12	2.2
Prescription opioids		
Total	453	
In combination		
With fentanyl	309	68.2
With cocaine	110	24.3
With heroin	106	23.4
With alcohol	55	12.1
With benzodiazepines	43	9.5
With methamphetamine	8	1.8
With phencyclidine	4	0.9

Figure 49. Combinations of Substances Related to Unintentional Drug-and Alcohol-Related Intoxication Deaths, Maryland, 2020 - Continued		
	Number	Percent
Alcohol		
Total	566	
In combination		
With fentanyl	425	75.1
With cocaine	152	26.9
With heroin	64	11.3
With prescription opioids	55	9.7
With phencyclidine	20	3.5
With benzodiazepines	18	3.2
With methamphetamine	8	1.4
Benzodiazepines		
Total	114	
In combination		
With fentanyl	81	71.1
With prescription opioids	43	37.7
With heroin	27	23.7
With cocaine	25	21.9
With alcohol	18	15.8
With methamphetamine	3	2.6
With phencyclidine	2	1.8
Phencyclidine		
Total	75	
In combination		
With fentanyl	54	72.0
With cocaine	21	28.0
With alcohol	20	26.7
With heroin	15	20.0
With prescription opioids	4	5.3
With benzodiazepines	2	2.7
With methamphetamine	1	1.3
Methamphetamine		
Total	76	
In combination		
With fentanyl	61	80.3
With cocaine	18	23.7
With heroin	12	15.8
With alcohol	8	10.5
With prescription opioids	8	10.5
With benzodiazepines	3	3.9
With phencyclidine	1	1.3

Figure 50. Age-Adjusted Mortality Rates[1,2] for Total Unintentional Intoxication Deaths by Place of Residence,[3] Maryland, 2017-2019.



[1]Age-adjusted to the 2000 U.S. standard population by the direct method.
[2]Since age-adjusted rates based on fewer than 20 deaths are considered unreliable, rates are only show for jurisdictions with 20 or more intoxication deaths over the five-year period.
[3]Rates are based on place of residence, not place of occurrence.

TABLES

TABLE 1. TOTAL NUMBER OF DRUG AND ALCOHOL-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	TOTAL INTOXICATION DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	671	799	858	1,041	1,259	2,089	2,282	2,406	2,379	2,799	16,583
NORTHWEST AREA	65	67	86	96	131	214	183	211	189	234	1,476
GARRETT	2	0	6	2	5	1	8	3	9	8	44
ALLEGANY	12	14	15	12	22	59	38	39	28	52	291
WASHINGTON	21	27	28	40	64	66	59	91	88	110	594
FREDERICK	30	26	37	42	40	88	78	78	64	64	547
BALTIMORE METRO AREA	420	519	557	678	841	1,402	1,549	1,731	1,652	1,860	11,209
BALTIMORE CITY	167	225	246	305	393	694	761	888	914	1,028	5,621
BALTIMORE COUNTY	107	119	144	170	220	336	367	388	350	394	2,595
ANNE ARUNDEL	79	83	78	101	112	195	214	241	208	251	1,562
CARROLL	8	29	24	38	40	47	55	72	56	46	415
HOWARD	21	24	29	21	26	46	51	41	37	57	353
HARFORD	38	39	36	43	50	84	101	101	87	84	663
NATIONAL CAPITAL AREA	86	104	111	128	140	231	283	216	251	342	1,892
MONTGOMERY	44	48	52	65	70	102	116	89	105	139	830
PRINCE GEORGE'S	42	56	59	63	70	129	167	127	146	203	1,062
SOUTHERN AREA	31	37	25	47	59	88	103	86	95	111	682
CALVERT	12	12	6	17	20	28	32	28	31	25	211
CHARLES	11	13	9	21	22	45	37	27	31	53	269
ST MARY'S	8	12	10	9	17	15	34	31	33	33	202
EASTERN SHORE AREA	69	72	79	92	88	154	164	162	192	252	1,324
CECIL	28	25	26	29	32	30	59	59	62	92	442
KENT	2	0	4	6	3	6	5	2	10	6	44
QUEEN ANNE'S	5	2	8	10	4	8	8	17	13	14	89
CAROLINE	11	4	2	7	3	10	11	7	12	17	84
TALBOT	1	5	7	4	5	10	11	10	14	17	84
DORCHESTER	2	5	5	0	1	6	12	7	11	17	66
WICOMICO	11	21	17	20	18	48	35	36	41	47	294
SOMERSET	3	3	4	3	6	8	4	8	10	16	65
WORCESTER	6	7	6	13	16	28	19	16	19	26	156

[1] Includes deaths that were the result of recent ingestion or exposure to alcohol or another type of drug, including heroin, cocaine, prescription opioids, benzodiazepines, and other prescribed and unprescribed drugs.
[2] Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 2. TOTAL NUMBER OF OPIOID-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	OPIOID-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	529	648	729	888	1,089	1,856	2,009	2,143	2,106	2,518	14,515
NORTHWEST AREA	53	53	74	81	118	198	157	189	168	219	1,310
GARRETT	1	0	4	2	4	0	4	3	6	5	29
ALLEGANY	8	10	11	11	20	55	36	33	23	48	255
WASHINGTON	16	20	26	34	57	63	51	83	80	105	535
FREDERICK	28	23	33	34	37	80	66	70	59	61	491
BALTIMORE METRO AREA	341	437	485	591	742	1,262	1,404	1,578	1,508	1,715	10,063
BALTIMORE CITY	142	189	212	275	354	628	692	814	851	964	5,121
BALTIMORE COUNTY	93	104	125	146	195	305	323	352	316	356	2,315
ANNE ARUNDEL	53	68	67	85	89	169	198	218	183	226	1,356
CARROLL	7	27	21	29	34	44	51	68	51	43	375
HOWARD	18	17	26	18	25	40	47	36	34	52	313
HARFORD	28	32	34	38	45	76	93	90	73	74	583
NATIONAL CAPITAL AREA	52	66	78	101	104	190	215	158	188	268	1,420
MONTGOMERY	28	36	40	53	59	84	91	64	86	109	650
PRINCE GEORGE'S	24	30	38	48	45	106	124	94	102	159	770
SOUTHERN AREA	26	32	24	40	48	74	94	71	82	93	584
CALVERT	10	11	5	16	19	25	27	25	25	19	182
CHARLES	10	12	9	16	17	36	34	19	26	42	221
ST MARY'S	6	9	10	8	12	13	33	27	31	32	181
EASTERN SHORE AREA	57	60	68	75	77	132	139	147	160	223	1,138
CECIL	24	22	22	25	26	28	57	58	53	85	400
KENT	1	0	4	3	3	4	4	2	10	6	37
QUEEN ANNE'S	4	2	7	9	4	6	6	16	11	13	78
CAROLINE	8	4	2	7	3	9	8	7	11	15	74
TALBOT	1	3	6	4	5	10	8	10	13	13	73
DORCHESTER	2	5	5	0	1	5	10	6	10	15	59
WICOMICO	10	17	14	15	17	44	28	30	29	39	243
SOMERSET	3	2	4	2	4	6	3	8	9	13	54
WORCESTER	4	5	4	10	14	20	15	10	14	24	120

[1] Includes deaths confirmed or suspected to be related to recent ingestion of opioids.
[2] Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 3. TOTAL NUMBER OF HEROIN-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	HEROIN-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	247	392	464	578	748	1,212	1,078	830	726	548	6,823
NORTHWEST AREA	23	27	40	53	80	119	72	68	58	44	584
GARRETT	1	0	2	1	3	0	1	1	1	1	11
ALLEGANY	3	6	3	5	13	34	14	15	9	14	116
WASHINGTON	8	11	14	21	38	39	22	29	25	20	227
FREDERICK	11	10	21	26	26	46	35	23	23	9	230
BALTIMORE METRO AREA	165	272	319	379	519	858	772	572	505	364	4,725
BALTIMORE CITY	76	131	150	192	260	454	380	286	279	205	2,413
BALTIMORE COUNTY	38	64	76	86	134	208	170	119	111	74	1,080
ANNE ARUNDEL	24	38	41	53	60	105	118	75	63	45	622
CARROLL	2	13	14	16	22	25	28	34	18	13	185
HOWARD	10	12	16	9	16	24	23	15	10	15	150
HARFORD	15	14	22	23	27	42	53	43	24	12	275
NATIONAL CAPITAL AREA	23	42	53	65	69	115	104	78	81	76	706
MONTGOMERY	11	22	28	33	37	48	52	34	39	32	336
PRINCE GEORGE'S	12	20	25	32	32	67	52	44	42	44	370
SOUTHERN AREA	15	18	13	28	29	48	45	31	30	24	281
CALVERT	5	6	2	13	15	17	17	8	10	3	96
CHARLES	6	5	5	10	8	22	16	11	12	15	110
ST MARY'S	4	7	6	5	6	9	12	12	8	6	75
EASTERN SHORE AREA	21	33	39	53	51	72	85	81	52	40	527
CECIL	8	11	11	15	16	19	37	40	16	12	185
KENT	1	0	0	2	1	1	1	0	3	2	11
QUEEN ANNE'S	2	2	5	7	1	4	5	8	3	4	41
CAROLINE	3	3	2	6	2	6	4	3	5	4	38
TALBOT	1	2	2	4	3	4	3	4	4	1	28
DORCHESTER	1	3	3	0	1	3	4	3	5	2	25
WICOMICO	3	9	11	12	13	21	20	12	9	8	118
SOMERSET	1	2	1	1	3	3	2	5	1	1	20
WORCESTER	1	1	4	6	11	11	9	6	6	6	61

[1] Includes deaths confirmed or suspected to be related to recent heroin use.
[2] Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 4. TOTAL NUMBER OF PRESCRIPTION OPIOID-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	PRESCRIPTION OPIOID-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	342	311	316	330	351	418	413	379	369	453	3,682
NORTHWEST AREA	38	30	35	33	39	56	35	34	33	38	371
GARRETT	1	0	2	2	1	0	1	1	1	1	10
ALLEGANY	5	5	8	6	6	15	9	5	5	8	72
WASHINGTON	11	9	11	16	20	23	8	19	17	18	152
FREDERICK	21	16	14	9	12	18	17	9	10	11	137
BALTIMORE METRO AREA	212	196	207	217	233	265	298	272	258	325	2,483
BALTIMORE CITY	82	74	86	84	105	113	123	128	134	168	1,097
BALTIMORE COUNTY	68	47	54	59	62	67	87	71	60	71	646
ANNE ARUNDEL	33	33	28	32	27	48	43	36	27	40	347
CARROLL	5	17	12	15	14	15	13	16	13	16	136
HOWARD	9	5	13	7	9	6	13	2	9	11	84
HARFORD	15	20	14	20	16	16	19	19	15	19	173
NATIONAL CAPITAL AREA	35	29	30	35	36	42	33	27	28	37	332
MONTGOMERY	20	18	16	19	23	26	19	16	15	16	188
PRINCE GEORGE'S	15	11	14	16	13	16	14	11	13	21	144
SOUTHERN AREA	15	18	12	19	19	25	26	22	23	24	203
CALVERT	7	6	3	7	6	11	5	6	5	5	61
CHARLES	5	7	5	9	8	10	11	8	7	8	78
ST MARY'S	3	5	4	3	5	4	10	8	11	11	64
EASTERN SHORE AREA	42	38	32	26	24	30	21	24	27	29	293
CECIL	20	18	12	12	10	8	8	5	6	10	109
KENT	1	0	4	2	2	0	2	0	0	0	11
QUEEN ANNE'S	2	0	3	3	3	2	2	4	0	3	22
CAROLINE	5	1	0	1	0	4	1	1	3	3	19
TALBOT	0	1	4	0	2	3	4	2	5	2	23
DORCHESTER	1	3	3	0	0	2	2	2	3	0	16
WICOMICO	7	9	4	3	5	7	0	5	5	7	52
SOMERSET	3	2	2	1	1	0	1	2	2	1	15
WORCESTER	3	4	0	4	1	4	1	3	3	3	26

[1] Includes deaths confirmed or suspected to be related to recent ingestion of one or more prescription opioids.
[2] Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 5. TOTAL NUMBER OF OXYCODONE-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	OXYCODONE-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	118	99	86	120	104	157	122	103	124	108	1,141
NORTHWEST AREA	11	13	12	10	11	25	16	13	18	15	144
GARRETT	0	0	1	0	0	0	0	0	1	1	3
ALLEGANY	0	2	3	3	2	7	3	2	2	5	29
WASHINGTON	5	2	5	5	6	11	2	7	9	2	54
FREDERICK	6	9	3	2	3	7	11	4	6	7	58
BALTIMORE METRO AREA	63	51	44	69	56	77	73	67	64	59	623
BALTIMORE CITY	15	15	11	20	18	22	23	21	22	21	188
BALTIMORE COUNTY	22	12	14	22	16	22	21	20	18	14	181
ANNE ARUNDEL	14	11	9	10	12	23	15	15	11	14	134
CARROLL	3	6	3	4	3	3	4	7	4	5	42
HOWARD	2	2	4	4	4	2	5	0	3	3	29
HARFORD	7	5	3	9	3	5	5	4	6	2	49
NATIONAL CAPITAL AREA	14	11	13	17	16	25	13	7	15	14	145
MONTGOMERY	9	8	7	11	8	16	8	4	6	7	84
PRINCE GEORGE'S	5	3	6	6	8	9	5	3	9	7	61
SOUTHERN AREA	10	10	6	11	13	13	14	10	16	13	116
CALVERT	4	5	3	3	3	7	3	1	4	4	37
CHARLES	4	3	1	5	8	4	7	5	4	3	44
ST MARY'S	2	2	2	3	2	2	4	4	8	6	35
EASTERN SHORE AREA	20	14	11	13	8	17	6	6	11	7	113
CECIL	9	4	6	6	3	2	2	0	2	2	36
KENT	0	0	1	0	1	0	0	0	0	0	2
QUEEN ANNE'S	1	0	1	1	2	1	0	1	0	1	8
CAROLINE	0	0	0	0	0	3	0	1	2	2	8
TALBOT	0	1	1	0	0	2	2	0	2	0	8
DORCHESTER	1	1	0	0	0	2	1	1	1	0	7
WICOMICO	5	5	1	2	1	5	0	2	3	0	24
SOMERSET	2	1	1	1	0	0	0	1	1	1	8
WORCESTER	2	2	0	3	1	2	1	0	0	1	12

[1] Includes deaths confirmed or suspected to be related to recent ingestion of oxycodone.
[2] Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 6: TOTAL NUMBER OF METHADONE-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	METHADONE-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	172	170	138	152	183	197	246	196	201	279	1,934
NORTHWEST AREA	14	14	8	20	14	12	11	14	10	17	134
GARRETT	0	0	1	1	0	0	0	0	0	0	2
ALLEGANY	4	1	1	3	2	4	3	2	2	1	23
WASHINGTON	5	4	3	10	6	5	4	10	6	14	67
FREDERICK	5	9	3	6	6	3	4	2	2	2	42
BALTIMORE METRO AREA	128	122	110	112	145	158	198	155	166	226	1,520
BALTIMORE CITY	65	54	57	54	78	82	87	85	98	131	791
BALTIMORE COUNTY	32	28	29	31	34	36	63	37	36	46	372
ANNE ARUNDEL	17	15	6	14	9	21	23	12	12	21	150
CARROLL	2	12	7	5	9	9	6	6	8	9	73
HOWARD	5	1	5	2	5	2	8	1	6	3	38
HARFORD	7	12	6	6	10	8	11	14	6	16	96
NATIONAL CAPITAL AREA	13	13	7	6	9	13	14	7	6	13	101
MONTGOMERY	6	7	3	5	6	7	6	4	4	5	53
PRINCE GEORGE'S	7	6	4	1	3	6	8	3	2	8	48
SOUTHERN AREA	3	5	2	7	6	6	9	7	6	11	62
CALVERT	2	2	0	2	3	2	3	4	0	1	19
CHARLES	0	1	1	4	2	2	3	2	2	4	21
ST MARY'S	1	2	1	1	1	2	3	1	4	6	22
EASTERN SHORE AREA	14	16	11	7	9	8	14	13	13	12	117
CECIL	9	10	4	4	3	3	4	5	4	6	52
KENT	1	0	2	1	1	0	2	0	0	0	7
QUEEN ANNE'S	1	0	1	0	1	1	2	3	0	0	9
CAROLINE	1	1	0	1	0	2	1	0	1	0	7
TALBOT	0	1	2	0	1	1	2	1	2	1	11
DORCHESTER	0	1	0	0	0	0	2	1	2	0	6
WICOMICO	1	1	2	0	2	0	0	1	2	3	12
SOMERSET	1	0	0	0	1	0	1	0	1	0	4
WORCESTER	0	2	0	1	0	1	0	2	1	2	9

1 Includes deaths confirmed or suspected to be related to recent ingestion of methadone.
2 Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 7: TOTAL NUMBER OF FENTANYL-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	FENTANYL-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	26	29	58	186	340	1,119	1,594	1,888	1,927	2,342	9,509
NORTHWEST AREA	6	3	7	8	32	109	119	166	146	200	796
GARRETT	1	0	0	0	2	0	2	2	5	5	17
ALLEGANY	1	1	1	1	5	29	29	29	19	44	159
WASHINGTON	1	1	4	1	14	31	39	70	70	95	326
FREDERICK	3	1	2	6	11	49	49	65	52	56	294
BALTIMORE METRO AREA	10	16	35	142	248	792	1,118	1,415	1,395	1,605	6,776
BALTIMORE CITY	2	4	12	72	120	419	573	758	810	920	3,690
BALTIMORE COUNTY	4	5	11	36	65	182	244	308	285	328	1,468
ANNE ARUNDEL	2	3	6	23	29	98	152	184	164	209	870
CARROLL	0	1	2	4	11	20	40	55	47	37	217
HOWARD	0	2	3	5	7	27	36	34	28	44	186
HARFORD	2	1	1	2	16	46	73	76	61	67	345
NATIONAL CAPITAL AREA	0	3	6	15	32	101	175	115	167	251	865
MONTGOMERY	0	2	0	8	17	43	72	40	76	102	360
PRINCE GEORGE'S	0	1	6	7	15	58	103	75	91	149	505
SOUTHERN AREA	3	1	4	9	9	32	74	60	74	79	345
CALVERT	1	0	0	5	2	11	22	23	23	16	103
CHARLES	1	1	3	1	4	17	26	14	24	37	128
ST MARY'S	1	0	1	3	3	4	26	23	27	26	114
EASTERN SHORE AREA	7	6	6	12	19	85	108	132	145	207	727
CECIL	2	0	0	1	7	9	44	52	49	81	245
KENT	0	0	0	1	0	3	3	2	10	6	25
QUEEN ANNE'S	0	0	1	1	0	4	5	16	10	12	49
CAROLINE	4	0	0	0	1	3	7	6	9	14	44
TALBOT	0	1	0	2	2	7	3	10	11	11	47
DORCHESTER	0	0	2	0	1	3	7	4	9	15	41
WICOMICO	1	4	1	7	1	34	24	24	26	34	156
SOMERSET	0	0	2	0	1	6	3	8	9	12	41
WORCESTER	0	1	0	0	6	16	12	10	12	22	79

[1] Includes deaths confirmed or suspected to be related to recent ingestion or exposure to pharmaceutical or nonpharmaceutical fentanyl.
[2] Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 8: TOTAL NUMBER OF COCAINE-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	COCAINE-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	148	153	154	198	221	464	691	891	869	921	4,710
NORTHWEST AREA	10	9	13	16	20	27	43	67	51	65	321
GARRETT	0	0	0	0	1	0	1	0	3	1	6
ALLEGANY	0	2	2	2	5	9	13	12	6	10	61
WASHINGTON	3	5	6	6	10	9	10	31	24	31	135
FREDERICK	7	2	5	8	4	9	19	24	18	23	119
BALTIMORE METRO AREA	97	108	102	138	167	348	522	693	647	666	3,488
BALTIMORE CITY	48	59	47	82	93	202	285	388	380	393	1,977
BALTIMORE COUNTY	19	17	27	28	38	80	123	132	138	135	737
ANNE ARUNDEL	18	13	12	19	19	31	66	91	72	89	430
CARROLL	3	7	7	2	6	8	14	23	24	8	102
HOWARD	5	7	5	3	6	7	16	19	9	16	93
HARFORD	4	5	4	4	5	20	18	40	24	25	149
NATIONAL CAPITAL AREA	24	22	25	29	16	44	62	49	74	86	431
MONTGOMERY	12	12	13	10	5	11	17	18	29	26	153
PRINCE GEORGE'S	12	10	12	19	11	33	45	31	45	60	278
SOUTHERN AREA	3	6	1	3	6	8	19	33	39	33	151
CALVERT	2	3	0	2	0	2	3	3	9	8	32
CHARLES	1	1	0	0	2	4	10	13	12	16	59
ST MARY'S	0	2	1	1	4	2	6	17	18	9	60
EASTERN SHORE AREA	14	8	13	12	12	37	45	49	58	71	319
CECIL	7	2	5	4	3	3	15	14	12	13	78
KENT	0	0	0	1	1	0	1	1	4	2	10
QUEEN ANNE'S	1	0	0	0	0	1	2	5	6	4	19
CAROLINE	1	1	0	1	0	5	2	1	2	2	15
TALBOT	0	0	3	0	1	2	2	3	6	5	22
DORCHESTER	1	1	1	0	0	1	7	2	5	6	24
WICOMICO	3	4	3	4	7	13	7	13	21	21	96
SOMERSET	0	0	0	0	0	4	2	6	2	7	21
WORCESTER	1	0	1	2	0	8	7	4	0	11	34

[1] Includes deaths confirmed or suspected to be related to recent use of cocaine.
[2] Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 9: TOTAL NUMBER OF BENZODIAZEPINE-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	BENZODIAZEPINE-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	68	73	69	103	91	126	146	127	107	114	1,024
NORTHWEST AREA	9	5	6	13	8	21	19	10	9	15	115
GARRETT	0	0	1	0	1	0	2	0	1	2	7
ALLEGANY	1	0	1	3	1	6	5	1	1	3	22
WASHINGTON	4	3	2	5	3	6	2	4	2	4	35
FREDERICK	4	2	2	5	3	9	10	5	5	6	51
BALTIMORE METRO AREA	39	49	44	66	56	78	98	90	64	69	653
BALTIMORE CITY	9	15	14	22	15	24	28	28	27	26	208
BALTIMORE COUNTY	9	12	16	24	18	29	25	32	17	21	203
ANNE ARUNDEL	14	11	3	9	11	9	27	16	11	10	121
CARROLL	0	1	3	3	4	1	4	4	3	2	25
HOWARD	4	2	5	0	6	8	5	1	1	4	36
HARFORD	3	8	3	8	2	7	9	9	5	6	60
NATIONAL CAPITAL AREA	9	6	7	12	8	12	15	15	17	17	118
MONTGOMERY	6	4	4	10	7	7	8	9	10	8	73
PRINCE GEORGE'S	3	2	3	2	1	5	7	6	7	9	45
SOUTHERN AREA	2	4	4	6	7	7	8	4	7	6	55
CALVERT	1	1	1	3	1	1	2	2	0	0	12
CHARLES	0	2	1	2	4	4	4	1	3	3	24
ST MARY'S	1	1	2	1	2	2	2	1	4	3	19
EASTERN SHORE AREA	9	9	8	6	12	8	6	8	10	7	83
CECIL	6	7	3	3	5	2	1	2	4	2	35
KENT	0	0	0	0	0	1	2	0	0	0	3
QUEEN ANNE'S	1	0	0	0	1	1	0	3	0	1	7
CAROLINE	0	0	0	0	0	0	1	0	1	2	4
TALBOT	0	0	3	0	1	1	1	0	1	0	7
DORCHESTER	0	1	1	0	0	1	0	0	1	0	4
WICOMICO	1	0	0	1	2	1	0	1	1	2	9
SOMERSET	0	1	1	0	0	0	0	0	0	0	2
WORCESTER	1	0	0	2	3	1	1	2	2	0	12

1 Includes deaths confirmed or suspected to be related to recent ingestion of a benzodiazepine or related drug with sedative effects.
2 Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 10: TOTAL NUMBER OF PHENCYCLIDINE-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	PHENCYCLIDINE-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	10	16	8	15	14	32	28	37	58	75	293
NORTHWEST AREA	0	1	0	1	2	4	1	4	3	5	21
GARRETT	0	0	0	0	0	0	0	0	0	0	0
ALLEGANY	0	0	0	0	0	0	0	0	0	0	0
WASHINGTON	0	0	0	0	1	0	1	0	0	2	4
FREDERICK	0	1	0	1	1	4	0	4	3	3	17
BALTIMORE METRO AREA	4	7	4	3	2	11	8	9	23	19	90
BALTIMORE CITY	1	2	1	1	1	2	2	3	8	5	26
BALTIMORE COUNTY	0	1	1	0	0	2	1	0	2	2	9
ANNE ARUNDEL	3	2	1	1	1	6	5	5	11	10	45
CARROLL	0	1	0	0	0	0	0	0	0	0	1
HOWARD	0	1	1	1	0	1	0	1	2	1	8
HARFORD	0	0	0	0	0	0	0	0	0	1	1
NATIONAL CAPITAL AREA	4	6	3	10	8	13	16	20	25	41	146
MONTGOMERY	2	1	0	1	1	2	2	4	2	10	25
PRINCE GEORGE'S	2	5	3	9	7	11	14	16	23	31	121
SOUTHERN AREA	1	2	1	1	2	3	2	4	6	8	30
CALVERT	1	0	0	0	1	0	2	1	2	3	10
CHARLES	0	2	1	1	0	3	0	3	4	4	18
ST MARY'S	0	0	0	0	1	0	0	0	0	1	2
EASTERN SHORE AREA	1	0	0	0	0	1	1	0	1	2	6
CECIL	0	0	0	0	0	0	0	0	1	0	1
KENT	0	0	0	0	0	0	0	0	0	0	0
QUEEN ANNE'S	0	0	0	0	0	0	0	0	0	0	0
CAROLINE	0	0	0	0	0	0	0	0	0	0	0
TALBOT	0	0	0	0	0	0	0	0	0	1	1
DORCHESTER	1	0	0	0	0	0	0	0	0	0	1
WICOMICO	0	0	0	0	0	1	1	0	0	1	3
SOMERSET	0	0	0	0	0	0	0	0	0	0	0
WORCESTER	0	0	0	0	0	0	0	0	0	0	0

1 Includes deaths confirmed or suspected to be related to recent ingestion of phencyclidine.
2 Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 11: TOTAL NUMBER OF METHAMPHETAMINE-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	METHAMPHETAMINE-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	0	2	6	3	10	18	28	32	41	76	216
NORTHWEST AREA	0	1	1	0	1	2	5	6	11	7	34
GARRETT	0	0	0	0	0	0	2	1	1	3	7
ALLEGANY	0	0	1	0	0	1	0	2	2	1	7
WASHINGTON	0	0	0	0	1	1	1	3	7	2	15
FREDERICK	0	1	0	0	0	0	2	0	1	1	5
BALTIMORE METRO AREA	0	1	3	1	4	12	12	13	13	28	87
BALTIMORE CITY	0	1	2	1	1	8	5	5	7	13	43
BALTIMORE COUNTY	0	0	1	0	0	1	1	4	4	7	18
ANNE ARUNDEL	0	0	0	0	0	0	2	2	2	1	7
CARROLL	0	0	0	0	1	0	1	1	0	2	5
HOWARD	0	0	0	0	2	2	1	1	0	1	7
HARFORD	0	0	0	0	0	1	2	0	0	4	7
NATIONAL CAPITAL AREA	0	0	2	0	4	3	4	4	6	9	32
MONTGOMERY	0	0	0	0	0	1	2	1	3	2	9
PRINCE GEORGE'S	0	0	2	0	4	2	2	3	3	7	23
SOUTHERN AREA	0	0	0	0	1	1	3	1	1	2	9
CALVERT	0	0	0	0	0	0	1	1	1	0	3
CHARLES	0	0	0	0	1	1	2	0	0	2	6
ST MARY'S	0	0	0	0	0	0	0	0	0	0	0
EASTERN SHORE AREA	0	0	0	2	0	0	4	8	10	30	54
CECIL	0	0	0	0	0	0	4	6	8	26	44
KENT	0	0	0	0	0	0	0	0	1	0	1
QUEEN ANNE'S	0	0	0	0	0	0	0	0	0	1	1
CAROLINE	0	0	0	1	0	0	0	0	1	1	3
TALBOT	0	0	0	0	0	0	0	0	0	0	0
DORCHESTER	0	0	0	0	0	0	0	0	0	0	0
WICOMICO	0	0	0	1	0	0	0	1	0	0	2
SOMERSET	0	0	0	0	0	0	0	0	0	0	0
WORCESTER	0	0	0	0	0	0	0	1	0	2	3

[1] Includes deaths confirmed or suspected to be related to recent ingestion of methamphetamine.
[2] Includes only deaths for which the manner of death was classified as accidental or undetermined.

TABLE 12: TOTAL NUMBER OF ALCOHOL-RELATED INTOXICATION DEATHS BY PLACE OF OCCURRENCE, 2011-2020.[1,2]

REGION AND POLITICAL SUBDIVISION	ALCOHOL-RELATED DEATHS										
	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	TOTAL
MARYLAND	161	195	239	270	310	582	517	472	423	566	3,735
NORTHWEST AREA	16	12	21	27	30	47	31	34	37	37	292
GARRETT	1	0	2	1	1	1	2	1	2	1	12
ALLEGANY	2	4	2	3	6	14	4	7	3	7	52
WASHINGTON	4	3	6	11	10	17	14	15	20	17	117
FREDERICK	9	5	11	12	13	15	11	11	12	12	111
BALTIMORE METRO AREA	99	126	154	166	215	403	334	339	273	343	2,452
BALTIMORE CITY	44	71	86	86	114	222	198	187	165	169	1,342
BALTIMORE COUNTY	22	24	32	39	52	81	71	80	53	91	545
ANNE ARUNDEL	21	15	22	18	27	56	37	44	34	49	323
CARROLL	4	4	4	9	6	12	9	10	6	7	71
HOWARD	4	6	6	6	5	14	7	5	4	11	68
HARFORD	4	6	4	8	11	18	12	13	11	16	103
NATIONAL CAPITAL AREA	28	38	35	36	32	67	86	51	58	102	533
MONTGOMERY	16	15	13	18	15	22	35	19	19	43	215
PRINCE GEORGE'S	12	23	22	18	17	45	51	32	39	59	318
SOUTHERN AREA	7	7	7	12	11	22	24	17	18	28	153
CALVERT	2	2	1	4	3	7	4	9	5	10	47
CHARLES	3	2	4	5	4	12	9	3	10	14	66
ST MARY'S	2	3	2	3	4	3	11	5	3	4	40
EASTERN SHORE AREA	11	12	22	29	22	43	42	31	37	56	305
CECIL	3	6	9	5	8	8	12	10	5	16	82
KENT	0	0	1	1	0	1	1	0	1	2	7
QUEEN ANNE'S	3	0	1	7	0	2	4	3	1	1	22
CAROLINE	1	0	1	2	0	5	4	1	2	3	19
TALBOT	0	2	2	0	0	0	5	4	5	6	24
DORCHESTER	0	1	0	0	1	1	2	1	5	5	16
WICOMICO	2	2	6	7	3	12	9	8	12	8	69
SOMERSET	1	1	1	2	2	3	1	0	1	4	16
WORCESTER	1	0	1	5	8	11	4	4	5	11	50

[1] Includes deaths confirmed or suspected to be related to recent ingestion of alcohol.
[2] Includes only deaths for which the manner of death was classified as accidental or undetermined.