

**(0.0.1)** Show that the localization of a ring  $R$  at a prime ideal  $\mathfrak{p}$  is a local ring. What is the maximal ideal?

*Solution.* I claim that the unique maximal ideal is the ideal of nonunits of  $R_{\mathfrak{p}}$ , more usually defined as

$$\mathfrak{m} = \left\{ \frac{a}{b} : \text{some element of } \mathfrak{p} \text{ divides } a \right\}$$

For any other ideal  $\mathfrak{i}$ , there are two options:

- (a) It contains only nonunits. Then obviously  $\mathfrak{i} \subset \mathfrak{m}$ .
- (b) It contains some element that is a unit. Since ideals are closed under multiplication, multiply by the inverse of the unit to get a 1 in the ideal, implying that  $\mathfrak{i} = (1)$ .

Hence all proper ideals are contained in  $\mathfrak{m}$ . □



# Chapter 1

## Rings and ideals

### 1.1 Nilradical and Jacobson radical

Define the *nilradical* of a ring  $R$  as

$$N_R := \sqrt{(0)_R}$$

equivalently,  $N(R)$  is the “ideal of nilpotents”. The main use of the nilradical of a ring (as far as I know!) is to eliminate the problems associated with nilpotents by looking at  $R/N_R$  instead of  $R$ : it seems plausible (at least intuitively) that this ring has none.

**(1.1.1)**  $N_R$  is an ideal.

*Proof.* Closure under  $R$ -multiplication is clear. As for additive closure, consider  $x, y \in N_R$ , such that  $x^m = y^n = 0$ . We must show that  $x + y$  is also a nilpotent.

Consider  $(x + y)^{m+n-1}$ . Every term of the binomial expansion contains either a power of  $x^{r \geq m}$  or  $y^{s \geq n}$  (else  $r + s < m + n - 1$ ), so the entire thing is zero.  $\square$

**(1.1.2)**  $R/N_R$  has no nonzero nilpotents.

*Proof.* Suppose  $\bar{x}$  is a nilpotent in  $R/N_R$ , with  $\bar{x}^n = 0$ . Then  $x^n$  is a nilpotent in  $R$  and is mapped to  $(0) \in R/N_R$ , implying that  $\bar{x} = 0$ .  $\square$

**(1.1.3)**