

# Practical Exercises for Exercise Collection

Sonja Hartnack, Terence Odoch & Muriel Buri

July 2019

## Exercise 1: Terminology

Group the following terminology items into the three categories:

- (1) sample & variables
- (2) hypothesis testing & statistical modelling
- (3) descriptive statistics

- |                           |                        |                       |                          |                             |
|---------------------------|------------------------|-----------------------|--------------------------|-----------------------------|
| • alternative hypothesis  | • degree of freedom    | • intercept           | • paired samples         | • single-sided test         |
| • anova                   | • dependent variable   | • IQR                 | • poisson                | • skewed data               |
| • barplot                 | • effect size          | • linear model        | • population             | • slope                     |
| • binary                  | • error                | • linear regression   | • predictor              | • standard deviation        |
| • binomial                | • explanatory variable | • logistic regression | • proportion             | • standard error            |
| • Bonferroni              | • factor               | • mean                | • $p$ -value             | • student $t$ -distribution |
| • boxplot                 | • Fisher's exact test  | • median              | • QQ-plot                | • treatment effect          |
| • categorical             | • histogram            | • multiple comparison | • quantile               | • $t$ -test                 |
| • Chisquare test          | • hypothesis testing   | • nominal             | • range                  | • two-sided test            |
| • confounding             | • hypothesis tests     | • normal              | • regression coefficient | • unpaired samples          |
| • contingency table       | • independent variable | • null hypothesis     | • residuals              | • variable                  |
| • continuous              | • integer              | • numeric             | • response               | • variance                  |
| • correlation coefficient | • interaction          | • observation         | • sample                 | • vector                    |
| • count                   |                        | • odds ratio          | • sampling variation     |                             |
| • data format             |                        | • ordinal             | • scatter plot           |                             |
| • data point              |                        | • outcome             | • significance           |                             |
| • data type               |                        |                       |                          |                             |

**Exercise 2: Getting to know R and chickwts**

- (a) Open R Studio.
- (b) Open a new R-Script.
- (c) Load data set chickwts.

```
# ?chickwts  
data(chickwts)  
head(chickwts)
```

**Exercise 3: Summary Statistics with chickwts**

- (a) Do summary statistics (numerically and graphically).
- (b) For advanced R users: Try an anova (are the assumptions fulfilled?) and a Tukey-Anscombe plot.  
Try a histogram with a density line on top. ...

**Exercise 4: Data import to R: perulung\_ems.csv**

- (a) Import the data set perulung\_ems.csv (taken from Kirkwood and Sterne, 2nd edition) into R.  
Data from a study of lung function among children living in a deprived suburb of Lima, Peru.  
Variables:
  - fev1: in liter, "Forced Expiratory Volume in 1 second" measured by a spirometer. This is the maximum volume of air which the children could breath out in 1 second
  - age: in years
  - height: in cm
  - sex: 0 = girl, 1 = boy
  - respsymp: respiratory symptoms experienced by the child over the previous 12 months
- (b) What *delimiter* do you need to choose?
- (c) Do summary statistics (numerically and graphically).
- (d) Plot a boxplot.

**Exercise 5: Defining a new data frame**

- (a) Create a data frame with 3 columns.

**Exercise 6: Get to know bacteria data set**

- (a) Install package MASS.
- (b) Load data set bacteria.
- (c) Describe in your own words what the data set bacteria contains.
- (d) Do summary statistic (numerically and graphically).

- (e) Select only observations collected during the second week.

### Exercise 7: Bracket types

What is conceptionally the difference between the bracket types [...] and (...)?

```
chickwts[, 2]  
summary(aov(weight ~ feed, data = chickwts))
```

### Exercise 8: Factor variables

- (a) How many levels has the factor variable `trt` from `bacteria`?
- (b) Define a new variable `trt.new` in which you combine the levels `drug` and `drug+` into one single level and label it as `treated`. The new variable `trt.new` should in the end have two levels: `placebo` and `treated`.
- (c) Do summary statistics for `placebo` and `treated` group.

### Exercise 9: Get to know `ToothGrowth` data set

- (a) Load data set `ToothGrowth`.
- (b) Do summary statistic (numerically and graphically).
- (c) Define additional column `dose.factor` by converting the numeric variable `dose` into a factor variable.
- (d) Are the tooth length measurements normally distributed within the treatment (`supp`: `VC` or `OJ`) and within in the different doses (`dose`: `0.5`, `1`, `2`)?

### Exercise 10: Summary Statistics

Apply the summary statistics to the `perulung_ems` and `ToothGrowth` data set.

### Exercise 11: Data Plausibility Checks

- (a) What can go wrong?
- (b) Identify different strategies for spotting these potential errors.
- Logical errors
  - Spelling mistakes
- (c) Import the data set `bacteria_plausibility_check.csv` to R.
- (d) Detect the **six** errors in the imported data set `bacteria_plausibility_check.csv` in R.
- (e) Find possible solutions in R how to handle these challenges.
- (f) Do all variables have the correct data type (numeric, integer, factor)? - If not, do correct / define them.

### Exercise 12: Missing Values

- (a) Check out the difference between the different missing values.

```

y1 <- c(2, 4, 3, NA, 6, 1)
y2 <- c("diseased", "healthy", NA, "NA")
y3 <- c(1, "NA", 0, 1, NaN)

#
is.na(y1)
which(is.na(y1))
is.na(y2)
which(is.na(y2))
is.na(y3)
which(is.na(y3))
is.nan(y3)

```

- (b) Create a vector with missing values and determine the mean and median.
- (c) If  $x = c(22, 3, 7, NA, NA, 67)$  what will be the output for the R statement `length(x)`?
- (d) If  $x = c(NA, 3, 14, NA, 33, 17, NA, 41)$  which line of R code removes all occurrences of NA in  $x$ .
- (e) If  $y = c(1, 3, 12, NA, 33, 7, NA, 21)$  what R statement will replace all occurrences of NA with 11?
- (f) If  $x = c(34, 33, 65, 37, 89, NA, 43, NA, 11, NA, 23, NA)$  then what will count the number of occurrences of NA in  $x$ ?
- (g) Create the vector  $x1$ . Then, find again the number of missing values and their position.

```
x1 <- c(rnorm(10,5,2), NA, 5:12, NA, 6, 7.5, NA)
```

- (h) Now, create the vector  $x2$  and assess the difference to  $x1$ .

```
x2 <- c(rnorm(10,5,2), NA, 5:12, NA, 6, 7.5, NA, log(-2))
```

- (i) What is the meaning of "NA" versus "NaN"?
- (j) Replace the missing values in  $x1$  with a 0. Check then that the NAs are no longer present. Try two different commands to coerce the NAs into 0.

### Exercise 13: Data import to R: `water_errors.csv`

- (a) Import the data set `water_errors.csv` to R: A data frame with 61 observations on the following 6 variables.
- **location:** a factor with levels `North` and `South` indicating whether the town is as north as Derby.
  - **town:** the name of the town.

- **mortality**: averaged annual mortality per 100.000 male inhabitants.
  - **hardness**: calcium concentration (in parts per million).
  - **smoker**: If there are any smokers living in town.
  - **num.of.cig**: In case, smokers live in town, what number of cigarettes do they smoke per day.
- (b) Detect the errors in the imported data set `water_errors.csv` in R.
- (c) Find possible solutions in R how to handle these challenges.
- (d) Do all variables have the correct data type (numeric, integer, factor)? - If not, do correct / define them.

#### Exercise 14: t-test in R

- (a) Load the data set `ToothGrowth` within R and apply the two-sided two sample t-test to suitable variables of the data set.

```
data(ToothGrowth)
```

- (b) Interpret the results.
- (c) Read in the data set `perulung_ems` and apply the two-sided t-test to suitable variables of the `perulung_ems` data set and interpret the results.

#### Exercise 15: Chi-square test in R

- (a) Apply the Chi-square test and the fisher exact test to the whole `bacteria` data set.
- (b) Apply the Chi-square test and the fisher exact test to the subset of `bacteria` containing only the observations taken in week 2 (cf. Exercise 3). Are there any issues?
- (c) Repeat this exercise by using the (previously defined) combined `trt.new` variable (cf. Exercise 5) with the two levels `treated` and `drug`.
- (d) Could you also obtain the odds ratios?
- (e) Try also a logistic regression in R. Ask Google for help!

#### Exercise 16: Outside plot frame

- (a) Type `demo(graphics)` in your console and press enter. This command shows you a nice demonstration of possible R graphics.

```
# After the demonstration us the following commands:  
dev.off()  
par(mfrow=c(1,1))
```

- (b) Change the x-axis and y-axis labelling of a boxplot plotting the `len` variable of the `ToothGrowth` data set.

```
data("ToothGrowth")
boxplot(ToothGrowth$len)
```

(c) How do you set a main title for your above plot?

(d) What does the following command do?

```
par(mfrow=c(2,2))
```

(e) We have six different feed types in `chickwts`. Try to plot two separate boxplots for casein and horsebean and set the same minimum and maximum for the y-axis. Use the function `subset` for doing so.

(f) How do you enlarge the font size of the axis as well as the axis labels of the following plot with the `perulung` data set?

```
lung <- read.csv("perulung_ems.csv", sep=";")
par(mfrow=c(1,1))
plot(lung$fev1, lung$height)
```

(g) Label the x-axis of the following plot with "Vitamin C in  $\mu\text{g}$ ". Use the greek letter for  $\mu$ .

```
plot(ToothGrowth$dose, ToothGrowth$len)
```

(h) Read <http://www.statmethods.net/advgraphs/parameters.html>.

### Exercise 17: Inside the square of the plot

(a) Type `demo(graphics)` in your console and press enter. This command shows you a nice demonstration of possible R graphics.

```
# After the demonstration us the following commands:
dev.off()
par(mfrow=c(1,1))
```

(b) Add a legend to the following barplot. Are there several different solutions for this?

```
library("MASS")
data("bacteria")
barplot(prop.table(table(bacteria$y, bacteria$trt),margin=1),
        beside=FALSE, ylim = c(0,0.8))
```

(c) Add a density line to this histogram.

```
hist(ToothGrowth$len, prob = TRUE, col = "grey", ylim = c(0, 0.05))
```

- (d) Add a **dotted red** linear regression line to the following plot.

```
plot(lung$height, lung$fev1)
```

- (e) Color the points in the following plot according to the sex variable.

```
plot(lung$height, lung$fev1)
```

- (f) Add two linear regression lines separately for female and male to the following plot.

```
plot(lung$height, lung$fev1)
```

- (g) Color the points in the following plot according to the supp variable. Use different point characters (pch) based on the supp variable.

```
plot(ToothGrowth$len, ToothGrowth$dose)
```

- (h) Read <http://www.statmethods.net/advgraphs/parameters.html>.

### Exercise 18:

- (a) Load the below data set and for further information check the command ?water.

```
# install.packages("HSAUR3")  
library("HSAUR3")  
data("water")  
str(water)  
head(water)  
summary(water)
```

- (b) Try to plot the variables mortality against hardness from the water data set.
- (c) Add a main title to the above plot (mortality against hardness).
- (d) Change the ...
- (a) font size of the axis annotation
  - (b) font size of the x- and y-axis labels
  - (c) the point sizes within the plot
- ... of the above plot (mortality against hardness).
- (e) Looking at the above plot: Do you think the two variables hardness and mortality correlate? What function do you use to find out the correlation coefficient? Do they have a positive or a negative correlation coefficient? How do you interpret the correlation coefficient in your own words?

- (f) In the water data set, can you graphically find out if there is a difference between the the two variables hardness and mortality conditional on the location (North, South).
- (g) Add a legend to the above plot so that you can easily differentiate the locations (North or South) of the observations.
- (h) Do a barplot of the variable location from the water data set.
- (i) ADDITIONAL: Try if any of these following plotting functions can be applied to the data sets perulung or ToothGrowth.

### Exercise 19: Anova in R

- (a) Open the .R file ANOVA\_with\_chickwts.R from your RCourse folder and have another look on how we applied the anova to the chickwts data set. Check line for line.
- (b) Load the ToothGrowth data set into R and encode the numeric variable dose as a factor variable. Define the new factor variable as dose.factor with the three levels low, med and high and add it to the data frame of ToothGrowth.

```
data(ToothGrowth)
```

- (c) Visualize the variable len per dose.factor level in a boxplot.
- (d) With the help of the R-commands written in the ANOVA\_with\_chickwts.R file, apply a analysis of variance (ANOVA) to the data set ToothGrowth

### Exercise 20: Linear Regression Model I

- (a) Reuse the commands from the lecture slides to fit a simple as well as a multiple linear regression model to the data set of perulung\_ems. Use fev1 as your response variable  $y$ .
- (b) Check the model assumptions.
- (c) Which model is best?

### Exercise 21: Linear Regression Model II

- (a) Load the ToothGrowth data set and run the following four linear regression models.

```
data(ToothGrowth)
ToothGrowth$dose.factor <- factor(ToothGrowth$dose, levels = c(0.5, 1.0, 2.0),
                                  labels = c("low", "med", "high"))
mod1 <- lm(len ~ dose.factor, data = ToothGrowth)
mod2 <- lm(len ~ supp, data = ToothGrowth)
mod3 <- lm(len ~ dose.factor + supp, data = ToothGrowth)
mod4 <- lm(len ~ dose.factor*supp, data = ToothGrowth)
```



- (b) Have a look at the summary of these models.
- (c) How do you interpret the model coefficients?
- (d) Which model is best?

### Exercise 22: Linear Regression Model III

- (a) Load the `water` data set and fit a multiple linear regression model. Use `mortality` as your response variable and add `hardness` and `location` as an explanatory variable.
- (b) Check the underlying model assumptions.
- (c) Add an interaction term between `hardness` and `location` to the above estimated multiple linear regression model.
- (d) Interpret the interaction coefficient `hardness:locationSouth`.
- (e) Check the underlying model assumptions.
- (f) Which one is the better model? With or without the interaction term?
- (g) How to derive confidence intervals for the regression coefficient of `hardness` and `location`?

### Exercise 23: Combining everything we have learnt

Hypothetical example from Kirkwood and Sterne, Medical Statistics, 2nd ed., p. 177

- (a) Read in the data set `lepto`. This study presents a serology survey of leptospira sero-prevalence in rural and urban areas of the west indies.
- (b) Encode the numeric variable `antibodies` as a factor with levels 0 and 1.
- (c) Make a crosstable with the risk factor `exposure` and `antibodies`.
- (d) Run a Chi-squared test, a Fisher's exact test and a logistic regression (`glm`) to assess if the exposure (living in rural vs. urban areas) is a risk factor.
- (e) Create a subset for `male` and `female` based on the variable `gender`.
- (f) Repeat the crosstable (2-by-2 table), Chi-squared test, Fisher's exact test and a logistic regression (`glm`) for the subsets **separately**.
- (g) Does the conclusion of your research question change with the analysis of the subsets? (Research question: Is the exposure (rural and urban areas) a risk factor?)
- (h) Fit a logistic regression model (`glm`) with `exposure` and `gender` as explanatory variables.