

L_p MINKOWSKI PROBLEM AND BRUNN-MINKOWSKI INEQUALITY FOR DUAL QUERMASSEINTEGRALS

XIAOJUAN CHEN, SHENGYU TANG, AND SINAN WANG

ABSTRACT. This paper studies the core problems in the L_p dual Brunn-Minkowski theory, encompassing the L_p Minkowski problem and L_p Brunn-Minkowski inequality for dual quermassintegrals. For the case $0 < p < q \leq n$, we establish C^0 estimates for the L_p dual Minkowski problem without symmetric assumptions, thereby resolving a related problem proposed by Böröczky-Chen-Liu-Saroglou in the smooth sense. We further prove the uniqueness of smooth solutions under appropriate conditions, provided the density function is sufficiently close to a constant in the Hölder norm. Finally, exploiting the fact that the uniqueness of the Minkowski type problem is equivalent to the validity of the Brunn-Minkowski inequality in a certain sense, we study the L_p Brunn-Minkowski inequality for dual quermassintegrals for origin-symmetric convex bodies with $p < q$.

1. INTRODUCTION

Convex geometric analysis, the analytically oriented branch of convex geometry, is built in part on the classical Brunn-Minkowski (or mixed volume) theory developed by Minkowski, Alexandrov, and other pioneers. In this classical framework, the quermassintegrals are core geometric invariants, which correspond to integrals of mean curvature for closed convex hypersurfaces in differential geometry, as well as integrals of projection areas in integral geometry. Viewed as functionals on the space of convex bodies in $\mathbb{R}^n$, the derivatives of the quermassintegrals yield surface area measures, whose characterization is a foundational problem in the field, epitomized by the Minkowski problem that links convex geometry to fully nonlinear partial differential equations. Here, tools such as the cosine transform (a variant of the Fourier transform) and elliptic partial differential equations on the unit sphere play crucial roles in the study of projections and surface area measures. Motivated by research on convex body sections in integral geometry, an analogous theory for mixed volumes was introduced in the 1970s by Lutwak [41], which revealed a striking duality in convex geometry and thus became known as the dual mixed volume theory, or

2020 *Mathematics Subject Classification.* 52A38, 35J60.

Key words and phrases. dual quermassintegral, Brunn-Minkowski inequality, L_p dual Minkowski problem.

the dual Brunn-Minkowski theory. The central invariants of this dual theory are the dual quermassintegrals, special dual mixed volumes where volume serves as both the classical and dual quermassintegrals.

1.1. Minkowski problem. The classical Minkowski problem is essentially the characterization of surface area measures, which was first studied by Minkowski for polytopes in 1897. In 1962, Firey [24] generalized the Minkowski sum to p -sum. Later, Lutwak [42] developed the corresponding L_p Brunn-Minkowski theory based on Firey's p -sum, and proposed L_p surface area measures, then utilized Aleksandrov's variational theory to solve the corresponding L_p Minkowski problem. The L_p Minkowski problem encompassed many important issues. When $p = 1$, it corresponds to the classical Minkowski problem [1, 44]; when $p = 0$, it is called the logarithmic Minkowski problem [9, 54]; when $p = -n$, it is known as the centro-affine Minkowski problem [21, 55].

The above Minkowski problems were proposed regarding the surface area measure. In convex geometry, there is another important measure, namely the curvature measure. The most typical curvature measure is the Aleksandrov integral curvature measure. Aleksandrov [2] obtained the existence of the corresponding solution to the Aleksandrov problem. In 2016, a breakthrough was made when Huang-Lutwak-Yang-Zhang [32] discovered the q -th dual curvature measure $\tilde{C}_q(K, \cdot)$, which can be viewed as differentials of dual quermassintegrals, and derived the related variational formulas. Then they proposed and solved the corresponding dual Minkowski problem. Subsequently, the dual Brunn-Minkowski theory flourished and achieved a series of significant results, see [10, 27, 29, 30, 40, 52, 53] and the references therein.

Lutwak-Yang-Zhang [43] further integrated the dual Brunn-Minkowski theory into the L_p Brunn-Minkowski theory, and introduced the L_p dual curvature measure $\tilde{C}_{p,q}(K, \cdot)$ to unify the L_p surface area measure and the q -th dual curvature measure. The associated Minkowski problem was called the L_p dual Minkowski problem. Subsequently, many researchers adopted variational methods to derive solutions with variational structure: Böröczky-Fodor [6] obtained the existence when $p > 1, q > 0, p \neq q$; Chen-Chen-Li [17] studied the existence of the solution under even assumptions when $q > 0, -q^* < p < 0$; Li-Liu-Lu [39] constructed counterexamples to demonstrate the nonuniqueness when $p < 0 < q$; Mui [45] studied the existence of the solution under even assumptions when $-1 < p < 0, q < 1 + p, p \neq q$, and was further extended by Böröczky-Kovács-Mui-Zhang [7] under group symmetric assumptions. After that, Shan [50] further established the existence of solutions from an algebraic perspective for group invariant measures and group invariant convex bodies when $p, q \in \mathbb{R}$.

When the L_p dual curvature measure has a density $\frac{d\tilde{C}_{p,q}(K, \cdot)}{d\mathcal{H}^{n-1}} = f$ with respect to spherical Lebesgue measure $\mathcal{H}^{n-1}$, then this problem is equivalent to solving the

following Monge-Ampère equation

$$h^{1-p}(h^2 + |\nabla h|^2)^{\frac{q-n}{2}} \det(\nabla^2 h + hI) = f \quad \text{on } S^{n-1}, \quad (1.1)$$

where ∇h and $\nabla^2 h$ denote the spherical gradient and the Hessian of the function h on S^{n-1} with respect to a moving orthonormal frame respectively, and I is the identity matrix. For equation (1.1), Huang-Zhao [34] used the continuity method to obtain the existence and the uniqueness of smooth solutions when $p > q$; Chen-Huang-Zhao [16] used the curvature flow method to obtain the existence of solutions when $pq > 0$ and f is even; Chen-Li [18] used the curvature flow method to obtain the existence and the uniqueness of solutions when $p > 0, q \neq n$ or $q \leq p < 0$.

In this paper, we mainly focus on the uniqueness of the L_p dual Minkowski problem when the density function is near the ball in C^α norm sense. The uniqueness of this viewpoint has become a hot research topic recently. Chen-Huang-Li-Liu [20] established the uniqueness of L_p Minkowski problem when $p \in (1 - \frac{c}{n^{\frac{3}{2}}}, 1)$ and the density function is even and close to the ball. Chen-Feng-Liu [19] further studied the uniqueness for logarithmic Minkowski problem when $n = 3$ without any symmetric assumptions. For higher dimensions, the uniqueness results were independently established by Böröczky-Saroglou [11] and Hu-Ivaki [31] using different approaches. Hu [30] established the uniqueness for symmetric solutions of the dual Minkowski problem when the density function is even, close to the ball, and $0 < q \leq n$ for $1 \leq n \leq 3$ or $n - 3 \leq q \leq n$ for $n > 3$. Cabezas-Moreno and Hu [13] proved the uniqueness for symmetric solutions of the L_p dual Minkowski problem when $1 < p < q \leq n$, which also developed by Böröczky-Chen-Liu-Saroglou [4] for $p > -1, q > 0, p < q < \min\{n, n + p\}$ with symmetric assumptions and for $p \in (-1, 1), q$ is close to n without symmetric assumptions.

It is worth noting that C^0 estimates are important to derive existence and uniqueness results for the above Minkowski problems and the corresponding Monge-Ampère equations. Recently, Böröczky-Chen-Liu-Saroglou [5] proposed the following problem for the L_p dual Minkowski problem.

Problem 1.1 (BCLS [5]). *For $n \geq 2, p \in (-1, 1), q > n - 1$ and $\lambda > 1$, does there exist a constant $C = C(n, p, q, \lambda) > 1$ such that if*

$$\lambda^{-1} \mathcal{H}^{n-1} \leq \tilde{C}_{p,q}(K, \cdot) \leq \lambda \mathcal{H}^{n-1}$$

holds on S^{n-1} for a convex body $K \in \mathcal{K}_o^n$, then the support function h_K of K satisfies

$$\max_{x \in S^{n-1}} h_K(x) \leq C \quad \text{and} \quad |K| \geq \frac{1}{C}.$$

In [5], they solved Problem 1.1 for $p \in [0, 1), q > p + 2$ in $\mathbb{R}^3$. They also verified Problem 1.1 in [4] when $n = 3, 4$ and q is very close to n , and higher dimension for

$-1 < p < q < \min\{n, n+p\}$ with $q > 0$ under symmetric assumptions. In our paper, we first answer Problem 1.1 when $q \in (1, n]$, $p \in (0, q)$ and the given data is smooth.

Theorem 1.2. *For $n \geq 2$, $q \in (1, n]$ and $p \in (0, q)$, let f be a smooth function on S^{n-1} . There exists a constant C depending on n, p, q, λ such that if*

$$\lambda^{-1} \mathcal{H}^{n-1} \leq \tilde{C}_{p,q}(K, \cdot) \leq \lambda \mathcal{H}^{n-1}$$

with $d\tilde{C}_{p,q}(K, \cdot) = f d\mathcal{H}^{n-1}$ for any $K \in \mathcal{K}_o^n$, then

$$\max_{x \in S^{n-1}} h_K(x) \leq C \text{ and } |K| \geq \frac{1}{C}.$$

Using the similar argument of Böröczky-Chen-Liu-Saroglou [4], we deduce the following uniqueness results. Along the way, we also use the uniqueness results of (1.1) in Chen-Li [18] when $f \equiv 1$ and $1 < p < q \leq n$.

Theorem 1.3. *Suppose $n \geq 2$, $\alpha \in (0, 1)$, $q \in (1, n]$, $p \in (0, q)$. There exists a constant $\epsilon > 0$ depending only on n, α, p and q such that, if f is a smooth function on S^{n-1} satisfying $\|f - 1\|_{C^\alpha(S^{n-1})} < \epsilon$, and*

- either, $p \in (0, 1)$ and $q \in (n - \epsilon, n]$,
- or $1 < p < q \leq n$,

then equation (1.1) has a unique smooth solution in $\mathcal{K}_o^n$.

1.2. Brunn-Minkowski inequality. The Brunn-Minkowski inequality, as another important topic in Brunn-Minkowski theory, states that for given convex bodies K and L in $\mathbb{R}^n$,

$$V_n((1 - \lambda)K + \lambda L)^{\frac{1}{n}} \geq (1 - \lambda)V_n(K)^{\frac{1}{n}} + \lambda V_n(L)^{\frac{1}{n}}, \quad \forall \lambda \in [0, 1],$$

with equality if and only if K and L are homothetic. This is equivalent to the log-concavity of the volume functional

$$V_n((1 - \lambda)K + \lambda L) \geq V_n(K)^{1-\lambda} V_n(L)^\lambda, \quad \forall \lambda \in [0, 1].$$

The relationship between the Brunn-Minkowski inequality and other inequalities in geometry and analysis, along with some applications, refer to the survey article by Gardner [25].

In 1962, Firey [24] generalized the Minkowski sum of convex bodies to the L_p -sum $(1 - \lambda)K +_p \lambda L$. When $p \geq 1$, it is defined as the convex body with support function $((1 - \lambda)h_K^p + \lambda h_L^p)^{\frac{1}{p}}$. Firey also established the corresponding L_p Brunn-Minkowski inequality

$$V_n((1 - \lambda)K +_p \lambda L)^{\frac{p}{n}} \geq (1 - \lambda)V_n(K)^{\frac{p}{n}} + \lambda V_n(L)^{\frac{p}{n}}, \quad \forall \lambda \in [0, 1], \quad (1.2)$$

with equality if and only if $K = L$.

A breakthrough came from Böröczky-Lutwak-Yang-Zhang [8], which studied the L_p Brunn-Minkowski inequality for $p \in [0, 1)$, covering the previously missing range. It is well-known that $((1-\lambda)h_K^p + \lambda h_L^p)^{\frac{1}{p}}$ cannot be a support function for any convex body when $p \in [0, 1)$. However, they found a natural generalization of L_p -sum as follows

$$(1-\lambda)K +_p \lambda L = \cap_{x \in \mathbb{S}^{n-1}} \{z \in \mathbb{R}^n : x \cdot z \leq ((1-\lambda)h_K^p(x) + \lambda h_L^p(x))^{\frac{1}{p}}\}.$$

When $p = 0$, it is actually the log Minkowski sum,

$$(1-\lambda)K +_0 \lambda L = \cap_{x \in \mathbb{S}^{n-1}} \{z \in \mathbb{R}^n : x \cdot z \leq h_K(x)^{1-\lambda} h_L(x)^\lambda\},$$

where K and L are convex bodies that contain the origin in their interiors. Meanwhile, when $p \in (0, 1)$, for two origin-symmetric convex bodies K and L in $\mathbb{R}^n$, they conjectured the L_p Brunn-Minkowski inequality (1.2) and the log Brunn-Minkowski inequality when $p = 0$,

$$V_n((1-\lambda)K +_0 \lambda L) \geq V_n(K)^{1-\lambda} V_n(L)^\lambda \quad \forall \lambda \in [0, 1].$$

Recently, Kolesnikov-Milman [38] introduced a local perspective: instead of proving the inequality for all convex bodies directly, they studied its infinitesimal version. Hence, they developed a local L_p Brunn-Minkowski inequality for $p \in [1 - \frac{c}{n^{\frac{3}{2}}}, 1)$, which stated that when $K, L \in \mathcal{K}_{+,e}^2$ satisfies

$$(1-\lambda)K +_p \lambda L \in \mathcal{K}_{+,e}^2, \quad \forall \lambda \in [0, 1],$$

where $\mathcal{K}_{+,e}^2$ is the class of origin-symmetric convex bodies with C^2 smooth boundary and strictly positive curvature, then (1.2) holds for $n \geq 2$. Moreover, Chen-Huang-Li-Liu [20] gave the equivalence of the L_p Brunn-Minkowski inequality (1.2) to the local version, by adapting the PDE methods. This result showed that studying the local inequality is not merely a technical simplification but a legitimate route to the global conjecture. It is worth to note that the local inequality could deduces local uniqueness for the even L_p Minkowski problem. Putterman [47] also proved the equivalence between local and global from a geometric perspective.

Corresponding to the near spherical uniqueness of Minkowski problem discussed in the preceding context, we further investigate the Brunn-Minkowski inequality in the near spherical setting. The study of near spherical Brunn-Minkowski inequality can be traced back to the work in Colesanti-Livshyts-Marsiglietti [22]. They investigated the Brunn-Minkowski inequality (and log version) for radially symmetric log-concave measures when the given two convex bodies are near the ball. Bianchini-Colesanti-Pagnini-Roncoroni [3] verified the validity of the L_p Brunn-Minkowski inequality for quermassintegrals of symmetric convex bodies in $\mathbb{R}^n$, under the condition that one body is the ball and the other one lies in a small neighborhood of the ball. Subsequently, Patsalos-Saroglou [46] improved the results in [3] to a symmetric convex

body near the ball, while the other one is arbitrary symmetric convex body. Furthermore, they proved the near spherical uniqueness of the corresponding Minkowski problem based on the near spherical Brunn-Minkowski inequality.

The Brunn-Minkowski inequality for dual quermassintegrals was conjectured by Lutwak and has been presented in various talks and seminars by the Lutwak-Yang-Zhang's group and their collaborators. It states that if K and L are convex bodies, then for $q > 0$,

$$\tilde{V}_q((1-\lambda)K + \lambda L)^{\frac{1}{q}} \geq (1-\lambda)\tilde{V}_q(K)^{\frac{1}{q}} + \lambda\tilde{V}_q(L)^{\frac{1}{q}}, \quad \forall \lambda \in [0, 1], \quad (1.3)$$

with equality if and only if K and L are dilates of each other. One can also see these details in [33]. Xi-Zhang [51] solved the conjecture when $q \in (0, 1]$ from a geometric way. Recently, based on the analytic tools developed by Kolesnikov-Milman [36, 37], Kolesnikov-Livshyts [35], and the estimate from Cordero-Erausquin and Rotem [23], Sadovskiy-Zhang [48] confirmed the conjecture (1.3) for $0 < q \leq n$ under symmetric assumptions, without the characterization of equality.

We mainly focus on the L_p Brunn-Minkowski inequality for dual quermassintegral, which is considered by Xi-Zhang [51] that, if it holds for any convex bodies K and L containing the origin in their interiors that

$$\tilde{V}_q((1-\lambda)K +_p \lambda L)^{\frac{p}{q}} \geq (1-\lambda)\tilde{V}_q(K)^{\frac{p}{q}} + \lambda\tilde{V}_q(L)^{\frac{p}{q}}, \quad \forall \lambda \in [0, 1] \quad (1.4)$$

for $q \neq 0$. They answered this question when $p \geq q$. The case $p < q$ is likewise both interesting and challenging, which constitutes the main focus of our final result.

For the case $p \geq 1$, we employ a classic mathematical argument to tackle the problem. This result follows naturally from the Brunn-Minkowski inequality for $p = 1$, i.e. the results established by Sadovskiy-Zhang [48]. This also explain the reason that our constraints on the parameter q and convex bodies coincide with those in their work. For $p \in (0, 1)$, we use the near spherical uniqueness of Minkowski problem to prove the near spherical Brunn-Minkowski inequality for dual quermassintegrals. We first prove the equivalence between the L_p Brunn-Minkowski inequality and the L_p Minkowski inequality for dual quermassintegral (the explicit form of the L_p Minkowski inequality is given in (5.3)). Building upon this equivalence, we show the L_p Minkowski inequality by using the near spherical uniqueness of the L_p Minkowski problem, and this argument requires one of the convex bodies to be sufficiently close to a ball. It is worth noting that [46] adopted the inverse direction of perspective to this paper, utilizing the Brunn-Minkowski inequality to establish the uniqueness of solutions to the Minkowski problem. In a certain sense, this phenomenon boils down that the uniqueness of the Minkowski problem is essentially equivalent to the validity of the Brunn-Minkowski inequality. Our results are stated as follows.

Theorem 1.4. *Let $n \geq 2$. Then for $K, L \in \mathcal{K}_e^n$, (1.4) holds when*

- either, $p \geq 1$ and $q \in (0, n]$,
- or, $p \in (0, 1)$, $q \in (p, n]$, $K, L \in \mathcal{K}_{+,e}^{2,\alpha}$, and for some $\epsilon > 0$, $\|h_K - 1\|_{C^{2,\alpha}} \leq \epsilon$.

This paper is organized as follows. In Section 2, we present some preliminaries for basic notations and convex bodies. For the L_p dual Minkowski problem, in Section 3, we establish C^0 estimates when $p \in (0, q)$, $q \in (1, n]$, which is Theorem 1.2, and in Section 4, we derive the uniqueness results when the density function near the ball and prove Theorem 1.3. In Section 5, we study the L_p dual Brunn-Minkowski inequality (1.4) and give the proof of Theorem 1.4.

2. PRELIMINARIES

In this section, some notations and introduction about convex bodies will be provided (see excellent reference [26, 49]) .

2.1. Basic Notations. Let $\mathbb{R}^n$ be the n -dimensional Euclidean space. The unit sphere in $\mathbb{R}^n$ is defined as S^{n-1} . $C(S^{n-1})$ represents the space of continuous functions on S^{n-1} , and will always be viewed as equipped with the max-norm metric

$$\|f - g\|_\infty = \max_{x \in S^{n-1}} |f(x) - g(x)|,$$

for $f, g \in C(S^{n-1})$. Denote by $C^+(S^{n-1})$ the set of positive continuous functions on S^{n-1} , and by $C_e^+(S^{n-1})$ the set of positive continuous even functions on S^{n-1} .

A convex body K in $\mathbb{R}^n$ is a compact convex set with nonempty interior. Let $\mathcal{K}_0^n$ be a class of convex bodies that contain the origin in their interiors in $\mathbb{R}^n$, $\mathcal{K}_e^n$ be a class of origin-symmetric convex bodies in $\mathbb{R}^n$, $\mathcal{K}_{+,e}^2$ be the subclass of $\mathcal{K}_e^n$ with C^2 smooth boundary and strictly positive curvature in $\mathbb{R}^n$, and $\mathcal{K}_{+,e}^{2,\alpha} \subset \mathcal{K}_{+,e}^2$ with $C^{2,\alpha}$ smooth boundary.

2.2. Convex bodies. Let K be a compact convex subset in $\mathbb{R}^n$, then the support function h_K of K is defined by

$$h_K(x) = \max\{x \cdot y : y \in K\}, \quad x \in \mathbb{R}^n. \quad (2.1)$$

The support function $h = h_K : \mathbb{R}^n \rightarrow \mathbb{R}$ is a continuous function homogeneous of degree 1. Suppose K contains the origin in its interior, then the radial function $\rho = \rho_K : \mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{R}$ is defined by

$$\rho_K(x) = \max\{\lambda : \lambda x \in K\}, \quad x \in \mathbb{R}^n \setminus \{0\}. \quad (2.2)$$

The radial function ρ_K is a continuous function homogeneous of degree -1 .

For each $f \in C^+(S^{n-1})$, the Wulff shape $[f]$ generated by f is the convex body defined by

$$[f] = \{x \in \mathbb{R}^n : x \cdot u \leq f(u), \text{ for all } u \in S^{n-1}\}.$$

It is obvious that $h_{[f]} \leq f$ and $[h] = K$ for each $K \in \mathcal{K}_0^n$.

Suppose $K_i \in \mathcal{K}_0^n$ is a sequence of convex bodies in $\mathbb{R}^n$, we say K_i converges to a compact convex subset $K \subset \mathbb{R}^n$ with respect to the Hausdorff metric provided that

$$\|h_{K_i}(u) - h_K(u)\|_\infty = \max_{u \in S^{n-1}} |h_{K_i}(u) - h_K(u)| \rightarrow 0, \quad i \rightarrow \infty.$$

If K contains the origin in its interior, the above formula is equivalent to

$$\|\rho_{K_i}(u) - \rho_K(u)\|_\infty = \max_{u \in S^{n-1}} |\rho_{K_i}(u) - \rho_K(u)| \rightarrow 0, \quad i \rightarrow \infty.$$

We use ν_K to denote the Gauss map that takes $x \in \partial K$ to its unique outer unit normal which is due to the convexity of K , and the map ν_K is almost everywhere defined on ∂K . We use ν_K^{-1} to denote the inverse Gauss map. Since K is not assumed to be strictly convex, the map ν_K^{-1} is set-valued map and for each set $\eta \subset S^{n-1}$,

$$\nu_K^{-1}(\eta) = \{x \in \partial K : \nu_K(x) \in \eta\}.$$

We employ the following renormalization for the Gauss and inverse Gauss map to enhance clarity. For $u \in S^{n-1}$, if ν_K is well-defined at $\rho_K(u)u \in \partial K$, then we write $\alpha_K(u)$ for $\nu_K(\rho_K(u)u)$. The reverse radial Gauss mapping α_K^{-1} is defined as follow:

$$\alpha_K^{-1}(\eta) = \{u \in S^{n-1} : \nu_K(\rho_K(u)) \in \eta\} \quad \text{for any } \eta \in S^{n-1}.$$

It is well known that the Jacobian of α_K^{-1} is

$$|\text{Jac} \alpha_K^{-1}| = \frac{h \det(\nabla^2 h + hI)}{(h^2 + |\nabla h|^2)^{\frac{n}{2}}}.$$

3. A PRIORI ESTIMATES

In this section, we provided uniform C^0 estimates of solutions to the L_p dual Minkowski problem with $p \in (0, q)$ and $q \in (1, n]$, which is Theorem 1.2.

Lemma 3.1. *Assume that $h \in C^3(S^{n-1})$ is a solution of (1.1) for $q \in (1, n]$ and $p \in (0, q)$, then there is a positive constant $0 < \alpha < 1$ such that*

$$\frac{|\nabla h|}{h^\alpha} \leq C, \tag{3.1}$$

where C depends only on $\|f\|_{C^1(S^{n-1})}$, $\min_{S^{n-1}} f$, $\max_{S^{n-1}} f$, α, n, p and q .

Proof. Consider the testing function $G = \frac{|\nabla h|}{h^\alpha}$, at its maximum $u_0 \in S^{n-1}$. Without loss of generality, we choose normal coordinates at u_0 such that

$$h_1 = |\nabla h| \tag{3.1a}$$

$$h_i = 0 \quad \text{for } i = 2, \dots, n-1. \tag{3.1b}$$

Writing equation (1.1) as

$$\log \det(h_{ij} + h\delta_{ij}) = \log \det(w_{ij}) = (p-1) \log h + \frac{n-q}{2} \log(h^2 + |\nabla h|^2) + \log f,$$

and differentiating it in direction $e_1 \in T_{u_0}S^{n-1}$, we obtain

$$w^{ij}w_{ij1} = \frac{f_1}{f} + (p-1)\frac{h_1}{h} + (n-q)\frac{hh_1 + h_1h_{11}}{h^2 + h_1^2}.$$

At u_0 ,

$$0 = (\log G)_i = \frac{h_i h_{1i}}{|\nabla h|^2} - \alpha \frac{h_i}{h} = \frac{h_{1i}}{h_1} - \alpha \frac{h_i}{h},$$

i.e.

$$h_{11} = \alpha \frac{h_1^2}{h}, \quad h_{1i} = 0 \quad \text{for } i \geq 2. \quad (3.2)$$

We may as well assume that $[h_{ij}(u_0)]$ is diagonal. Noticing that $(\log G)_{ij}$ is nonpositive definite at u_0 ,

$$\begin{aligned} 0 &\geq w^{ij}(\log G)_{ij} = w^{ij}\left(\frac{h_i h_{1j}}{h_1^2} + \frac{h_{1ij}}{h_1} - \frac{2h_{1i}h_{1j}}{h_1^2} - \frac{\alpha h_{ij}}{h} + \frac{\alpha h_i h_j}{h^2}\right) \\ &\geq w^{ij}\left(\frac{h_{1ij}}{h_1} - \frac{h_{1i}h_{1j}}{h_1^2} - \frac{\alpha h_{ij}}{h} + \frac{\alpha h_i h_j}{h^2}\right) \\ &= \frac{1}{h_1}w^{ij}w_{ij1} - w^{11} + (\alpha - \alpha^2)w^{11}\frac{h_1^2}{h^2} - \frac{\alpha(n-1)}{h} + \alpha \sum_i w^{ii}, \end{aligned}$$

where we have used $h_{1ij} = w_{1ij} - \delta_{1i}h_j = w_{ij1} - \delta_{1i}h_j$ and $w_{ij} = h_{ij} + h\delta_{ij}$. Since

$$\begin{aligned} w^{11} &= \frac{1}{h_{11} + h} = \frac{h}{\alpha h_1^2 + h^2} \\ w^{ij}w_{ij1} &= \frac{(p-1)h_1}{h} + \frac{f_1}{f} + (n-q)\frac{hh_1 + h_1h_{11}}{h^2 + h_1^2}. \end{aligned}$$

Thus for $q \leq n$,

$$\begin{aligned} 0 &\geq \frac{p-1}{h} + \frac{n-q}{h} \frac{h^2 + \alpha h_1^2}{h^2 + h_1^2} + \frac{f_1}{fh_1} + \frac{(\alpha - \alpha^2)h_1^2 - h^2}{\alpha h h_1^2 + h^3} - \frac{\alpha(n-1)}{h} + \alpha \sum_i w^{ii} \\ &\geq \frac{p-1}{h} + \frac{f_1}{fh_1} + \frac{(\alpha - \alpha^2)h_1^2 - h^2}{\alpha h h_1^2 + h^3} - \frac{\alpha(n-1)}{h} + \alpha \sum_i w^{ii}. \end{aligned} \quad (3.3)$$

We divide it into two cases to deal with the inequality according to value of $h(u_0)$. It is to say that there exists a positive constant M depending only on p, q, n, f and to be determined such that

- Case 1: $h(u_0) \leq M$;
- Case 2: $h(u_0) \geq M$.

Case 1: $h(u_0) \leq M$: Subcase(1_A): $h_1 \leq \frac{h}{\alpha}$, then $\frac{h_1}{h^\alpha} \leq \frac{h^{1-\alpha}}{\alpha} \leq \frac{M^{1-\alpha}}{\alpha}$.

Subcase(1_B): $h_1 \geq \frac{h}{\alpha}$, then $\frac{h^2}{\alpha h_1^2 + h^2} = \frac{1}{\alpha(\frac{h_1}{h})^2 + 1} \leq \frac{\alpha}{1+\alpha} < \alpha$, combining this and (3.3), we have

$$\begin{aligned}
0 &\geq \frac{p-1}{h} + \frac{f_1}{fh_1} + \frac{(\alpha - \alpha^2)h_1^2 - h^2}{\alpha h h_1^2 + h^3} - \frac{\alpha(n-1)}{h} + \alpha \sum_i w^{ii} \\
&= \frac{p-1}{h} + \frac{f_1}{fh_1} + \frac{1}{h} \left[1 - \alpha - \frac{(2-\alpha)h^2}{\alpha h_1^2 + h^2} \right] - \frac{\alpha(n-1)}{h} + \alpha \sum_i w^{ii} \\
&\geq \frac{1}{h} [p - (2+n)\alpha + \alpha^2] + \frac{f_1}{fh_1}.
\end{aligned} \tag{3.4}$$

Here we set $p \geq (n+3)\alpha$, then it yields

$$0 \geq \frac{\alpha^2}{h} - \frac{|\nabla f|}{fh_1},$$

so

$$\frac{h_1}{h^\alpha} \leq \frac{|\nabla f|}{\alpha^2 f} M^{1-\alpha}. \tag{3.5}$$

Case 2: $h(u_0) \geq M$. We divide Case 2 in three subcases as follows:

Subcase(2_A): $h_1 \geq C_1 h$, where C_1 is a positive constant, and chosen so that

$$\frac{\alpha(n-1)}{2} C_1^{\frac{q-1}{n-1}} \left(1 + \frac{1}{C_1^2} \right)^{-\frac{n-q}{2(n-1)}} M^{\frac{q-p}{n-1}} (\max f)^{-\frac{1}{n-1}} > \max \left\{ \frac{2-\alpha}{C_1 \alpha}, \frac{|\nabla f|}{f} \right\}. \tag{3.6}$$

Since $h_1 \geq C_1 h$, thus

$$\frac{h}{\alpha h_1^2 + h^2} \leq \frac{h_1}{C_1(\alpha h_1^2 + h^2)} \leq \frac{1}{C_1 h_1} \frac{h_1^2 + \frac{1}{\alpha} h^2}{\alpha h_1^2 + h^2} = \frac{1}{\alpha C_1 h_1}. \tag{3.7}$$

Combining (3.7) and (3.3), we deduce that

$$\begin{aligned}
0 &\geq \frac{p-1}{h} + \frac{f_1}{fh_1} + \frac{1-\alpha}{h} - (2-\alpha) \frac{h}{\alpha h_1^2 + h^2} - \frac{\alpha(n-1)}{h} + \alpha \sum_i w^{ii} \\
&\geq \frac{p-n\alpha}{h} + \left(\frac{f_1}{fh_1} + \frac{\alpha}{2} \sum_i w^{ii} \right) + \left(\frac{\alpha}{2} \sum_i w^{ii} - \frac{2-\alpha}{\alpha C_1 h_1} \right),
\end{aligned}$$

which will lead to a contradiction if

$$\left(\frac{f_1}{fh_1} + \frac{\alpha}{2} \sum_i w^{ii} \right) + \left(\frac{\alpha}{2} \sum_i w^{ii} - \frac{2-\alpha}{\alpha C_1 h_1} \right) > 0. \quad (3.8)$$

Indeed, (3.2) tells us that $w_{ij} = h_{ij} + h\delta_{ij}$ may be diagonal at u_0 ,

$$\sum_i w^{ii} \geq (n-1) \det(w_{ij})^{-\frac{1}{n-1}} = (n-1) (h^{p-1} (h^2 + h_1^2)^{\frac{n-q}{2}} f)^{-\frac{1}{n-1}}. \quad (3.9)$$

Then (3.6), (3.9) and the fact $\frac{h_1}{C_1} \geq h \geq M$ give that

$$\begin{aligned} \frac{f_1}{fh_1} + \frac{\alpha}{2} \sum_i w^{ii} &\geq -\frac{|\nabla f|}{fh_1} + \frac{\alpha(n-1)}{2} (h^{p-1} (h^2 + h_1^2)^{\frac{n-q}{2}} f)^{-\frac{1}{n-1}} \\ &\geq \frac{1}{h_1} \left(-\frac{|\nabla f|}{f} + \frac{\alpha(n-1)}{2} h_1 h^{-\frac{p-1}{n-1}} \left(1 + \frac{1}{C_1^2} \right)^{-\frac{n-q}{2(n-1)}} h_1^{-\frac{n-q}{n-1}} f^{-\frac{1}{n-1}} \right) \\ &\geq \frac{1}{h_1} \left(-\frac{|\nabla f|}{f} + \frac{\alpha(n-1)}{2} C_1^{\frac{q-1}{n-1}} \left(1 + \frac{1}{C_1^2} \right)^{-\frac{n-q}{2(n-1)}} M^{\frac{q-p}{n-1}} f^{-\frac{1}{n-1}} \right) > 0 \end{aligned}$$

provided $1 < q \leq n$ and $p < q$. On the other hand, by (3.6) we have

$$\begin{aligned} \frac{\alpha}{2} \sum_i w^{ii} - \frac{2-\alpha}{\alpha C_1 h_1} &\geq \frac{1}{h_1} \left(\frac{\alpha(n-1)}{2} h_1 \left(h^{p-1} (h^2 + h_1^2)^{\frac{n-q}{2}} f \right)^{-\frac{1}{n-1}} - \frac{2-\alpha}{C_1 \alpha} \right) \\ &\geq \frac{1}{h_1} \left(\frac{\alpha(n-1)}{2} h_1 h^{-\frac{p-1}{n-1}} \left(1 + \frac{1}{C_1^2} \right)^{-\frac{n-q}{2(n-1)}} h_1^{-\frac{n-q}{n-1}} f^{-\frac{1}{n-1}} - \frac{2-\alpha}{C_1 \alpha} \right) \\ &\geq \frac{1}{h_1} \left(\frac{\alpha(n-1)}{2} C_1^{\frac{q-1}{n-1}} \left(1 + \frac{1}{C_1^2} \right)^{-\frac{n-q}{2(n-1)}} M^{\frac{q-p}{n-1}} f^{-\frac{1}{n-1}} - \frac{2-\alpha}{C_1 \alpha} \right) \\ &> 0. \end{aligned}$$

The above two inequalities imply (3.8).

Subcase(2_B): $C_2 h^{1-\frac{q-p}{n-1}} \leq h_1 \leq C_1 h$ where C_2 is chosen such that

$$\frac{\alpha(n-1)}{2} (1 + C_1^2)^{-\frac{n-q}{2(n-1)}} C_2 > \max \frac{|\nabla f|}{f^{1-\frac{1}{n-1}}}. \quad (3.10)$$

Furthermore, we now choose $M \gg C_1 \gg 1$ such that

$$\frac{\alpha(n-1)}{2} (1 + C_1^2)^{-\frac{n-q}{2(n-1)}} M^{\frac{q-p}{n-1}} (\max f)^{-\frac{1}{n-1}} > 2. \quad (3.11)$$

From (3.3), we get

$$\begin{aligned} 0 &\geq \frac{p-1}{h} + \frac{f_1}{fh_1} + \frac{1-\alpha}{h} - (2-\alpha) \frac{h}{\alpha h_1^2 + h^2} - \frac{\alpha(n-1)}{h} + \alpha \sum_i w^{ii} \\ &\geq \frac{p-n\alpha}{h} + \left(\frac{f_1}{fh_1} + \frac{\alpha}{2} \sum_i w^{ii} \right) + \left(\frac{\alpha}{2} \sum_i w^{ii} - \frac{2}{h} \right), \end{aligned}$$

which leads to a contradiction provided that

$$\left(\frac{f_1}{fh_1} + \frac{\alpha}{2} \sum_i w^{ii} \right) + \left(\frac{\alpha}{2} \sum_i w^{ii} - \frac{2}{h} \right) > 0. \quad (3.12)$$

Then combining (3.9), (3.10) and $C_2 h^{1-\frac{q-p}{n-1}} \leq h_1 \leq C_1 h$

$$\begin{aligned} \frac{f_1}{fh_1} + \frac{\alpha}{2} \sum_i w^{ii} &\geq -\frac{|\nabla f|}{fh_1} + \frac{\alpha(n-1)}{2} (h^{p-1} (h^2 + h_1^2)^{\frac{n-q}{2}} f)^{-\frac{1}{n-1}} \\ &\geq \frac{1}{h_1} \left(-\frac{|\nabla f|}{f} + \frac{\alpha(n-1)}{2} (1 + C_1^2)^{-\frac{n-q}{2(n-1)}} h_1 h^{-\frac{n-q+p-1}{n-1}} f^{-\frac{1}{n-1}} \right) \\ &\geq \frac{1}{h_1} \left(-\frac{|\nabla f|}{f} + \frac{\alpha(n-1)}{2} (1 + C_1^2)^{-\frac{n-q}{2(n-1)}} C_2 f^{-\frac{1}{n-1}} \right) > 0 \end{aligned}$$

provided $p < q$ and $q \leq n$. On the other hand, by (3.11) and $h_1 \leq C_1 h$, we have

$$\begin{aligned} \frac{\alpha}{2} \sum_i w^{ii} - \frac{2}{h} &\geq \frac{\alpha(n-1)}{2} \left(h^{p-1} (h^2 + h_1^2)^{\frac{n-q}{2}} f \right)^{-\frac{1}{n-1}} - \frac{2}{h} \\ &\geq \frac{1}{h} \left(\frac{\alpha(n-1)}{2} (1 + C_1^2)^{-\frac{n-q}{2(n-1)}} h^{1-\frac{n-q+p-1}{n-1}} f^{-\frac{1}{n-1}} - 2 \right) \\ &\geq \frac{1}{h} \left(\frac{\alpha(n-1)}{2} (1 + C_1^2)^{-\frac{n-q}{2(n-1)}} M^{\frac{q-p}{n-1}} f^{-\frac{1}{n-1}} - 2 \right) > 0 \end{aligned}$$

provided $p < q$ and $q \leq n$. The above inequalities directly deduce (3.12).

Subcase(2_C): $h_1 \leq C_2 h^{1-\frac{q-p}{n-1}}$, then

$$\frac{h_1}{h^\alpha} \leq C_2 \frac{h^{1-\alpha}}{h^{\frac{q-p}{n-1}}} \leq \frac{C_2}{M^{\frac{q-p}{n-1}}} h^{1-\alpha}. \quad (3.13)$$

So we get

$$\frac{|\nabla h|}{h^\alpha} \leq \frac{C_2}{M^{\frac{q-p}{n-1}}} h^{1-\alpha}, \quad \text{for any } x \in S^{n-1}. \quad (3.14)$$

Let Γ be a great circle connecting x_1 and x_2 with length less than or equal to π , here $h(x_1) = \min h$ and $h(x_2) = \max h$. Then

$$h(x_2)^{1-\alpha} - h(x_1)^{1-\alpha} = \int_{\Gamma} dh^{1-\alpha} \leq (1-\alpha) \int_{\Gamma} \frac{|\nabla h|}{h^\alpha} \leq \frac{C_2(1-\alpha)\pi}{M^{\frac{q-p}{n-1}}} h(x_2)^{1-\alpha}. \quad (3.15)$$

From equation (1.1), we have for $p < q$

$$h(x_1) = \min h \leq (\max f)^{\frac{1}{q-p}}.$$

This with (3.15) imply the upper bound of h provided that

$$\frac{C_2(1-\alpha)\pi}{M^{\frac{q-p}{n-1}}} \leq \frac{1}{2}. \quad (3.16)$$

So we get our desired upper bound for $\frac{h_1}{h^\alpha}$ from (3.14).

Finally, we would like to claim that our requirements for the constants C_1 , C_2 and M are not contradictory in Case 2, i.e., (3.6) to (3.11), and (3.10) to (3.16).

For (3.6) to (3.11), we first ask that, for any $C_1 > 0$, choose M large so that (3.11) holds. Then,

$$\begin{aligned} \text{left-hand side of (3.6)} &= \frac{\alpha(n-1)}{2} C_1^{\frac{q-1}{n-1}} \left(1 + \frac{1}{C_1^2}\right)^{-\frac{n-q}{2(n-1)}} M^{\frac{q-p}{n-1}} (\max f)^{-\frac{1}{n-1}} \\ &= \frac{\alpha(n-1)}{2} C_1 (1 + C_1^2)^{-\frac{n-q}{2(n-1)}} M^{\frac{q-p}{n-1}} (\max f)^{-\frac{1}{n-1}} \\ &= C_1 \cdot \text{left-hand side of (3.11)}. \end{aligned}$$

Thus, we only need to choose C_1 large enough, so that (3.6) is established.

For (3.10), (3.16), we ask that, for any $C_1 > 0$, choose C_2 large enough, so that (3.10) holds. By choosing M large again, (3.16) is true. We finish the proof. $\square$

Lemma 3.1 implies the upper bound of h .

Lemma 3.2. (*Upper bound of Theorem 1.2*) Assume that h is a solution to equation (1.1) for $p \in (0, 1]$ and $q \in (1, n]$, then there exists constant $C(f, n, p, q)$ such that

$$\sup_{S^{n-1}} h \leq C. \quad (3.17)$$

Proof. Let Γ be a great circle connecting x_1 and x_2 with length less than or equal to π , where $h(x_1) = \min h$ and $h(x_2) = \max h$. Then by (3.1) in Lemma 3.1,

$$h(x_2)^{1-\alpha} - h(x_1)^{1-\alpha} = \int_{\Gamma} dh^{1-\alpha} \leq (1-\alpha) \int_{\Gamma} \frac{|\nabla h|}{h^\alpha} \leq C.$$

On the other hand, from the equation (1.1), we have $h_{ij} \geq 0$ at x_1 and

$$h(x_1) = \min h \leq (\max f)^{\frac{1}{q-p}}.$$

Thus

$$h(x_2) \leq C.$$

□

We next use the following trick to get the lower bound of volume.

Lemma 3.3. *(Lower bound of volume in Theorem 1.2) Suppose h is a solution to the equation (1.1) when $p \in (0, 1]$ and $q \in (1, n]$, then there exists constant $C(f, n, p, q)$ such that*

$$V_n([h]) > C.$$

Proof. Let $K = [h]$. Then by Hölder's inequality,

$$V_n(K) = \frac{1}{n} \int_{S^{n-1}} \rho_K^n \geq \frac{1}{n} (\omega_{n-1})^{\frac{q-n}{q}} \left(\int_{S^{n-1}} \rho_K^q \right)^{\frac{n}{q}} \quad (3.18)$$

provided $0 < q \leq n$. Since h is a solution to equation (1.1), then

$$\begin{aligned} \int_{\alpha_K^{-1}(\eta)} \rho^q(\xi) d\xi &= \int_{\eta} \rho^q(\alpha_K^{-1}(x)) \text{Jac}(\alpha_K^{-1}) dx \\ &= \int_{\eta} \rho^{q-n}(\alpha_K^{-1}(x)) h(x) \det(h_{ij} + h\delta_{ij}) dx \\ &= \int_{\eta} h^p(x) f(x) dx \end{aligned} \quad (3.19)$$

for any $\eta \subset S^{n-1}$. By the definition of support function,

$$h(x) \geq h_{\max} \langle x, u_0 \rangle_+$$

with $h(u_0) = h_{\max}$. Moreover, we have $h_{ij} \leq 0$ at u_0 . From the equation (1.1), we have

$$h(u_0)^{n-1} \geq f(u_0) h(u_0)^{p-1+n-q},$$

which means

$$h_{\max} \geq C \quad (3.20)$$

when $p < q$. From (3.19), we have

$$\begin{aligned}
\int_{S^{n-1}} \rho^q(\xi) d\xi &= \int_{S^{n-1}} h^p(x) f(x) dx \\
&\geq (\min_{S^{n-1}} f) \int_{S^{n-1}} h^p(x) dx \\
&\geq (\min_{S^{n-1}} f) h_{\max}^p \int_{S^{n-1}} \langle x, u_0 \rangle_+^p dx \\
&\geq C,
\end{aligned} \tag{3.21}$$

where the last inequality is because (3.20), and the integral $\int_{S^{n-1}} \langle x, u_0 \rangle_+^p dx$ is positive constant. Combining (3.18) and (3.21), the volume of K is bounded from below. $\square$

4. UNIQUENESS NEAR THE BALL

After obtaining C^0 estimates of the form given in Section 3, we apply the method in Böröczky-Chen-Liu-Saroglou [4] to derive the uniqueness result Theorem 1.3. For the sake of completeness, we repeat some lemmas and steps from [4].

Lemma 4.1. (BCLS [4]) *Let $\lambda > 1$, $p \in (-1, 1)$, $q_m > 1$ tend to n , and $K_m \in \mathcal{K}_o^n$ tend to $K_\infty \in \mathcal{K}_o^n$ as m tends to infinity where $\lambda^{-1} \mathcal{H}^{n-1} \leq \tilde{C}_{p,q_m,K_m} \leq \lambda \mathcal{H}^{n-1}$. Then $\tilde{C}_{p,q_m,K_m}$ tends weakly to S_{p,K_∞} .*

Now we prove Theorem 1.3.

Proof of Theorem 1.3. We begin first with the case $p \in [1, q)$ and $q \in (1, n]$. Consider a family of operators $\Psi : C_+^{2,\alpha}(S^{n-1}) \rightarrow C^\alpha(S^{n-1})$ defined by

$$\Psi(h) := h^{1-p} (h^2 + |\nabla h|^2)^{\frac{q-n}{2}} \det(\nabla^2 h + hI). \tag{4.1}$$

It is easy to calculate that the linear operator of Ψ at $h_0 \equiv 1$, denoted by $L_{h_0} : C^{2,\alpha}(S^{n-1}) \rightarrow C^\alpha(S^{n-1})$ is

$$L_{h_0}(\psi) = \Delta_{S^{n-1}} \psi + (q - p)\psi.$$

The spherical Laplacian $\Delta_{S^{n-1}}$ admits discrete eigenvalues

$$-\lambda_k = k^2 + (n - 2)k, \quad k = 0, 1, 2, \dots.$$

It is easy to find that L_{h_0} is invertible, provided $p \in [1, q)$ and $q \in (1, n]$. Indeed, the eigenvalues $\tilde{\lambda}_k$ of L_{h_0} are

$$\tilde{\lambda}_k = -k^2 - (n - 2)k + q - p, \quad k = 0, 1, 2, \dots,$$

and $\tilde{\lambda}_0 = q - p > 0$, $\tilde{\lambda}_1 = 1 - n + q - p < 0$ and $\tilde{\lambda}_k < 0$ when $k = 2, 3, \dots$. By inverse function theorem, there exists a neighborhood $\mathcal{N}$ of h_0 such that, if $h_1, h_2 \in \mathcal{N}$ and

$$\Psi(h_1) = \Psi(h_2),$$

then $h_1 = h_2$. Due to this, to prove Theorem 1.3 it is sufficient to prove that, there exists constant $\epsilon > 0$ depending only on n, α, p and q such that if the smooth function f satisfies $\|f - 1\|_{C^\alpha(S^{n-1})} < \epsilon$, then for any convex body K satisfying

$$d\tilde{C}_{p,q}(K, \cdot) = f d\mathcal{H}^{n-1},$$

then the support function $h_K \in \mathcal{N}$. Suppose not, then there exist a sequence function $f_i \rightarrow 1$ in $C^\alpha(S^{n-1})$ and a sequence of convex bodies K_i such that

$$d\tilde{C}_{p,q}(K_i, \cdot) = f_i d\mathcal{H}^{n-1},$$

but $h_{K_m} \notin \mathcal{N}$ when m is sufficiently large. By Theorem 1.2, we have $\|h_{K_m}\|_{L^\infty(S^{n-1})} \leq C$ and $|K_m| \geq C^{-1}$ for some positive constant C . Thus, using Blaschke selection theorem, K_m converges to a convex body K_∞ up to a subsequence. By the weak convergence of $\tilde{C}_{p,q}(K, \cdot)$, then

$$d\tilde{C}_{p,q}(K_\infty) = \mathcal{H}^{n-1}.$$

By using the uniqueness result in [18] when $p > 1$ and $q \leq n$, K_∞ must be the ball centered at origin. That is to say,

$$\|h_K - 1\|_{L^\infty(S^{n-1})} \rightarrow 0 \quad \text{as } m \rightarrow \infty.$$

By Caffarelli's $C^{2,\alpha}$ results [14, 15], $\|h_{K_m}\|_{C^{2,\alpha}(S^{n-1})} \leq C_0$ for some positive constant C_0 . Thus,

$$\|h_{K_m}^{p-1}(h_{K_m}^2 + |\nabla h_{K_m}|^2)^{\frac{n-q}{2}} f_m - 1\|_{C^\alpha(S^{n-1})} < \epsilon_0 \rightarrow 0 \text{ as } \epsilon \rightarrow 0.$$

Let $\omega = h_{K_m} - 1$, $a_{ij} = \int_0^1 U_t^{ij} dt$, where U_t^{ij} is the cofactor matrix of

$$\nabla^2(1 + t\omega) + (1 + t\omega)I.$$

Then

$$\begin{aligned} h_{K_m}^{p-1}(h_{K_m}^2 + |\nabla h_{K_m}|^2)^{\frac{n-q}{2}} f_m - 1 &= \det(\nabla^2 h_{K_m} + h_{K_m} I) - 1 \\ &= \sum_{i,j=1}^n a_{ij}(\nabla_{ij}^2 \omega + \omega \delta_{ij}). \end{aligned}$$

Note that $C^{-1}I \leq a_{ij} \leq CI$ for some $C > 0$, thus ω satisfies a uniformly elliptic linear equation. By Schauder estimates,

$$\|h_{K_m} - 1\|_{C^{2,\alpha}} \leq C \left(\|h_{K_m} - 1\|_{L^\infty} + \|h_{K_m}^{p-1}(h_{K_m}^2 + |\nabla h_{K_m}|^2)^{\frac{n-q}{2}} f_m - 1\|_{C^\alpha} \right) \rightarrow 0$$

as $m \rightarrow \infty$. This is a contradiction with $h_{K_m} \notin \mathcal{N}$.

As for the case when $n - \epsilon < q \leq n$ and $p \in (0, 1)$, we need to use the Lemma 4.1 and the uniqueness result in Brendle-Choi-Daskalopoulos [12]. One can also see these details in [4]. We complete the proof. $\square$

5. THE L_p DUAL BRUNN-MINKOWSKI INEQUALITY

5.1. Case $p \geq 1$. Recently, Sadovskiy-Zhang [48] proved that Lutwak's conjecture holds in the symmetric case. In this subsection, we use the classical trick to prove the L_p Brunn-Minkowski inequality when $p \geq 1$ which is stated in Theorem 1.4.

We first states Sadovskiy-Zhang's result in [48].

Theorem 5.1 (Sadovskiy-Zhang [48]). *Let K, L be origin-symmetric convex bodies in $\mathbb{R}^n$. Then for $0 < q \leq n$,*

$$\tilde{V}_q((1-\lambda)K + \lambda L)^{\frac{1}{q}} \geq (1-\lambda)\tilde{V}_q(K)^{\frac{1}{q}} + \lambda\tilde{V}_q(L)^{\frac{1}{q}} \quad \forall \lambda \in [0, 1]. \quad (5.1)$$

with equality if K and L are dilates of each other.

We are now in a position to prove the first item in Theorem 1.4 by a classical trick.

Proof of the first item in Theorem 1.4. Let $\bar{K} = \tilde{V}_q(K)^{-\frac{1}{q}}K$, $\bar{L} = \tilde{V}_q(L)^{-\frac{1}{q}}L$, and $\lambda = \frac{\tilde{V}_q(L)^{\frac{p}{q}}}{\tilde{V}_q(K)^{\frac{p}{q}} + \tilde{V}_q(L)^{\frac{p}{q}}}$. By Jensen's inequality, for $p \geq 1, \lambda \in [0, 1]$,

$$h_{(1-\lambda)K +_p \lambda L} = [(1-\lambda)h_K^p + \lambda h_L^p]^{\frac{1}{p}} \geq (1-\lambda)h_K + \lambda h_L = h_{(1-\lambda)K + \lambda L},$$

thus

$$(1-\lambda)K + \lambda L \subseteq (1-\lambda)K +_p \lambda L. \quad (5.2)$$

Combining with (5.1) and (5.2),

$$\begin{aligned} \frac{\tilde{V}_q(K +_p L)^{\frac{1}{q}}}{(\tilde{V}_q(K)^{\frac{p}{q}} + \tilde{V}_q(L)^{\frac{p}{q}})^{\frac{1}{p}}} &= \tilde{V}_q \left(\frac{K +_p L}{(\tilde{V}_q(K)^{\frac{p}{q}} + \tilde{V}_q(L)^{\frac{p}{q}})^{\frac{1}{p}}} \right)^{\frac{1}{q}} \\ &= \tilde{V}_q((1-\lambda)\bar{K} +_p \lambda\bar{L})^{\frac{1}{q}} \\ &\geq \tilde{V}_q((1-\lambda)\bar{K} + \lambda\bar{L})^{\frac{1}{q}} \\ &\geq (1-\lambda)\tilde{V}_q(\bar{K})^{\frac{1}{q}} + \lambda\tilde{V}_q(\bar{L})^{\frac{1}{q}} = 1. \end{aligned}$$

Hence for $p \geq 1, q \in (0, n]$,

$$\tilde{V}_q(K +_p L)^{\frac{p}{q}} \geq \tilde{V}_q(K)^{\frac{p}{q}} + \tilde{V}_q(L)^{\frac{p}{q}}.$$

By the homogeneity of the above inequality, we immediately get

$$\tilde{V}_q((1-\lambda)K +_p \lambda L)^{\frac{p}{q}} \geq (1-\lambda)\tilde{V}_q(K)^{\frac{p}{q}} + \lambda\tilde{V}_q(L)^{\frac{p}{q}}.$$

□

5.2. Case $p \in (0, 1)$. In this subsection, our goal is to prove the near spherical L_p dual Brunn-Minkowski inequality when $p \in (0, 1)$ which is stated in Theorem 1.4. Inspired by the method in [8, 20], we first obtain the L_p dual Minkowski inequality, which is Theorem (5.2). Then in Theorem 5.4 we prove the equivalence between the L_p Brunn-Minkowski inequality (1.4) and the L_p Minkowski inequality (5.3) for dual quermassintegrals. Finally, we prove Theorem 1.4.

We first state our result on the L_p Minkowski inequality for dual quermassintegrals.

Theorem 5.2. (*L_p dual Minkowski inequality*) For $n \geq 2$, $\alpha \in (0, 1)$, $p \in (0, 1)$ and $q \in (p, n]$. There exists a constant $\epsilon_0 \in (0, 1)$ that depends only on n, α, p and q such that, if $K, L \in \mathcal{K}_{+,e}^{2,\alpha}$ with $\|h_K - 1\|_{C^{2,\alpha}} \leq \epsilon_0$, then

$$\left(\int_{S^{n-1}} \left(\frac{h_L}{h_K} \right)^p d\bar{C}_{q,K} \right)^{\frac{1}{p}} \geq \left(\frac{\tilde{V}_q(L)}{\tilde{V}_q(K)} \right)^{\frac{1}{q}}, \quad (5.3)$$

where $\bar{C}_{q,K} := \frac{\tilde{C}_{q,K}}{\tilde{V}_q(K)}$.

To prove the Theorem 5.2, we first consider a minimization problem.

Lemma 5.3. *The minimization problem*

$$\min_{\phi \in C_{+,e}^{2,\alpha}(S^{n-1})} \Phi(\phi)$$

has a minimizer h_{L_0} , which is the support function of some $L_0 \in \mathcal{K}_{+,e}^{2,\alpha}$. The functional $\Phi : C_{+,e}^{2,\alpha}(S^{n-1}) \rightarrow \mathbb{R}$ is defined by

$$\Phi(\phi) = \frac{1}{\tilde{V}_q([\phi])^{\frac{p}{q}}} \int_{S^{n-1}} \left(\frac{\phi}{h_K} \right)^p d\tilde{C}_{q,K}$$

for the given $K \in \mathcal{K}_{+,e}^{2,\alpha}$.

Proof. Let $h_{[\phi]}$ be the support function of convex body $[\phi]$. Since $\phi \in C_{+,e}^{2,\alpha}(S^{n-1})$, $[\phi] \in \mathcal{K}_{+,e}^{2,\alpha}$ and $h_{[\phi]} \in C_{+,e}^{2,\alpha}(S^{n-1})$. Moreover,

$$h_{[\phi]} \leq \phi, \tilde{V}_q([\phi]) = \tilde{V}_q(h_{[\phi]}) \Rightarrow \Phi(h_{[\phi]}) \leq \Phi(\phi)$$

which means that the minimizer of $\Phi(h_{[\phi]})$ is actually the minimiser of $\Phi(\phi)$. Note that $\tilde{V}_q$ is homogeneous of degree q which yields

$$\Phi(t\phi) = \frac{1}{\tilde{V}_q([t\phi])^{\frac{p}{q}}} \int_{S^{n-1}} \left(\frac{t\phi}{h_K} \right)^p d\tilde{C}_{q,K} = \Phi(\phi).$$

This fact allows us to add a limitation of $\tilde{V}_q(L) = 1$. Therefore,

$$\begin{aligned} s &:= \min_{\phi \in C_e^+(\mathbb{S}^{n-1})} \Phi(\phi) \\ &= \min_{h_L \in C_e^+(\mathbb{S}^{n-1})} \Phi(h_L) \\ &= \min \left\{ \int_{\mathbb{S}^{n-1}} \left(\frac{h_L}{h_K} \right)^p d\tilde{C}_{q,K} : L \in \mathcal{K}_e^n, \tilde{V}_q(L) = 1 \right\}. \end{aligned}$$

Denote $\int_{\mathbb{S}^{n-1}} \left(\frac{h_L}{h_K} \right)^p d\tilde{C}_{q,K}$ as $F(L)$. Assume L_j be a minimizing sequence such that $F(L_j) \rightarrow s$ as $j \rightarrow \infty$. Next, we need to prove that there exists a uniform constant $C > 0$, such that for any j , $h_{L_j} \leq C$.

We suppose to the contrary that, passing to a subsequence, $R_j := \|h_{L_j}(e_j)\|_{L^\infty} \rightarrow \infty$ as $j \rightarrow \infty$. Since $K \in \mathcal{K}_e^n$, we have $h_K^{-p} d\tilde{C}_{q,K} = f d\mathcal{H}^{n-1}$ for some positive continuous function f . Therefore,

$$\begin{aligned} F(L_j) &= \int_{\mathbb{S}^{n-1}} \left(\frac{h_{L_j}}{h_K} \right)^p d\tilde{C}_{q,K} = \int_{\mathbb{S}^{n-1}} h_{L_j}^p f d\mathcal{H}^{n-1} \\ &\geq R_j^p \int_{\mathbb{S}^{n-1}} (x, e_j)_+^p f d\mathcal{H}^{n-1} \geq C R_j^p \rightarrow \infty \end{aligned}$$

as $j \rightarrow \infty$, which contradicts the fact that $F(L_j) \rightarrow s$ as $j \rightarrow \infty$.

By the Blaschke selection theorem, there exists a convex set $L_0 \in \mathcal{K}^n$ and a subsequence L_{j_k} such that $L_{j_k} \rightarrow L_0$ in Hausdorff distance. We next show that $L_0 \in \mathcal{K}_e^n$. Since $\tilde{V}_q(L_j) = 1$, then the Hölder's inequality yields

$$\begin{aligned} 1 = \tilde{V}_q(L_j) &= \frac{1}{n} \int_{\mathbb{S}^{n-1}} \rho_{L_j}^q(u) du \\ &\leq \frac{1}{n} \left(\int_{\mathbb{S}^{n-1}} \rho_{L_j}^q(u) du \right)^{\frac{q}{n}} \left(\int_{\mathbb{S}^{n-1}} 1 du \right)^{1-\frac{q}{n}} \\ &= n^{\frac{q}{n}-1} (\omega_{n-1})^{1-\frac{q}{n}} V_n(L_j)^{\frac{q}{n}}. \end{aligned} \tag{5.4}$$

Hence, there exists a constant $C > 0$ such that $V_n(L_j) \geq C$ for all j . Let v_j be a direction where h_{L_j} attains its minimum. The upper bound of h_{L_j} shows that there exists $R > 0$ such that $L_j \subseteq B_R$. Consider the symmetric strip

$$S_{L_j} := \{x \in \mathbb{R}^n : |\langle x, v_j \rangle| \leq h_{L_j}(v_j)\}.$$

It is obvious that $L_j \subseteq S_{L_j} \cap B_R$. We could deduce

$$V_n(L_j) \leq V_n(S_{L_j} \cap B_R) = C_0 R^{n-1} \min h_{L_j}$$

for some positive constant C_0 . Combining this with $V_n(L_j) \geq C$, we obtain a uniform positive lower bound for $\min h_{L_j}$,

$$\min h_{L_j} \geq \frac{C}{C_0 R^{n-1}} > 0.$$

Therefore, $L_0 \in \mathcal{K}_e^n$. Since $h_{L_{j_k}} \rightarrow h_{L_0}$ uniformly, we have $F(L_{j_k}) \rightarrow F(L_0)$. Hence $F(L_0) = s$ and L_0 is the desired minimiser. $\square$

This lemma puts us in a position to prove Theorem 5.2

Proof of Theorem 5.2. By Lemma 5.3, L_0 is the minimiser. For any $f \in C_e^+(\mathbb{S}^{n-1})$, define

$$q_t := \log h_{L_0}(x) + tf(x) + o(t, x), \text{ for } t \in (-\epsilon, \epsilon),$$

then $q_t \in C_e^+(\mathbb{S}^{n-1})$ for all $t \in (-\epsilon, \epsilon)$. Let L_t be the Wulff shape associated with q_t , one has $L_t \in \mathcal{K}_e^n$ and L_t is actually the minimiser L_0 when $t = 0$.

Since h_{L_0} is the minimiser of $\Phi(\phi)$, then by the variational formula in [32], we have

$$\begin{aligned} 0 &= \frac{d}{dt} \Big|_{t=0} \log \Phi(q_t) \\ &= \frac{d}{dt} \Big|_{t=0} \log \left[\frac{1}{\tilde{V}_q([q_t])^{\frac{p}{q}}} \int_{\mathbb{S}^{n-1}} \left(\frac{h_{L_t}}{h_K} \right)^p d\tilde{C}_{q,K} \right] \\ &= \frac{d}{dt} \Big|_{t=0} \left[-\frac{p}{q} \log \tilde{V}_q([q_t]) + \log \left(\int_{\mathbb{S}^{n-1}} \left(\frac{h_{L_t}}{h_K} \right)^p d\tilde{C}_{q,K} \right) \right] \\ &= \frac{p \int_{\mathbb{S}^{n-1}} \left(\frac{h_{L_0}}{h_K} \right)^{p-1} \frac{h_{L_0}}{h_K} f d\tilde{C}_{q,K}}{\int_{\mathbb{S}^{n-1}} \left(\frac{h_{L_0}}{h_K} \right)^p d\tilde{C}_{q,K}} - \frac{p \int_{\mathbb{S}^{n-1}} f d\tilde{C}_{q,L_0}}{\tilde{V}_q(L_0)}. \end{aligned}$$

From the arbitrariness of f , it follows that

$$h_K^{-p} d\tilde{C}_{q,K} = \frac{\int_{\mathbb{S}^{n-1}} \left(\frac{h_{L_0}}{h_K} \right)^p d\tilde{C}_{q,K}}{\tilde{V}_q(L_0)} h_{L_0}^{-p} d\tilde{C}_{q,L_0} = h_{kL_0}^{-p} d\tilde{C}_{q,kL_0},$$

where the positive rescaling constant

$$k = \left(\frac{\tilde{V}_q(L_0)}{\int_{\mathbb{S}^{n-1}} \left(\frac{h_{L_0}}{h_K} \right)^p d\tilde{C}_{q,K}} \right)^{\frac{1}{p-q}}.$$

Note that $K \in \mathcal{K}_{+,e}^{2,\alpha}$. By Caffarelli's regularity results [15, Theorem 4], we have $kL_0 \in \mathcal{K}_{+,e}^{2,\alpha}$. Since $\|h_K - 1\|_{C^{2,\alpha}} \leq \epsilon$, it means that

$$\left\| \frac{h_K^{-p} d\tilde{C}_{q,K}}{d\mathcal{H}^{n-1}} - 1 \right\|_{C^\alpha(S^{n-1})} < \epsilon_0$$

for some $\epsilon_0 > 0$. By the uniqueness result of L_p dual Minkowski problem in [4, Theorem 1.5], it follows that $L_0 = k^{-1}K \in \mathcal{K}_{+,e}^{2,\alpha}$. Therefore, for any $L \in \mathcal{K}_e^n$, we have $h_L \in C_e^+(S^{n-1})$. Since $L_0 = k^{-1}K$ is a minimizer of Φ and Φ is homogeneous of degree zero, we obtain

$$\Phi(K) = \Phi(k^{-1}K) \leq \Phi(h_L),$$

and finally,

$$\left(\int_{S^{n-1}} \left(\frac{h_L}{h_K} \right)^p d\tilde{V}_{q,K} \right)^{\frac{1}{p}} \geq \left(\frac{\tilde{V}_q(L)}{\tilde{V}_q(K)} \right)^{\frac{1}{q}},$$

which is desired. $\square$

We finally prove the equivalence between the L_p Minkowski inequality and the L_p Brunn–Minkowski inequality for the dual quermassintegral when $p, q > 0$, and the method is from [8]. Notably, this equivalence holds without symmetry or smoothness assumptions.

Lemma 5.4. *Suppose $p, q > 0$. For $K, L \in \mathcal{K}_o^n$, the L_p Minkowski inequality (5.3) and the L_p Brunn–Minkowski inequality (1.4) for the dual quermassintegral are equivalent.*

Proof. Suppose $K, L \in \mathcal{K}_o^n$. For $0 \leq \lambda \leq 1$, let

$$Q_\lambda = (1 - \lambda)K +_p \lambda L,$$

i.e., Q_λ is the Wulff shape associated with the function $q_\lambda = ((1 - \lambda)h_K^p + \lambda h_L^p)^{\frac{1}{p}}$. It will be convenient to consider h_{Q_λ} as being defined for λ in the open interval $(-\varepsilon_0, 1 + \varepsilon_0)$, where $\varepsilon_0 > 0$ is chosen so that for $\lambda \in (-\varepsilon_0, 1 + \varepsilon_0)$, the function q_λ is strictly positive.

Step 1: (5.3) $\Rightarrow$ (1.4).

Assume that the L_p Minkowski inequality (5.3) holds. Then

$$\begin{aligned}
\tilde{V}_q(Q_\lambda) &= \int_{\mathbb{S}^{n-1}} 1 d\tilde{C}_{q,Q_\lambda} \\
&= \int_{\mathbb{S}^{n-1}} h_{Q_\lambda}^p h_{Q_\lambda}^{-p} d\tilde{C}_{q,Q_\lambda} \\
&= \int_{\mathbb{S}^{n-1}} ((1-\lambda)h_K^p + \lambda h_L^p) h_{Q_\lambda}^{-p} d\tilde{C}_{q,Q_\lambda} \\
&= (1-\lambda) \int_{\mathbb{S}^{n-1}} \left(\frac{h_K}{h_{Q_\lambda}} \right)^p d\tilde{C}_{q,Q_\lambda} + \lambda \int_{\mathbb{S}^{n-1}} \left(\frac{h_L}{h_{Q_\lambda}} \right)^p d\tilde{C}_{q,Q_\lambda} \\
&\geq (1-\lambda) \tilde{V}_q(Q_\lambda) \left(\frac{\tilde{V}_q(K)}{\tilde{V}_q(Q_\lambda)} \right)^{\frac{p}{q}} + \lambda \tilde{V}_q(Q_\lambda) \left(\frac{\tilde{V}_q(L)}{\tilde{V}_q(Q_\lambda)} \right)^{\frac{p}{q}} \\
&= \tilde{V}_q(Q_\lambda)^{1-\frac{p}{q}} [(1-\lambda) \tilde{V}_q(K)^{\frac{p}{q}} + \lambda \tilde{V}_q(L)^{\frac{p}{q}}].
\end{aligned}$$

Hence

$$\tilde{V}_q(Q_\lambda) \geq [(1-\lambda) \tilde{V}_q(K)^{\frac{p}{q}} + \lambda \tilde{V}_q(L)^{\frac{p}{q}}]^{\frac{q}{p}}, \quad (5.5)$$

which is actually (1.4).

Step 2: (1.4) $\Rightarrow$ (5.3).

Define the function $f : [0, 1] \rightarrow (0, \infty)$, given by $f(\lambda) = \tilde{V}_q(Q_\lambda)^{\frac{p}{q}}$. (5.5) shows that $f(\lambda) \geq (1-\lambda)f(0) + \lambda f(1)$, which is not enough, because we want to verify that f is a concave function, i.e., $f((1-\lambda)\sigma + \lambda\tau) \geq (1-\lambda)f(\sigma) + \lambda f(\tau)$ for $\sigma, \tau \in [0, 1]$. When $p \geq 1$, $f(\lambda)$ is concave. However, when $p < 1$, it is not obvious, so a property of Wulff shapes is needed. For given $\sigma, \tau \in [0, 1]$, let

$$K_\sigma = (1-\sigma) \cdot K +_p \sigma \cdot L, \quad K_\tau = (1-\tau) \cdot K +_p \tau \cdot L.$$

Since K_σ is the Wulff shape of the function $((1-\sigma)h_K^p + \sigma h_L^p)^{\frac{1}{p}}$, we have

$$h_{K_\sigma} \leq ((1-\sigma)h_K^p + \sigma h_L^p)^{\frac{1}{p}}.$$

If $\lambda \in [0, 1]$ and $\alpha = (1-\lambda)\sigma + \lambda\tau$, this gives

$$\begin{aligned}
(1-\lambda)h_{K_\sigma}^p + \lambda h_{K_\tau}^p &\leq (1-\lambda)[(1-\sigma)h_K^p + \sigma h_L^p] + \lambda[(1-\tau)h_K^p + \tau h_L^p] \\
&= [(1-\lambda)(1-\sigma) + \lambda(1-\tau)]h_K^p + [(1-\lambda)\sigma + \lambda\tau]h_L^p \\
&= (1-\alpha)h_K^p + \alpha h_L^p.
\end{aligned}$$

Thus, $[(1-\lambda)h_{K_\sigma}^p + \lambda h_{K_\tau}^p]^{\frac{1}{p}} \leq [(1-\alpha)h_K^p + \alpha h_L^p]^{\frac{1}{p}}$, and taking the Wulff shapes of these functions allows us to conclude that

$$(1-\lambda)K_\sigma +_p \lambda K_\tau \subseteq (1-\alpha)K +_p \alpha L.$$

This gives

$$\begin{aligned}
f((1-\lambda)\sigma + \lambda\tau) &= \tilde{V}_q((1-\alpha) \cdot K +_p \alpha \cdot L)^{\frac{p}{q}} \\
&\geq \tilde{V}_q((1-\lambda) \cdot K_\sigma +_p \lambda \cdot K_\tau)^{\frac{p}{q}} \\
&\geq (1-\lambda)\tilde{V}_q(K_\sigma)^{\frac{p}{q}} + \lambda\tilde{V}_q(K_\tau)^{\frac{p}{q}} \\
&= (1-\lambda)f(\sigma) + \lambda f(\tau),
\end{aligned}$$

which is the desired concavity of f , i.e., $f(1) - f(0) \leq f'(0)$.

From $f(\lambda) = \tilde{V}_q(Q_\lambda)^{\frac{p}{q}}$, we have

$$f'(\lambda)|_{\lambda=0} = \frac{p}{q} \tilde{V}_q(Q_\lambda)^{\frac{p}{q}-1} \tilde{V}_q(Q_\lambda)'|_{\lambda=0}.$$

Since Q_λ is the Wulff shape of the function $q_\lambda^p = ((1-\lambda)h_K^p + \lambda h_L^p)$, and

$$q_\lambda^p = h_K^p + \lambda(h_L^p - h_K^p) = h_K^p \left[1 + \lambda \left(\frac{h_L^p - h_K^p}{h_K^p} \right) \right],$$

it follows immediately that, for $\lambda \in [0, 1]$,

$$\log q_\lambda = \log h_K + \frac{1}{p} \log \left[1 + \lambda \left(\frac{h_L^p - h_K^p}{h_K^p} \right) \right] = \log h_K + \frac{\lambda}{p} \left[\left(\frac{h_L}{h_K} \right)^p - 1 \right] + o(\lambda).$$

Therefore, combining the perturbation $g = \frac{1}{p} \left[\left(\frac{h_L}{h_K} \right)^p - 1 \right]$ with the variational formulation in [32], we have

$$\begin{aligned}
\frac{d}{d\lambda} \tilde{V}_q(Q_\lambda)|_{\lambda=0} &= q \int_{\mathbb{S}^{n-1}} g(u) d\tilde{C}_{q,K}(u) \\
&= \frac{q}{p} \int_{\mathbb{S}^{n-1}} \left[\left(\frac{h_L}{h_K} \right)^p - 1 \right] d\tilde{C}_{q,K}(u) \\
&= \frac{q}{p} \int_{\mathbb{S}^{n-1}} \left(\frac{h_L}{h_K} \right)^p d\tilde{C}_{q,K}(u) - \frac{q}{p} \tilde{V}_q(K).
\end{aligned}$$

Therefore, the concavity of f yields

$$\begin{aligned}
\tilde{V}_q(K)^{\frac{p-q}{q}} \left[\int_{\mathbb{S}^{n-1}} \left(\frac{h_L}{h_K} \right)^p d\tilde{C}_q(K, u) - \tilde{V}_q(K) \right] &= f'(0) \\
&\geq f(1) - f(0) = \tilde{V}_q(L)^{\frac{p}{q}} - \tilde{V}_q(K)^{\frac{p}{q}},
\end{aligned}$$

which gives the L_p dual Minkowski inequality. $\square$

Proof of the second item in Theorem 1.4. Combining Theorem 5.2 and Lemma 5.4, we complete the proof. $\square$

ACKNOWLEDGEMENT

The authors would like to express sincere gratitude to professor Yong Huang for his constant guidance.

REFERENCES

- [1] A. D. Aleksandrov, On the theory of mixed volumes. III. Extension of two theorems of Minkowski on convex polyhedra to arbitrary convex bodies, *Mat. Sb. N. S.* **3** (1938), 27–46.
- [2] A. D. Aleksandrov, Existence and uniqueness of a convex surface with a given integral curvature, *C. R. (Doklady) Acad. Sci. URSS (N.S.)* **35** (1942), 131–134. MR0007625
- [3] C. Bianchini, A. Colesanti, D. Pagnini and A. Roncoroni, On p -Brunn-Minkowski inequalities for intrinsic volumes, with $0 \leq p < 1$, *Math. Ann.* **387** (2023), no. 1-2, 321–352.
- [4] K. J. Böröczky, S. Chen, W. Liu and C. Saroglou, Uniqueness in the near isotropic L_p dual Minkowski problem, *arXiv:2505.01066*, 2025.
- [5] K. J. Böröczky, S. Chen, W. Liu and C. Saroglou, Compactness of the L_p dual Minkowski problem in $\mathbb{R}^3$, *arXiv:2505.17219*, 2025.
- [6] K. J. Böröczky and F. Fodor, The L_p dual Minkowski problem for $p > 1$ and $q > 0$, *J. Differential Equations* **266** (2019), no. 12, 7980–8033. MR3944247
- [7] K. J. Böröczky, A. Kovács, S. Mui and G. Zhang, Dual Curvature Density Equation with Group Symmetry, *J. Differential Equations*, **465** (2026), Paper No. 114197, 23 pp.
- [8] K. J. Böröczky, E. Lutwak, D. Yang and G. Zhang, The log-Brunn-Minkowski inequality, *Adv. Math.* **231** (2012), no. 3-4, 1974–1997. MR2964630
- [9] K. J. Böröczky, E. Lutwak, D. Yang and G. Zhang, The logarithmic Minkowski problem, *J. Amer. Math. Soc.* **26** (2013), no. 3, 831–852. MR3037788
- [10] K. J. Böröczky, E. Lutwak, D. Yang, G. Zhang and Y. Zhao, The dual Minkowski problem for symmetric convex bodies, *Adv. Math.* **356** (2019), 106805, 30 pp. MR4008522
- [11] K. J. Böröczky and C. Saroglou, Uniqueness when the L_p curvature is close to be a constant for $p \in [0, 1)$, *Calc. Var. Partial Differential Equations* **63** (2024), no. 6, Paper No. 154, 26 pp. MR4765819
- [12] S. Brendle, K. Choi and P. Daskalopoulos, Asymptotic behavior of flows by powers of the Gaussian curvature, *Acta Math.* **219** (2017), no. 1, 1–16. MR3765656
- [13] C. Cabezas-Moreno and J. Hu, The L_p dual Christoffel-Minkowski problem for $1 < p < q \leq k + 1$ with $1 \leq k \leq n$, *Calc. Var. Partial Differential Equations* **64** (2025), no. 7, Paper No. 229, 29 pp. MR4949561
- [14] L. Caffarelli, A localization property of viscosity solutions to the Monge-Ampère equation and their strict convexity, *Ann. of Math. (2)* **131** (1990), no. 1, 129–134. MR1038359
- [15] L. Caffarelli, Interior $W^{2,p}$ estimates for solutions of the Monge-Ampère equation, *Ann. of Math. (2)* **131** (1990), no. 1, 135–150. MR1038360
- [16] C. Chen, Y. Huang and Y. Zhao, Smooth solutions to the L_p dual Minkowski problem, *Math. Ann.* **373** (2019), no. 3-4, 953–976. MR3953117
- [17] H. Chen, S. Chen and Q. Li, Variations of a class of Monge-Ampère-type functionals and their applications, *Anal. PDE* **14** (2021), no. 3, 689–716. MR4259871
- [18] H. Chen and Q. Li, The L_p dual Minkowski problem and related parabolic flows, *J. Funct. Anal.* **281** (2021), no. 8, Paper No. 109139, 65 pp. MR4271790
- [19] S. Chen, Y. Feng and W. Liu, Uniqueness of solutions to the logarithmic Minkowski problem in $\mathbb{R}^3$, *Adv. Math.* **411** (2022), part A, Paper No. 108782, 18 pp. MR4512403

- [20] S. Chen, Y. Huang, Q. Li, J. Liu, The L_p -Brunn-Minkowski inequality for $p < 1$, *Adv. Math.* **368** (2020), 107166, 21 pp. MR4088419
- [21] K. S. Chou and X. J. Wang, The L_p -Minkowski problem and the Minkowski problem in centroaffine geometry, *Adv. Math.* **205** (2006), no. 1, 33–83. MR2254308
- [22] A. Colesanti, G. V. Livshyts and A. Marsiglietti, On the stability of Brunn-Minkowski type inequalities, *J. Funct. Anal.* **273** (2017), no. 3, 1120–1139. MR3653949
- [23] D. Cordero-Erausquin and L. Rotem, Improved log-concavity for rotationally invariant measures of symmetric convex sets, *Ann. Probab.* **51** (2023), no. 3, 987–1003. MR4583060
- [24] W. J. Firey, p -means of convex bodies, *Math. Scand.* **10** (1962), 17–24. MR0141003
- [25] R. J. Gardner, The Brunn-Minkowski inequality, *Bull. Amer. Math. Soc. (N.S.)* **39** (2002), no. 3, 355–405.
- [26] R. J. Gardner, *Geometric tomography*, second edition, *Encyclopedia of Mathematics and its Applications*, 58, Cambridge Univ. Press, New York, 2006. MR2251886
- [27] M. Henk and H. Pollehn, Necessary subspace concentration conditions for the even dual Minkowski problem, *Adv. Math.* **323** (2018), 114–141. MR3725875
- [28] J. Hosle, A. V. Kolesnikov and G. V. Livshyts, On the L_p -Brunn-Minkowski and dimensional Brunn-Minkowski conjectures for log-concave measures, *J. Geom. Anal.* **31** (2021), no. 6, 5799–5836. MR4267627
- [29] J. Hu, Uniqueness of solutions to the dual Minkowski problem, arXiv:2502.18032, 2025.
- [30] J. Hu, The dual Minkowski problem for positive indices, *Int. Math. Res. Not. IMRN* **2025**, no. 13, Paper No. rnaf192, 17 pp. MR4929415
- [31] Y. Hu and M. N. Ivaki, Stability of the cone-volume measure with near constant density, *Int. Math. Res. Not. IMRN* **2025**, no. 6, Paper No. rnaf062, 9 pp. MR4880436
- [32] Y. Huang, E. Lutwak, D. Yang and G. Zhang, Geometric measures in the dual Brunn-Minkowski theory and their associated Minkowski problems, *Acta Math.* **216** (2016), no. 2, 325–388. MR3573332
- [33] Y. Huang, D. Yang and G. Zhang, Minkowski problems for geometric measures, *Bull. Amer. Math. Soc. (N.S.)* **62** (2025), no. 3, 359–425. MR4926873
- [34] Y. Huang and Y. Zhao, On the L_p dual Minkowski problem, *Adv. Math.* **332** (2018), 57–84. MR3810248
- [35] A. V. Kolesnikov and G. V. Livshyts, On the Gardner-Zvavitch conjecture: symmetry in inequalities of Brunn-Minkowski type, *Adv. Math.* **384** (2021), Paper No. 107689, 23 pp. MR4238914
- [36] A. V. Kolesnikov and E. Milman, Brascamp-Lieb-type inequalities on weighted Riemannian manifolds with boundary, *J. Geom. Anal.* **27** (2017), no. 2, 1680–1702. MR3625169
- [37] A. V. Kolesnikov and E. Milman, Poincaré and Brunn-Minkowski inequalities on the boundary of weighted Riemannian manifolds, *Amer. J. Math.* **140** (2018), no. 5, 1147–1185. MR3862061
- [38] A. V. Kolesnikov and E. Milman, Local L^p -Brunn-Minkowski inequalities for $p < 1$, *Mem. Amer. Math. Soc.* **277** (2022), no. 1360, v+78 pp. MR4438690
- [39] Q. Li, J. Liu and J. Lu, Nonuniqueness of solutions to the L_p dual Minkowski problem, *Int. Math. Res. Not. IMRN* **2022**, no. 12, 9114–9150. MR4436203
- [40] Q. Li, W. Sheng and X. J. Wang, Flow by Gauss curvature to the Aleksandrov and dual Minkowski problems, *J. Eur. Math. Soc. (JEMS)* **22** (2020), no. 3, 893–923. MR4055992
- [41] E. Lutwak, Dual mixed volumes, *Pacific J. Math.* **58** (1975), no. 2, 531–538. MR0380631
- [42] E. Lutwak, The Brunn-Minkowski-Firey theory. I. Mixed volumes and the Minkowski problem, *J. Differential Geom.* **38** (1993), no. 1, 131–150. MR1231704

- [43] E. Lutwak, D. Yang and G. Zhang, L_p dual curvature measures, *Adv. Math.* **329** (2018), 85–132. MR3783409
- [44] H. Minkowski, *Ausgewählte Arbeiten zur Zahlentheorie und zur Geometrie*, Teubner-Archiv zur Mathematik, 12, Teubner, Leipzig, 1989. MR1064523
- [45] S. Mui, On the L^p dual Minkowski problem for $-1 < p < 0$, *Calc. Var. Partial Differential Equations* **63** (2024), no. 8, Paper No. 215, 19 pp. MR4797707
- [46] K. Patsalos and C. Saroglou, A note on the L_p -Brunn-Minkowski inequality for intrinsic volumes and the L_p -Christoffel-Minkowski problem, arXiv:2411.17896v4, 2026.
- [47] E. Putterman, Equivalence of the local and global versions of the L^p -Brunn-Minkowski inequality, *J. Funct. Anal.* **280** (2021), no. 9, Paper No. 108956, 20 pp. MR4220744
- [48] S. Sadosky and G. Zhang, Brunn-Minkowski and reverse isoperimetric inequalities for dual quermassintegrals, *Adv. Math.* **480** (2025), part A, Paper No. 110456, 14 pp. MR4941129
- [49] R. Schneider, *Convex bodies: the Brunn-Minkowski theory*, second expanded edition, *Encyclopedia of Mathematics and its Applications*, 151, Cambridge Univ. Press, Cambridge, 2014. MR3155183
- [50] J. Shan, The L_p dual Minkowski problem for group-invariant convex bodies, *Adv. Math.* **498** (2026), Paper No. 111038. MR5074845
- [51] D. Xi and Z. Zhang, The L_p Brunn-Minkowski inequalities for dual quermassintegrals, *Proc. Amer. Math. Soc.* **150** (2022), no. 7, 3075–3086. MR4428890
- [52] Y. Zhao, The dual Minkowski problem for negative indices, *Calc. Var. Partial Differential Equations* **56** (2017), no. 2, Paper No. 18, 16 pp. MR3605843
- [53] Y. Zhao, Existence of solutions to the even dual Minkowski problem, *J. Differential Geom.* **110** (2018), no. 3, 543–572. MR3880233
- [54] G. Zhu, The logarithmic Minkowski problem for polytopes, *Adv. Math.* **262** (2014), 909–931. MR3228445
- [55] G. Zhu, The centro-affine Minkowski problem for polytopes, *J. Differential Geom.* **101** (2015), no. 1, 159–174. MR3356071

INSTITUTE OF MATHEMATICS, HUNAN UNIVERSITY, CHANGSHA, 410082, CHINA

Email address: cxj@hnu.edu.cn

INSTITUTE OF MATHEMATICS, HUNAN UNIVERSITY, CHANGSHA, 410082, CHINA

Email address: tsy@hnu.edu.cn

INSTITUTE OF MATHEMATICS, HUNAN UNIVERSITY, CHANGSHA, 410082, CHINA

Email address: wangsinan@hnu.edu.cn