

Simultaneous Graph Learning and Matrix Recovery

Seyyid Emre Sofuoglu, Selin Aviyente

March 11, 2022

1 Methods

Our aim is to learn the underlying manifold of a data on a Cartesian Product Graph and recovery missing or grossly corrupted entries. The objective can be written in convex terms as:

$$\begin{aligned} & \text{minimize}_{L, S, \Phi_1, \Phi_2} \text{tr}(L^\top \Phi_1 L) + \text{tr}(L \Phi_2 L^\top) + \|S\|_1, \\ & \text{s.t. } \mathcal{P}_\Omega[Y] = \mathcal{P}_\Omega[L + S], \Phi_1, \Phi_2 \in \mathcal{S}, \end{aligned} \quad (1)$$

where Φ_1, Φ_2 are Graph Laplacians for the row graph and column graph, respectively, and \mathcal{S} is the space of undirected graph Laplacians. When the graph constraints are explicitly written, (1) becomes:

$$\begin{aligned} & \text{minimize}_{L, S, \Phi_1, \Phi_2} \alpha_1 \text{tr}(L^\top \Phi_1 L) + \alpha_2 \text{tr}(L \Phi_2 L^\top) + \|S\|_1 + \|\Phi_1\|_F^2 + \|\Phi_2\|_F^2, \\ & \text{s.t. } \mathcal{P}_\Omega[Y] = \mathcal{P}_\Omega[L + S], \Phi_i = \Phi_i^\top, \Phi_i \mathbf{1} = 0, \text{tr}(\Phi_i) = 2I_i. \end{aligned} \quad (2)$$

Since the graph Laplacians are assumed to be symmetric, the upper triangle portion is enough to summarize the conditions on them. This allows us to rewrite (2) as:

$$\begin{aligned} & \text{minimize}_{L, S, \phi_i \leq 0, \mathbf{d}_{\phi_i}} \sum_{i=1}^2 \alpha_i (2\mathbf{l}_i^\top \phi_i + \mathbf{d}_i^\top \mathbf{d}_{\phi_i} + \beta_{1,i} f(\mathbf{d}_{\phi_i}) + \beta_{2,i} g(\phi_i)) + \|S\|_1, \\ & \text{s.t. } \mathcal{P}_\Omega[Y] = \mathcal{P}_\Omega[L + S], P_i \phi_i = -\mathbf{d}_{\phi_i}, \end{aligned} \quad (3)$$

where $\mathbf{l}_i = \text{upper}(L_{(i)} L_{(i)}^\top)$, $\phi_i = \text{upper}(\Phi_i)$, $\mathbf{d}_i = \text{diag}(L_{(i)} L_{(i)}^\top)$, $\mathbf{d}_{\phi_i} = \text{diag}(\Phi_i)$, and $P_i \in \mathbb{R}^{I_i \times I_i(I_i-1)/2}$ is a matrix such that $P_i \text{upper}(W_i) = W_i \mathbf{1} - \text{diag}(W_i)$. $f(\cdot)$ is a function that controls the degree distribution and $g(\cdot)$ is a function that controls the sparsity of learned graph.

1.1 Optimization

(2) will be solved using ADMM. For (2) to be optimized, the graph variable L can be separated into two variables L_i , for $i \in \{1, 2\}$ to avoid large inverses. Corresponding augmented Lagrangian is written as:

$$\begin{aligned} & \|S\|_1 + \lambda_1 \|\mathcal{P}_\Omega[Y - L - S - \Gamma_1]\|_F^2 + \sum_{i=1}^2 \alpha_i \text{tr}(L_i^\top \Phi_i L_i) + \beta_{1,i} f(\mathbf{d}_{\phi_i}) + \beta_{2,i} g(\phi_i) + \lambda_{2,i} \|L_{(i)} - L_i - \Gamma_{2,i}\|_F^2 \\ & + \lambda_{3,i} \|P_i \phi_i + \mathbf{d}_{\phi_i} - \Gamma_{3,i}\|_F^2 + \lambda_{4,i} (\mathbf{1}^\top \mathbf{d}_{\phi_i} - I_i - \gamma_{4,i})^2 \end{aligned} \quad (4)$$

The variables are updated according to the following:

$$\mathcal{P}_\Omega[L^{t+1}] = \mathcal{P}_\Omega \left[\frac{1}{\lambda_1 + \sum_{i=1}^2 \lambda_{2,i}} \left(\lambda_1 Y - S^t - \Gamma_1^t + \sum_{i=1}^2 \lambda_{2,i} (L_{(i)}^t - \Gamma_{2,i}^t) \right) \right]. \quad (5)$$

$$\mathcal{P}_{\Omega^\perp}[L^{t+1}] = \mathcal{P}_{\Omega^\perp} \left[\sum_{i=1}^2 \frac{\lambda_{2,i}}{\sum_{i=1}^2 \lambda_{2,i}} (L_{(i)}^t - \Gamma_{2,i}^t) \right] \quad (6)$$

$$S^{t+1} = \sigma(Y - L^{t+1} - \Gamma_1^t, \lambda_1), \quad (7)$$

where $\sigma(\cdot)$ is the soft thresholding operator.

$$L_i^{t+1} = L_{i,inv}^t (L_i^{t+1} - \Gamma_{2,i}^t), \quad (8)$$

where $L_{i,inv} = (\mathbf{I} + \frac{\alpha_i}{\lambda_{2,i}} \Phi_i^t)^{-1}$.

$$\phi_i^{t+1} = \Pi_{\mathbb{R}_-^{I_i(I_i-1)/2}} \left[[-\beta_{2,i} \mathbf{I} + \lambda_{3,i} P_i^\top P_i]^{-1} (\mathbf{l}_i^{t+1} + \lambda_{3,i} P_i^\top (\mathbf{d}_{\phi_i}^t - \Gamma_{3,i}^t)) \right], \quad (9)$$

where $\Pi_{\mathbb{R}_-^{I_i(I_i-1)/2}}[\cdot]$ is an operator that projects to the negative orthant.

$$\mathbf{d}_{\phi_i}^{t+1} = [\beta_{1,i} \mathbf{I} + \lambda_{4,i} \mathbf{1} \mathbf{1}^\top]^{-1} [(\lambda_{4,i} I_i - \gamma_{4,i}^t) \mathbf{1} - \alpha_i \mathbf{d}_{\mathbf{l}_i}^{t+1} + \Gamma_{3,i}^t] \quad (10)$$

Finally, dual variables are updated as follows:

$$\Gamma_1^{t+1} = \Gamma_1^t - \mathcal{P}_\Omega[Y - L^{t+1} - S^{t+1}] \quad (11)$$

$$\Gamma_{2,i}^{t+1} = \Gamma_{2,i}^t - (L_{(i)}^{t+1} - L_i^{t+1}) \quad (12)$$

$$\Gamma_{3,i}^{t+1} = \Gamma_{3,i}^t - (P_i \phi_i + \mathbf{d}_{\phi_i}) \quad (13)$$

$$\gamma_{4,i}^{t+1} = \gamma_{4,i}^t - (\mathbf{1}^\top \mathbf{d}_{\phi_i} - I_i) \quad (14)$$