

Ordinary differential equations

Nguyen Thu Huong



School of Applied Mathematics and Informatics
Hanoi University of Science and Technology

October 28, 2020

ODEs

Differential
equations

Motivation

Basic concepts

First order
differential
equations

Separable Equations

Homogeneous
equationsExact differential
equations

Linear equations

Bernoulli equations

1 Differential equations

- Motivation
- Basic concepts

2 First order differential equations

- Separable Equations
- Homogeneous equations
- Exact differential equations
- Linear equations
- Bernoulli equations

Content

1 Differential equations

- Motivation
- Basic concepts

2 First order differential equations

- Separable Equations
- Homogeneous equations
- Exact differential equations
- Linear equations
- Bernoulli equations

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Mathematical models of many phenomena in physics, biology, economy, . . . result in ordinary differential equations.

- 1 Population model: $\frac{dN}{dt} = kN$.
- 2 Vibration of a spring: $mx'' + kx = 0$.
- 3 Electrical circuits: $\frac{dI}{dt} + \frac{R}{L}I = \frac{E}{L}$.
- 4 Oscillation equation of a pendulum: $x''(t) + \frac{g}{l} \sin x = 0$.

ODEs

Differential
equations

Motivation

Basic concepts

First order
differential
equations

Separable Equations

Homogeneous
equationsExact differential
equations

Linear equations

Bernoulli equations

Definition

An **ordinary differential equation** is an equation involving an unknown function (of one variable) and its derivatives.

$$F(x, y, y', y'', \dots, y^{(n)}) = 0,$$

where x is a variable, $y = y(x)$ is the function in search, and $y', y'', \dots, y^{(n)}$ are the derivatives of y .

Definition

The **order** of an ODE is the order of the highest derivative appearing in the equation.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Example

$$1 \quad y''' - 3xy' + y^2 = 0.$$

$$2 \quad y'y'' - y^3 \cos x + xy = 0.$$

$$3 \quad xy'' - (1 + x^2)y' + 5y = \tan x.$$

$$4 \quad \sin y \frac{dy}{dx} - 2x^3y + x^4 = 0.$$

$$5 \quad e^x \frac{d^3u}{dx^3} + 2 \left(\frac{du}{dx} \right)^2 = x^3.$$

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Definition

A **linear** ODE is an ODE where F is linear with respect to $y, y', y'', \dots, y^{(n)}$.

General form

$$y^{(n)} + a_1(x)y^{(n-1)} + \dots + a_{n-1}(x)y' + a_n(x)y = f(x),$$

where $a_1(x), \dots, a_{n-1}(x), a_n(x), f(x)$ are given functions.

ODEs

Differential
equations

Motivation

Basic concepts

First order
differential
equations

Separable Equations

Homogeneous
equationsExact differential
equations

Linear equations

Bernoulli equations

Definition

A **solution** of an ODE is a function $y = y(x)$, $x \in I$, which satisfies the equation identically for all $x \in I$.

General solution of an ODE is the set of all solutions depending on parameters, which can be found once additional conditions are given.

A **particular solution** of an ODE is any solution obtained from the general solution by specifying values of the parameters.

A **singular solution** of an ODE is a solution that cannot be obtained from the general solution.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

The solutions can be given in explicit / implicit forms, or by parametrization

- Explicitly: general solution $y = \varphi(x, C)$; particular solution $y = \varphi(x, C_0)$.
- Implicitly: general integral $\Phi(x, y, C) = 0$; particular integral $\Phi(x, y, C_0) = 0$.
- Parametrization:
$$\begin{cases} x = x(t, C) \\ y = y(t, C). \end{cases}$$

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Example

1 $y' = f(x)$, the general solution is $y = \int f(x)dx + C$.

2 Oscillation equation of a spring $kx'' + mx = 0$.

- General solution $x(t) = C_1 \cos \omega t + C_2 \sin \omega t$.
- Observe at the time of release $t = 0$: e.g. $x(0) = A_0$, $x'(0) = 0$.

We obtain a particular solution $x(t) = A_0 \cos(\omega t)$,
 $C_1 = A_0$, $C_2 = 0$.

Initial value and Boundary value problems

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

- Initial conditions \Rightarrow IVP.
Example: Oscillation of a spring $x(0) = A, x'(0) = 0$.
- Boundary conditions \Rightarrow BVP.
Example: Oscillation of a string $x'(0) = x'(1) = 0$.

Content

1 Differential equations

- Motivation
- Basic concepts

2 First order differential equations

- Separable Equations
- Homogeneous equations
- Exact differential equations
- Linear equations
- Bernoulli equations



Separable equations



ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

General form: $f(x)dx = g(y)dy$

Integrating both sides of the equation:

$$\int f(x)dx = \int g(y)dy \Rightarrow F(x) = G(y) + C,$$

where $F(x)$, $G(y)$ are antiderivatives of $f(x)$, $g(y)$ respectively.



ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Example (20182)

Solve the ODE $y' = 2xy^2$.

- $y = 0$ is a solution of the equation.
- $y \neq 0$, the equation becomes $\frac{dy}{y^2} = 2xdx$. Integrating both sides, we get

$$\int \frac{dy}{y^2} = \int 2xdx \Rightarrow -\frac{1}{y} = x^2 + C.$$

Hence, the solutions are $y = -\frac{1}{x^2 + C}$ and $y = 0$.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Example (20181)

Solve the following problem $y' = 3 + xy + x + 3y, y(0) = 1$.

The equation is equivalent to $\frac{dy}{dx} = (x + 3)(y + 1)$.

- $y + 1 = 0 \Rightarrow y = -1$ does not satisfy the condition $y(0) = 1$, hence it is not a solution.
- $y + 1 \neq 0$, ta có

$$\begin{aligned}\frac{dy}{y+1} &= (x+3)dx \Rightarrow \int \frac{dy}{y+1} = \int (x+3)dx \\ &\Rightarrow \ln|y+1| = \frac{x^2}{2} + 3x + \ln|C|, C \neq 0 \\ &\Rightarrow y+1 = Ce^{\frac{x^2}{2}+3x}.\end{aligned}$$

Plugging the condition in the solution, we obtain $C = 2$.

Hence, the solution is $y + 1 = 2e^{\frac{x^2}{2}+3x}$.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

General form: $\frac{dy}{dx} = f(x, y)$ where $f(tx, ty) = f(x, y)$.

Or $y' = g\left(\frac{y}{x}\right)$

We transform it into a separable equation as follows:

- Making a substitution $y = ux$.
- The resulting equation is $x \frac{du}{dx} = f(u) - u \Rightarrow u(x)$.
- Substituting back we get $y(x)$.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Example (20181)

Solve the following problem $y' = \frac{-x + 2y}{x}$, $y(1) = 2$.

Set $y = x.u$, the equation becomes

$$xu' + u = -1 + 2u \Leftrightarrow x \frac{du}{dx} = u - 1.$$

$y(1) = 2$ so $u(1) = \frac{y(1)}{1} = 2$. $u = 1$ does not satisfy the condition, hence it is not a solution of the problem.

$u \neq 1$, the equation can be rewritten as

$$\begin{aligned} \frac{du}{u-1} &= \frac{dx}{x} \Rightarrow \int \frac{du}{u-1} = \int \frac{dx}{x} \\ &\Rightarrow \ln |u-1| = \ln |x| + \ln |C|, (C \neq 0) \Rightarrow \frac{y}{x} - 1 = Cx. \end{aligned}$$

Using $y(1) = 2$, we get $C = 1$. The solution is $y = x(x + 1)$.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

General form

$$P(x, y)dx + Q(x, y)dy = 0,$$

where $P(x, y)$, $Q(x, y)$ are continuous functions and have continuous first partial derivatives on some rectangle D of the plane and $\frac{\partial P}{\partial y} = \frac{\partial Q}{\partial x}$.

Under these conditions, we can find a function $u(x, y)$ such that

$$P = \frac{\partial u}{\partial x}, Q = \frac{\partial u}{\partial y}.$$

The equation reads $du = 0$, hence the general solution is given implicitly by:

$$u(x, y) = C.$$

ODEs

Differential
equations

Motivation

Basic concepts

First order
differential
equations

Separable Equations

Homogeneous
equationsExact differential
equations

Linear equations

Bernoulli equations

$u(x, y)$ is given by:

$$\begin{aligned} u(x, y) &= \int_{x_0}^x P(x, y_0) dx + \int_{y_0}^y Q(x, y) dy \\ &= \int_{x_0}^x P(x, y) dx + \int_{y_0}^y Q(x_0, y) dy. \end{aligned}$$

where $(x_0, y_0) \in D$.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Example (CK20181)

Solve the ODE $(3x^2 - 6xy)dx - (3x^2 + 4y^3)dy = 0$.

- $P(x, y) = 3x^2 - 6xy$, $Q = -3x^2 - 4y^3$.
 $P'_y = Q'_x = -6x \Rightarrow$ exact differential equation.
- The general integral is given by:

$$\begin{aligned}
 u(x, y) &= \int_0^x P(x, 0)dx + \int_0^y Q(x, y)dy = C \\
 &\Leftrightarrow \int_0^x 3x^2 dx + \int_0^y (-3x^2 - 4y^3)dy = C \\
 &\Leftrightarrow x^3 - (3x^2y + y^4) \Big|_0^y = C \\
 &\Leftrightarrow x^3 - 3x^2y - y^4 = C.
 \end{aligned}$$

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

We can also find $u(x, y)$ by solving the system

$$\begin{cases} u'_x = 3x^2 - 6xy \\ u'_y = -3x^2 - 4y^3. \end{cases}$$

The first equation yields that

$$u = \int (3x^2 - 6xy) dx = x^3 - 3x^2y + C(y).$$

Plugging into the second equation, we get

$$u'_y = -3x^2 + C'(y) = -3x^2 - 4y^3,$$

we obtain $C'(y) = -4y^3 \Rightarrow C(y) = -y^4$.

Hence, $u = x^3 - 3x^2y - y^4$, the general integral is $x^3 - 3x^2y - y^4 = C$.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

In general, the equation $P(x, y)dx + Q(x, y)dy = 0$ is not an exact DE.

A function $\alpha(x, y)$ is called **integrating factor** if

$$\alpha(x, y)[P(x, y)dx + Q(x, y)dy] = 0$$

is an exact DE, which means $\frac{\partial(\alpha P)}{\partial y} = \frac{\partial(\alpha Q)}{\partial x}$.

Particular cases of the integrating factor

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

- If $\frac{Q'_x - P'_y}{Q} = \varphi(x) \Rightarrow \alpha(x, y) = \alpha(x) = e^{-\int \varphi(x) dx}$
- If $\frac{Q'_x - P'_y}{P} = \psi(y) \Rightarrow \alpha(x, y) = \alpha(y) = e^{\int \psi(y) dy}$

Example (20182)

Solve the problem $e^y dx + (9y + 4xe^y) dy = 0, y(1) = 0$.

$P = e^y, Q = 9y + 4xe^y \Rightarrow \frac{Q'_x - P'_y}{P} = \frac{4e^y - e^y}{e^y} = 3$ hence, an integrating factor is $\alpha(y) = e^{3y}$.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Multiplying through by e^{3y} , we obtain
 $e^{4y} dx + (9ye^{3y} + 4xe^{4y}) dy = 0$ (exact equation).

$$u(x, y) = \int_{x_0}^x P(x, y_0) dx + \int_{y_0}^y Q(x, y) dy = C$$

In particular $y(x_0) = y_0$, we get $C = 0$.

The integral of the problem is

$$\int_1^x dx + \int_0^y (9ye^{3y} + 4xe^{4y}) dy = 0.$$



ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

General form

$$y' + p(x)y = q(x),$$

where $p(x), q(x)$ are continuous function on $I \subset \mathbb{R}$.

ODEs

$$y' + p(x)y = q(x) \Rightarrow (p(x)y - q(x))dx + dy = 0.$$

- $P = p(x)y - q(x)$, $Q = 1$, $\frac{Q'_x - P'_y}{Q} = -p(x)$, an integrating factor is $\alpha(x) = e^{\int p(x)dx}$.
- Multiplying both sides by $\alpha(x)$, we get

$$\begin{aligned}(y' + p(x)y)e^{\int p(x)dx} &= q(x)e^{\int p(x)dx} \\ \Leftrightarrow \left(ye^{\int p(x)dx} \right)' &= q(x)e^{\int p(x)dx} \\ \Rightarrow y &= e^{-\int p(x)dx} \left(\int q(x)e^{\int p(x)dx} dx + C \right).\end{aligned}$$

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

The general solution is given by

$$y = \left(\int q(x) e^{\int p(x) dx} dx + C \right) e^{-\int p(x) dx}.$$

Example (20182)

Solve the ODE $y' - 2y \tan x = 2 \sin 2x$.

$$\begin{aligned}
 y &= e^{\int 2 \tan x dx} \left(\int 2 \sin 2x e^{-\int 2 \tan x dx} dx + C \right) \\
 \Rightarrow y &= \frac{1}{\cos^2 x} \left(\int 2 \sin 2x \cos^3 x dx + C \right) \\
 \Rightarrow y &= \frac{C - \cos^4 x}{\cos^2 x}.
 \end{aligned}$$

Structure of the general solutions of linear equations

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

The equation $y' + p(x)y = q(x)$ has the general solution

$$y = \left(\int q(x) e^{\int p(x) dx} dx + C \right) e^{-\int p(x) dx} = y^* + \bar{y},$$

where

- $\bar{y} = C e^{-\int p(x) dx}$ is the general solution of the corresponding homogeneous equation $y' + p(x)y = 0$.
- y^* is a particular solution of the given inhomogeneous equation.

Variation of constants:

We look for y^* of the form $y^* = C(x) e^{-\int p(x) dx}$ and substitute in the equation to find $C(x)$.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

General form:

$$y' + p(x)y = q(x)y^\alpha, \alpha \neq 0, 1.$$

- 1 Verify whether $y = 0$ is a solution.
- 2 $y \neq 0$, set $v = y^{1-\alpha}$, the equation becomes

$$v' + (1 - \alpha)p(x)v = (1 - \alpha)q(x),$$

which is a linear equation.

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

The resulting equation has the general solution given by

$$v = \bar{v} + v^* = \left(\int (1-\alpha)q(x)e^{\int (1-\alpha)p(x)dx} dx + C \right) e^{-\int (1-\alpha)p(x)dx}.$$

Substitute back, we have $y = v^{\frac{1}{1-\alpha}}$.

Example

Solve the ODE $y' + xy = x^3y^3$.

Bernoulli equation, $\alpha = 3$.

- $y = 0$ is a solution.
- $y \neq 0$. Dividing both sides by y^3 , we obtain

$$\frac{y'}{y^3} + x \frac{1}{y^2} = x^3.$$

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Set $z = \frac{1}{y^2} \Rightarrow z' = -\frac{2y'}{y^3}$, the equation becomes

$$-\frac{z'}{2} + xz = x^3 \Leftrightarrow z' - 2xz = -2x^3.$$

(linear equation in z).

The general solution is

$$\begin{aligned} \frac{1}{y^2} &= e^{\int 2x dx} \left(C - 2 \int x^3 e^{-\int 2x dx} dx \right) \\ &= e^{x^2} \left(C - \int x^2 e^{-x^2} d(x^2) \right) \\ &= e^{x^2} \left(C + (x^2 + 1)e^{-x^2} \right). \end{aligned}$$

ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Theorem

Assume that $f(x, y): D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ is **continuous** on D , and $(x_0, y_0) \in D$. Then, in some interval $(x_0 - h, x_0 + h)$, there exists one solution $y = y(x)$ of the IVP

$$\begin{cases} y' = f(x, y), & x \in U_\varepsilon(x_0), \\ y(x_0) = y_0. \end{cases}$$

If additionally $\frac{\partial f}{\partial y}(x, y)$ is **continuous** in D then the solution is **unique**.



ODEs

Differential equations

Motivation

Basic concepts

First order differential equations

Separable Equations

Homogeneous equations

Exact differential equations

Linear equations

Bernoulli equations

Solve the following ODEs

1 $xy' = y \ln \frac{y}{x}.$

2 $y' - 2y \tan x + y^2 \sin^2 x = 0.$

3 $2y'\sqrt{x} = \sqrt{1 - y^2}.$

4 $(e^x + y + \sin y)dx + (e^y + x + x \cos y)dy = 0.$

5 $y' = \sin(y - x - 1).$

6 $(x^2 + y)dx + (x - 2y)dy = 0.$

7 $y' + y \cos x = \sin x \cos x.$

8 $y' + \frac{y}{x} = x^2 y^4.$

9 $\tan y dx - x \ln x dy = 0.$