

25 YEARS ANNIVERSARY
SOICT

ĐẠI HỌC BÁCH KHOA HÀ NỘI
VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG



ĐẠI HỌC BÁCH KHOA HÀ NỘI
VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG

PRESENTATION TITLE

Presentation Subtitle


IV. Stochastic processes

- Definitions
- Stationary processes
- Spectral density function
- Ergodicity of stochastic processes

IV. Stochastic processes

4.1. Definitions

- Stochastic processes
 - Let ξ denote the random outcome of an experiment. To every such outcome suppose a waveform $X(t, \xi)$ is assigned.
 - The collection of such waveforms form a stochastic process.



IV. Stochastic processes

4.1. Definitions

- The set of $\{\xi_k\}$ and the time index t can be continuous or discrete (countably infinite or finite) as well.
- For fixed $\xi_i \in S$ (the set of all experimental outcomes), $X(t, \xi_i)$ is a specific time function.
- For fixed t , $X_1=X(t_1, \xi_i)$ is a random variable. The ensemble of all such realizations $X(t, \xi)$ over time represents the stochastic process $X(t)$.

IV. Stochastic processes

4.1. Definitions

- Example
 - $X(t) = A \cos(\omega_0 t + \varphi)$
 - Where φ is a uniformly distributed random variable in $(0, 2\pi)$ represents a stochastic process.
- Some stochastic processes
 - Brownian motion,
 - Stock market fluctuations,
 - Various queuing systems.

IV. Stochastic processes

4.1. Definitions

- If $X(t)$ is a stochastic process, then for fixed t , $X(t)$ represents a random variable.
- Its distribution function is given by

$$F_x(x, t) = P\{X(t) \leq x\}$$

- Notice that $F_x(x, t)$ depends on t , since for a different t , we obtain a different random variable.

$$f_x(x, t) \triangleq \frac{dF_x(x, t)}{dx}$$

- Derivative of $F_x(x, t)$ represents the first-order probability density function of the process $X(t)$.

IV. Stochastic processes

4.1. Definitions

- For $t = t_1$ and $t = t_2$,
 - $X(t)$ represents two different random variables $X_1 = X(t_1)$ and $X_2 = X(t_2)$ respectively.
 - Their joint distribution is given by

$$F_x(x_1, x_2, t_1, t_2) = P\{X(t_1) \leq x_1, X(t_2) \leq x_2\}$$

- Their joint density function is:

$$f_x(x_1, x_2, t_1, t_2) \triangleq \frac{\partial^2 F_x(x_1, x_2, t_1, t_2)}{\partial x_1 \partial x_2}$$

- And represents the second-order density function of the process $X(t)$.

IV. Stochastic processes

4.1. Definitions

- Similarly $f_x(x_1, \dots, x_n, t_1, \dots, t_n)$ represents the n^{th} order density function of the process $X(t)$.
- Complete specification of the stochastic process $X(t)$ requires the knowledge of $f_x(x_1, \dots, x_n, t_1, \dots, t_n)$ for all t_i , $i = 1, \dots, n$ and for all n . (an almost impossible task in reality).

IV. Stochastic processes

4.1. Definitions

- Characteristics of stochastic processes
 - Mean of a stochastic process

$$\mu(t) \triangleq E\{X(t)\} = \int_{-\infty}^{+\infty} x f_X(x, t) dx$$

- $\mu(t)$ represents the mean value of a process $X(t)$. In general, the mean of a process can depend on the time index t .
- Autocorrelation function of a process $X(t)$ is defined as

$$R_{XX}(t_1, t_2) \triangleq E\{X(t_1)X^*(t_2)\} = \iint x_1 x_2^* f_X(x_1, x_2, t_1, t_2) dx_1 dx_2$$

• It represents the interrelationship between the random variables $X_1 = X(t_1)$ and $X_2 = X(t_2)$ generated from the process $X(t)$.

IV. Stochastic processes

4.1. Definitions

- Properties of autocorrelation function

$$1. \quad R_{xx}(t_1, t_2) = R_{xx}^*(t_2, t_1) = [E\{X(t_2)X^*(t_1)\}]^*$$

$$2. \quad R_{xx}(t, t) = E\{|X(t)|^2\} \geq 0.$$

(Average instantaneous power)

- 3. $R_{xx}(t_1, t_2)$ represents a nonnegative definite function, i.e., for *any* set of constants $\{a_i\}_{i=1}^n$

$$\sum_{i=1}^n \sum_{j=1}^n a_i a_j^* R_{xx}(t_i, t_j) \geq 0.$$

- This equation follows by noticing that: $E\{|Y|^2\} \geq 0$ and

$$Y = \sum_{i=1}^n a_i X(t_i).$$

IV. Stochastic processes

4.1. Definitions

- The **autocovariance** function of the process $X(t)$.

$$C_{xx}(t_1, t_2) = R_{xx}(t_1, t_2) - \mu_x(t_1)\mu_x^*(t_2)$$

- Examples
 - Given

$$z = \int_{-T}^T X(t)dt.$$

$$\begin{aligned} E[|z|^2] &= \int_{-T}^T \int_{-T}^T E\{X(t_1)X^*(t_2)\}dt_1dt_2 \\ &= \int_{-T}^T \int_{-T}^T R_{xx}(t_1, t_2)dt_1dt_2 \end{aligned}$$

IV. Stochastic processes

4.1. Definitions

- Consider process $X(t)$

$$X(t) = a \cos(\omega_0 t + \varphi), \quad \varphi \sim U(0, 2\pi).$$

$$\mu_x(t) = E\{X(t)\} = aE\{\cos(\omega_0 t + \varphi)\}$$

$$= a \cos \omega_0 t E\{\cos \varphi\} - a \sin \omega_0 t E\{\sin \varphi\} = 0,$$

$$\text{since } E\{\cos \varphi\} = \frac{1}{2\pi} \int_0^{2\pi} \cos \varphi d\varphi = 0 = E\{\sin \varphi\}.$$

$$R_{xx}(t_1, t_2) = a^2 E\{\cos(\omega_0 t_1 + \varphi) \cos(\omega_0 t_2 + \varphi)\}$$

$$= \frac{a^2}{2} E\{\cos \omega_0(t_1 - t_2) + \cos(\omega_0(t_1 + t_2) + 2\varphi)\}$$

$$= \frac{a^2}{2} \cos \omega_0(t_1 - t_2).$$

IV. Stochastic processes

4.2. Stationary processes

- Strict sense and wide sense stationarity
 - Stationarity
 - Stationary processes exhibit statistical properties that are invariant to shift in the time index.
 - Thus, for example, second-order stationarity implies that the statistical properties of the pairs $\{X(t_1), X(t_2)\}$ and $\{X(t_1+c), X(t_2+c)\}$ are the same for *any* c .
 - Similarly first-order stationarity implies that the statistical properties of $X(t_i)$ and $X(t_i+c)$ are the same for any c .

IV. Stochastic processes

4.2. Stationary processes

- Strict sense stationarity
 - In strict terms, the statistical properties of a stochastic process are governed by the joint probability density function.
 - A process is n^{th} -order **Strict-Sense Stationary (S.S.S)** if

for any c , where the left side represents the joint density function of the random variables $X_1 = X(t_1), \dots, X_n = X(t_n)$ and the right side corresponds to the joint density function of the random variables $X'_1 = X(t_1 + c), \dots, X'_n = X(t_n + c)$.

- A process $X(t)$ is said to be **strict-sense stationary** if equation above is true for all $t_i, i=1, \dots, n; n=1, 2, \dots$ and any c .

IV. Stochastic processes

4.2. Stationary processes

- First order strict sense stationary process
 - For any c

$$f_x(x, t) \equiv f_x(x, t + c)$$

- In particular, if $c = -t$, then

$$f_x(x, t) = f_x(x)$$

- That means, the first-order density of $X(t)$ is independent of t . In that case

$$E[X(t)] = \int_{-\infty}^{+\infty} x f(x) dx = \mu, \text{ a constant.} \quad (1)$$

IV. Stochastic processes

4.2. Stationary processes

- Second order strict sense stochastic process
 - From definition, we have:

$$f_x(x_1, x_2, t_1, t_2) \equiv f_x(x_1, x_2, t_1 + c, t_2 + c)$$

- for any c. If c is chosen so as $c = -t_2$, we get

- $f_x(x_1, x_2, t_1, t_2) \equiv f_x(x_1, x_2, t_1 - t_2)$
- The second order density function of a strict sense stationary process depends only on the difference of the time indices $t_1 - t_2 = \tau$.

IV. Stochastic processes

4.2. Stationary processes

- The autocorrelation function is given by

$$\begin{aligned} R_{xx}(t_1, t_2) &\triangleq E\{X(t_1)X^*(t_2)\} \\ &= \iint x_1 x_2^* f_x(x_1, x_2, \tau = t_1 - t_2) dx_1 dx_2 \\ &= R_{xx}(t_1 - t_2) \triangleq R_{xx}(\tau) = R_{xx}^*(-\tau), \quad (2) \end{aligned}$$

- The autocorrelation function of a second order strict-sense stationary process depends only on the difference of the time indices $t_2 - t_1 = \tau$

IV. Stochastic processes

4.2. Stationary processes

- Wide sense stationarity
 - A process $X(t)$ is said to be **Wide-Sense Stationary** if:
 - $E\{X(t)\} = \mu$ and $E\{X(t_1)X^*(t_2)\} = R_{xx}(t_1 - t_2)$,
 - Since these equations follow from (1) and (2), strict-sense stationarity always implies wide-sense stationarity.
 - In general, the converse is *not true*
 - Exception: the Gaussian process (normal process).
 - This follows, since if $X(t)$ is a Gaussian process, then by definition $X_1=X(t_1), \dots, X_n=X(t_n)$ are jointly Gaussian random variables for any t_1, \dots, t_n whose joint characteristic function is given by

$$\phi_x(\omega_1, \omega_2, \dots, \omega_n) = e^{j \sum_{k=1}^n \mu(t_k) \omega_k - \sum_{l,k}^n \sum_{l,k} C_{xx}(t_l, t_k) \omega_l \omega_k / 2}$$

IV. Stochastic processes

4.2. Stationary processes

- Examples

- The process:

$$X(t) = a \cos(\omega_0 t + \varphi), \quad \varphi \sim U(0, 2\pi).$$

- This process is wide sense stationary but not strict sense stationary.

- If the process $X(t)$ has zero mean,


then σ_z^2 is reduced to:

$$z = \int_{-T}^T X(t) dt.$$

$$\sigma_z^2 = E\{|z|^2\} = \int_{-T}^T \int_{-T}^T R_{xx}(t_1 - t_2) dt_1 dt_2.$$

- As t_1 and t_2 varies from $-T$ to T , so $\tau = t_2 - t_1$ varies from $-2T$ to $2T$.

- $R_{xx}(\tau)$ is a constant over the shaded region in the figure on the left.



IV. Stochastic processes

4.3. Power spectrum

- Power spectrum

- For a deterministic signal $x(t)$


- The spectrum is well defined: If $X(\omega)$ represents its Fourier transform,

$$X(\omega) = \int_{-\infty}^{+\infty} x(t) e^{-j\omega t} dt,$$

- then $|X(\omega)|^2$ represents its energy spectrum. This follows from Parseval's theorem since the signal energy is given by

$$\int_{-\infty}^{+\infty} x^2(t) dt = \frac{1}{2\pi} \int_{-\infty}^{+\infty} |X(\omega)|^2 d\omega = E.$$

- Thus $|X(\omega)|^2 \Delta\omega$ represents the signal energy in the band $(\omega, \omega + \Delta\omega)$



IV. Stochastic processes

4.3. Power spectrum

- For stochastic processes,
 - A direct application of Fourier transform generates a sequence of random variables for every ω
 - For a stochastic process, $E\{|X(t)|^2\}$ represents the ensemble average power (instantaneous energy) at the instant t .
 - Partial Fourier transform of a process $X(t)$ based on $(-T, T)$ is given by

$$X_T(\omega) = \int_{-T}^T X(t) e^{-j\omega t} dt$$

- The power distribution associated with that realization based on $(-T, T)$ is represented by

$$\frac{|X_T(\omega)|^2}{2T} = \frac{1}{2T} \left| \int_{-T}^T X(t) e^{-j\omega t} dt \right|^2$$

IV. Stochastic processes

4.3. Power spectrum

- The average power distribution based on $(-T, T)$ is ensemble average of power distribution for ω

$$P_T(\omega) = E \left\{ \frac{|X_T(\omega)|^2}{2T} \right\} = \frac{1}{2T} \int_{-T}^T \int_{-T}^T E\{X(t_1)X^*(t_2)\} e^{-j\omega(t_1-t_2)} dt_1 dt_2$$
$$= \frac{1}{2T} \int_{-T}^T \int_{-T}^T R_{xx}(t_1, t_2) e^{-j\omega(t_1-t_2)} dt_1 dt_2$$

- We have this represents the power distribution of $X(t)$ based on $(-T, T)$.
- If $X(t)$ is assumed to be w.s.s, then
- and we have $R_{xx}(t_1, t_2) = R_{xx}(t_1 - t_2)$

$$P_T(\omega) = \frac{1}{2T} \int_{-T}^T \int_{-T}^T R_{xx}(t_1 - t_2) e^{-j\omega(t_1-t_2)} dt_1 dt_2.$$

IV. Stochastic processes

4.3. Power spectrum

- Let $\tau = t_2 - t_1$, we obtain

$$P_T(\omega) = \frac{1}{2T} \int_{-2T}^{2T} R_{xx}(\tau) e^{-j\omega\tau} (2T - |\tau|) d\tau$$

$$= \int_{-2T}^{2T} R_{xx}(\tau) e^{-j\omega\tau} \left(1 - \frac{|\tau|}{2T}\right) d\tau \geq 0$$

- This is the power distribution of the w.s.s. process $X(t)$ based on $(-T, T)$.
- Letting $T \rightarrow \infty$, we obtain

$$S_{xx}(\omega) = \lim_{T \rightarrow \infty} P_T(\omega) = \int_{-\infty}^{+\infty} R_{xx}(\tau) e^{-j\omega\tau} d\tau \geq 0$$

- $S_{xx}(\omega)$ is the *power spectral density* of the w.s.s process $X(t)$.

IV. Stochastic processes

4.3. Power spectrum

- Khinchin-Wiener theorem
 - The autocorrelation function and the power spectrum of a w.s.s process form a Fourier transform pair.

$$R_{xx}(\omega) \xleftrightarrow{\text{F.T.}} S_{xx}(\omega) \geq 0.$$

$$R_{xx}(\tau) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} S_{xx}(\omega) e^{j\omega\tau} d\omega$$


- For $\tau=0$, $S_{xx}(\omega) = \lim_{T \rightarrow \infty} P_T(\omega) = \int_{-\infty}^{+\infty} R_{xx}(\tau) e^{-j\omega\tau} d\tau \geq 0$

$$\frac{1}{2\pi} \int_{-\infty}^{+\infty} S_{xx}(\omega) d\omega = R_{xx}(0) = E\{|X(t)|^2\} = P, \quad \text{the total power.}$$

IV. Stochastic processes

4.3. Power spectrum

- The area under $S_{xx}(\omega)$ represents the total power of the process $X(t)$, and hence $S_{xx}(\omega)$ truly represents the power spectrum.



- The nonnegative-definiteness property of the auto-correlation function translates into the “nonnegative” property for its Fourier transform (power spectrum).

$$R_{xx}(\tau) \text{ nonnegative - definite} \Leftrightarrow S_{xx}(\omega) \geq 0.$$

IV. Stochastic processes

4.3. Power spectrum

- If $X(t)$ is a real w.s.s process, then

$$R_{XX}(\tau) = R_{XX}(-\tau)$$

- So that

$$\begin{aligned} S_{XX}(\omega) &= \int_{-\infty}^{+\infty} R_{XX}(\tau) e^{-j\omega\tau} d\tau \\ &= \int_{-\infty}^{+\infty} R_{XX}(\tau) \cos \omega\tau d\tau \end{aligned}$$

- The power spectrum is an even function, (in addition to being real and nonnegative).

IV. Stochastic processes

4.4. Ergodicity

- Time averages

- Given wide-sense stationary process $X(t)$.

- Time averages

- Mean

$$\bar{x} = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T x(t) dt$$

- Autocorrelation

$$R(\tau) = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T x(t + \tau) x(t) dt$$

- These limits are random variables.
- Problems:

$$\bar{x} \stackrel{?}{=} E\{x(t)\} \quad R(\tau) \stackrel{?}{=} E\{x(t + \tau)x(t)\}$$

IV. Stochastic processes

4.4. Ergodicity

- Ergodicity
 - $X(t)$ is ergodic if in the most general form if all its statistics can be determined from a single function $X(t, \zeta)$ of the process.
 - $X(t)$ is ergodic if time averages equal ensemble averages (expected values)

IV. Stochastic processes

4.4. Ergodicity

- Ergodicity of the mean
 - Time average of a given process $X(t)$

$$n_T = \frac{1}{2T} \int_{-T}^T x(t) dt$$

- n_T is a random variable.
- Since $E\{X(t)\}$ is a constant, we have

$$E\{n_T\} = E\{X(t)\} = \eta$$

- The variance of n_T is given by:

$$\sigma_{n_T}^2 = \frac{1}{T} \int_0^{2T} \left(1 - \frac{\tau}{2T}\right) [R(\tau) - \eta^2] d\tau$$

- $R(\tau)$ is the autocorrelation of $X(t)$.
- If this variance tends to zero with $T \rightarrow \infty$, then n_T tends to its expected value.

IV. Stochastic processes

4.4. Ergodicity

- Ergodic theorem for $E\{X(t)\}$

$$\lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T \mathbf{x}(t) dt = E\{\mathbf{x}(t)\} = \boldsymbol{\eta}$$

iff

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^{2T} \left(1 - \frac{\tau}{2T}\right) [R(\tau) - \boldsymbol{\eta}^2] d\tau = 0$$

IV. Stochastic processes

4.4. Ergodicity

- Ergodicity of autocorrelation
 - We form the average:

$$R_T(\lambda) = \frac{1}{2T} \int_{-T}^T x(t + \lambda)x(t)dt$$

- We have

$$E\{R_T(\lambda)\} = \frac{1}{2T} \int_{-T}^T E\{x(t + \lambda)x(t)\}dt = R(\lambda)$$

- For a given λ , $R_T(\lambda)$ is the time average of the process $\Phi(t)=x(t+\lambda)x(t)$
- The mean of the process $\Phi(t)$ is given by

$$E\{\Phi(t)\}=E\{x(t+\lambda)x(t)\} = R(\lambda)$$

IV. Stochastic processes

4.4. Ergodicity

- Its autocorrelation
 $R_{\Phi\Phi}(\tau) = E\{x(t+\lambda+\tau)x(t+\tau)x(t+\lambda)x(t)\}$
- Hence with $w(t) = \Phi(t)$, we have
- Ergodicity theorem for autocorrelation
 - For a given λ

$$\lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T x(t + \lambda)x(t) dt = E\{x(t + \lambda)x(t)\} = R(\lambda)$$

iff

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^{2T} \left(1 - \frac{\tau}{2T}\right) [R_{\Phi\Phi}(\tau) - R^2(\lambda)] d\tau = 0$$

IV. Stochastic processes

4.4. Ergodicity

- Ergodicity of the distribution function
 - We determine first order distribution $F(x) = E\{X(t) \leq x\}$ of a given process $X(t)$ by a suitable time average.
 - Consider the process
$$y(t) = \begin{cases} 1 & \text{if } x(t) \leq x \\ 0 & \text{if } x(t) > x \end{cases}$$
 - Its mean is given by
 - Its autocorrelation $E\{y(t)\} = 1.P\{x(t) \leq x\} = F(x)$
 - Where $F(x, x; \tau)$ is the second order distribution of $x(t)$

$$E\{y(t + \tau)y(t)\} = 1.P\{x(t + \tau) \leq x, x(t) \leq x\} = F(x, x; \tau)$$

IV. Stochastic processes

4.4. Ergodicity

- We form the time average

$$y_T = \frac{1}{2T} \int_{-T}^T y(t) dt$$

- We have

$$E\{y_T\} = E\{y(t)\} = F(x)$$

- The variance of y_T is given by

$$\frac{1}{T} \int_0^{2T} \left(1 - \frac{\tau}{2T}\right) [R(\tau) - \eta^2] d\tau = 0$$

- Where $R(\tau)$ and η are replaced by $F(x, x; \tau)$ and $F(x)$

IV. Stochastic processes

4.4. Ergodicity

- Ergodic theorem for distribution function
 - For a given x ,

$$\lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T y(t) dt = F(x)$$

iff

$$\lim_{T \rightarrow \infty} \frac{1}{T} \int_0^{2T} \left(1 - \frac{\tau}{2T}\right) [F(x, x; \tau) - F^2(x)] d\tau = 0$$



25 YEARS ANNIVERSARY
SOICT

VIỆN CÔNG NGHỆ THÔNG TIN VÀ TRUYỀN THÔNG
SCHOOL OF INFORMATION AND COMMUNICATION TECHNOLOGY

**Thank you for
your attentions!**



soict.hust.edu.vn/



fb.com/groups/soict

