

# Power series

Nguyen Thu Huong



School of Applied Mathematics and Informatics  
Hanoi University of Science and Technology

October 9, 2020

Power series

Nguyen Thu  
Huong

Definition

Theorems on  
power series

Expansion of  
functions into  
power series

## 1 Definition

## 2 Theorems on power series

## 3 Expansion of functions into power series

# Content

1 Definition

2 Theorems on power series

3 Expansion of functions into power series

## Definition

A **power series** (centered at  $x_0$ ) is a **function series** of the form

$$\sum_{n=0}^{\infty} a_n(x - x_0)^n = a_0 + a_1(x - x_0) + \dots + a_n(x - x_0)^n + \dots$$

where  $a_n$  are constants,  $x$  is the variable.

Consider  $x - x_0$  as  $X$ , in the following we consider power series of the form  $\sum_{n=0}^{\infty} a_n x^n$ .

## Example

$$\sum_{n=0}^{\infty} x^n = 1 + x + x^2 + \dots = \frac{1}{1 - x}, \quad |x| < 1.$$

## Power series

Nguyen Thu  
Huong

## Definition

Theorems on  
power seriesExpansion of  
functions into  
power series

## Theorem (Abel Theorem)

*If the series  $\sum_{n=0}^{\infty} a_n x^n$  converges at  $x_0 \neq 0$  then the series converges absolutely at all  $x$  that  $|x| < |x_0|$ .*

*If the series  $\sum_{n=0}^{\infty} a_n x^n$  diverges at  $x_1 \neq 0$  then the series diverges at all  $x$  that  $|x| > |x_1|$ .*

Proof.

## Power series

Nguyen Thu  
Huong

## Definition

Theorems on  
power seriesExpansion of  
functions into  
power series

The series  $\sum_{n=0}^{\infty} a_n x^n$  always converges at  $x = 0$ .

$\exists R > 0$  such that the power series converges absolutely in  $(-R, R)$  and diverges in  $(-\infty, -R) \cup (R, \infty)$ .

At the end points  $x = \pm R$ , the series may converge or diverge.

## Definition

$R$  is called the **radius of convergence** of the series.

$(-R; R)$  is called the **interval of convergence** of the series.

## Theorem

*Radius of convergence* of the series  $\sum_{n=1}^{\infty} a_n x^n$  is determined by

$$R = \lim_{n \rightarrow \infty} \frac{|a_n|}{|a_{n+1}|} \text{ or } R = \lim_{n \rightarrow \infty} \frac{1}{\sqrt[n]{|a_n|}}.$$

## Example

Find the domain of convergence

$$\text{a) } \sum_{n=1}^{\infty} \frac{x^n}{n+2} \quad \text{b) } \sum_{n=1}^{\infty} n! x^n \quad \text{c) } \sum_{n=1}^{\infty} \frac{(-1)^n x^{2n}}{(2n)!}$$

# Content

1 Definition

2 Theorems on power series

3 Expansion of functions into power series



## Proposition

Assume that  $\sum_{n=0}^{\infty} a_n x^n = S(x)$  has the radius of convergence  $R \neq 0$ . Then

- 1  $\sum_{n=0}^{\infty} a_n x^n$  converges uniformly on  $[a; b] \subset (-R; R)$ .
- 2  $S(x)$  is continuous on  $(-R, R)$ .
- 3  $S(x)$  is integrable on  $[a, b] \subset (-R, R)$ .

$$\int \left( \sum_{n=0}^{\infty} a_n x^n \right) dx = \sum_{n=0}^{\infty} \frac{a_n x^{n+1}}{n+1} + C.$$

- 4  $S(x)$  is differentiable on  $(a, b) \subset (-R, R)$ .

$$\left( \sum_{n=0}^{\infty} a_n x^n \right)' = \sum_{n=1}^{\infty} n a_n x^{n-1}.$$

## Remark

These series have the same radius of convergence  $R$ . But their domains of convergence might be different, because of the convergence at the endpoints  $x = \pm R$ .

## Example

Find the sum

$$1 \quad \sum_{n=1}^{\infty} (-1)^n \frac{x^{2n+1}}{2n+1}$$

$$2 \quad \sum_{n=0}^{\infty} (3n+1)x^n.$$

## Example

Find the sum  $\sum_{n=0}^{\infty} \frac{(-1)^n(3n+1)}{8^n}.$

# Content

1 Definition

2 Theorems on power series

3 Expansion of functions into power series

## Definition

Let  $f(x)$  be an infinitely differentiable function at  $x_0$ .  
The **Taylor series** of  $f(x)$  at  $x_0$  is the series

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n.$$

If  $x_0 = 0$ , the series

$$\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n$$

is called the **Maclaurin series** of  $f(x)$ .

## Power series

Nguyen Thu  
Huong

## Definition

Theorems on  
power seriesExpansion of  
functions into  
power series

## Example

Consider  $f(x) = \begin{cases} e^{-\frac{1}{x^2}} & \text{if } x \neq 0 \\ 0 & \text{if } x = 0. \end{cases}$

$f(x)$  has derivatives of all orders and  $f^{(n)}(0) = 0$ , the Taylor series of  $f(x)$  is 0.

## Remark

The Taylor series of  $f(x)$  at  $x_0$  may converge or diverge. In case it converges, the sum may not equal  $f(x)$ .

## Theorem

Let  $f(x)$  have the derivatives of all orders in  $I = (x_0 - R; x_0 + R)$ . If there is  $M > 0$  such that  $|f^{(n)}(x)| \leq M$  for all  $x \in I$ ,  $n \in \mathbb{N}$ . Then the Taylor series

$\sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$  converges to  $f(x)$  in  $(x_0 - R; x_0 + R)$ .

## Example

Expand  $f(x) = e^x$  into power series.

## Power series

Nguyen Thu  
Huong

## Definition

Theorems on  
power seriesExpansion of  
functions into  
power series

- $e^x = \sum_{n=0}^{\infty} \frac{x^n}{n!}, x \in \mathbb{R}.$
- $\sin x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n+1}}{(2n+1)!}, x \in \mathbb{R}.$
- $\cos x = \sum_{n=0}^{\infty} (-1)^n \frac{x^{2n}}{(2n)!}, x \in \mathbb{R}.$
- $\frac{1}{1-x} = \sum_{n=0}^{\infty} x^n, |x| < 1.$
- $(1+x)^\alpha = \sum_{n=0}^{\infty} \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!} x^n, |x| < 1.$

## Power series

Nguyen Thu  
Huong

Definition

Theorems on  
power seriesExpansion of  
functions into  
power series

## Example

Expand the following functions into Maclaurin series

$$1 \quad f(x) = \frac{1}{x^2 - 3x + 2}.$$

$$2 \quad f(x) = \ln(1 + x).$$

$$3 \quad f(x) = \arctan x.$$

$$4 \quad f(x) = \frac{1}{(1 - x)^2}.$$

## Example

Expand  $f(x) = \ln x$  into Taylor series near  $x = 1$ .