

Implicit functions

Nguyen Thu Huong



School of Applied Mathematics and Informatics
Hanoi University of Science and Technology

January 21, 2021

Content

Implicit
functions

Higher order
partial
derivatives
and total
differentials

Taylor's
theorem

1 Implicit functions

2 Higher order partial derivatives and total differentials


3 Taylor's theorem

Content

- 1 Implicit functions
- 2 Higher order partial derivatives and total differentials
- 3 Taylor's theorem

Implicit function $y = y(x)$ given by $f(x, y) = 0$

- Given $x^2 + y^2 - 1 = 0$, we solve for $y(x) = \pm\sqrt{1 - x^2}$.
- Given the equation $x^3 + y^3 - 3xy = 0$.



For all $x \in [0, \sqrt[3]{2}]$, the equation determines three y such that $f(x, y) = 0$. Hence, the equation $f(x, y) = 0$ determines one or more implicit functions $y = y(x)$.

Derivative of the function $y = y(x)$ given by $f(x, y) = 0$

Implicit
functions

Higher order
partial
derivatives
and total
differentials

Taylor's
theorem

Theorem (Existence)

Let $f(x, y): D \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ have continuous partial derivatives in a neighborhood of $M_0(x_0, y_0)$, $f(M_0) = 0$ and $f'_y(M_0) \neq 0$.

Then, the equation $f(x, y) = 0$ determines in a neighborhood of x_0 a continuously differentiable implicit function $y = y(x)$ that $y(x_0) = y_0$. Moreover,

$$y'(x) = -\frac{f'_x(x, y)}{f'_y(x, y)}.$$

It holds $f(x, y) = 0 \Leftrightarrow y = y(x)$, or $f(x, y(x)) = 0$.

Implicit function $z = z(x, y)$ determined by $f(x, y, z) = 0$

Implicit
functions

Higher order
partial
derivatives
and total
differentials

Taylor's
theorem

Theorem (Existence)

Let $f(x, y, z): D \subset \mathbb{R}^3 \rightarrow \mathbb{R}$ have continuous partial derivatives in a neighborhood of $M_0(x_0, y_0, z_0)$, $f(M_0) = 0$ and $f'_z(M_0) \neq 0$. Then, the equation $f(x, y, z) = 0$ determines in a neighborhood of (x_0, y_0) an implicit function $z = z(x, y)$ such that $z(x_0, y_0) = z_0$ and z has continuous partial derivatives. Moreover,

$$z'_x(x, y) = -\frac{f'_x(M)}{f'_z(M)}, \quad z'_y(x, y) = -\frac{f'_y(M)}{f'_z(M)}.$$

Example

- 1 Assume that the function $y = y(x)$ is given by the equation $xe^y + x^3 + 2x^2y - 2 = 0$. Compute $y'(1)$.
- 2 Let $z = z(x, y)$ be determined by the equation $2x^3y - x^2 + xe^z + 2xyz = 2$. Compute $z(1; 1)$, $z'_x(1; 1)$, $z'_y(1; 1)$.
- 3 Let $y = y(x)$ be determined by the equation $xe^y + 2x^3 - 2x^2y + 3 = 0$. Approximate $y(-1, 01)$.
- 4 Let $z = z(x, y)$ be determined by the equation $z^3 + x^2y + xy^2 = 3xyz$.
 - a) Compute $z(1; -1)$, $z'_x(1; -1)$, $z'_y(1; -1)$.
 - b) Approximate $z(0, 99; -1, 01)$.

Theorem

Let $f(M_0) = g(M_0) = 0$, where f, g have continuous partial derivatives near $M_0(x_0, y_0, z_0, u_0, v_0)$ and

$$\frac{D(f, g)}{D(u, v)}(M_0) = \begin{vmatrix} f'_u & f'_v \\ g'_u & g'_v \end{vmatrix}(M_0) \neq 0.$$

Then, the system $\begin{cases} f(x, y, z, u, v) = 0 \\ g(x, y, z, u, v) = 0 \end{cases}$ determines two implicit functions $u = u(x, y, z)$, $v = v(x, y, z)$ near (x_0, y_0, z_0) such that $u(x_0, y_0, z_0) = u_0$, $v(x_0, y_0, z_0) = v_0$. The functions u, v have continuous partial derivatives.

Differentiating w.r.t. x both sides of the equations of the system

$$\begin{cases} f'_x + f'_u \cdot u'_x + f'_v \cdot v'_x = 0 \\ g'_x + g'_u \cdot u'_x + g'_v \cdot v'_x = 0 \end{cases}$$

We obtain

$$u'_x = -\frac{\frac{D(f,g)}{D(x,v)}}{\frac{D(f,g)}{D(u,v)}}, \quad v'_x = -\frac{\frac{D(f,g)}{D(u,x)}}{\frac{D(f,g)}{D(u,v)}}.$$

Similarly, we can compute u'_y, u'_z, v'_y, v'_z .

Content

- 1 Implicit functions
- 2 Higher order partial derivatives and total differentials
- 3 Taylor's theorem

Definition

Second order partial derivatives of $f(x, y)$ are:

$$f''_{xx} = \frac{\partial^2 f}{\partial x^2} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial x} \right)$$

$$f''_{yy} = \frac{\partial^2 f}{\partial y^2} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial y} \right)$$

$$f''_{xy} = \frac{\partial^2 f}{\partial y \partial x} = \frac{\partial}{\partial y} \left(\frac{\partial f}{\partial x} \right)$$

$$f''_{yx} = \frac{\partial^2 f}{\partial x \partial y} = \frac{\partial}{\partial x} \left(\frac{\partial f}{\partial y} \right)$$

Example

- 1 Compute all the second order partial derivatives of the function $z = \arctan(x^2y)$.
- 2 Compute the second order partial derivatives at $(0,0)$ of the function $f(x,y) = \begin{cases} xy \frac{x^2 - y^2}{x^2 + y^2} & \text{if } (x,y) \neq (0,0), \\ 0 & \text{if } (x,y) = (0,0). \end{cases}$

Schwarz theorem

Implicit
functions

Higher order
partial
derivatives
and total
differentials

Taylor's
theorem

Theorem

Assume that $z(x, y)$ has partial derivatives f''_{xy} và f''_{yx} near $M_0(x_0, y_0)$ which are continuous at M_0 . Then,

$$f''_{xy}(M_0) = f''_{yx}(M_0).$$

Higher order total differentials

Definition

Higher order total differential of $z = f(x, y)$:

$$d^n z = d(d^{n-1} z), \quad n \geq 2.$$

If x, y are independent variables, then $d^n z = \left(\frac{\partial}{\partial x} dx + \frac{\partial}{\partial y} dy \right)^n f$

Example

Second order total differential

$$d^2 z = \left(\frac{\partial}{\partial x} dx + \frac{\partial}{\partial y} dy \right)^2 f = \frac{\partial^2 f}{\partial x^2} (dx)^2 + 2 \frac{\partial^2 f}{\partial x \partial y} dx dy + \frac{\partial^2 f}{\partial y^2} (dy)^2.$$

Note: Higher order total differentials do not fulfill the invariance property.

Example

- 1 Given $z = 2^x \sin y$. Compute the second order differential d^2z and $d^2z(0, \pi)$.
- 2 Given $z = \arctan(xy)$. Compute the second order differential d^2z and $d^2z(1, -1)$.

Content

- 1 Implicit functions
- 2 Higher order partial derivatives and total differentials
- 3 Taylor's theorem**

Taylor theorem for multivariable functions

Theorem

Let $f(x, y)$ have continuous partial derivatives up to order $n + 1$ in an open set $U \subset \mathbb{R}^2$, $M_0(x_0, y_0), M(x_0 + \Delta x, y + \Delta y) \in U$. There exists $0 < \theta < 1$ such that

$$\begin{aligned} f(x_0 + \Delta x, y + \Delta y) - f(x_0, y_0) = & df(x_0, y_0) + \frac{1}{2!} d^2 f(x_0, y_0) + \dots \\ & + \frac{1}{n!} d^n f(x_0, y_0) + \frac{1}{(n+1)!} d^{n+1} f(x_0 + \theta \Delta x, y + \theta \Delta y). \end{aligned}$$

Example

Compute the second degree Taylor polynomial of the function $f(x, y) = e^{xy}$ at $(0, 1)$.