ANOVA a Misure Ripetute

Effetti Entro i Sggetti

	Correzione della Sfericità	Somma dei Quadrati	gdl	Media Quadratica	F	p	η^2	η^2_p
Training	Nessuna	4.903	1	4.9031	87.67	<.001	0.334	0.907
	Greenhouse-Geisser	4.903	1.00	4.9031	87.67	<.001	0.334	0.907
Residuo	Nessuna	0.503	9	0.0559				
	Greenhouse-Geisser	0.503	9.00	0.0559				
Blocco	Nessuna	1.874	2	0.9372	12.72	<.001	0.128	0.586
	Greenhouse-Geisser	1.874	1.75	1.0741	12.72	<.001	0.128	0.586
Residuo	Nessuna	1.327	18	0.0737				
	Greenhouse-Geisser	1.327	15.71	0.0845				
Training ✻ Blocco	Nessuna	1.523	2	0.7617	9.97	.001	0.104	0.525
	Greenhouse-Geisser	1.523	1.13	1.3507	9.97	.009	0.104	0.525
Residuo	Nessuna	1.376	18	0.0764				
	Greenhouse-Geisser	1.376	10.15	0.1355				

Nota. Somma dei quadrati Tipo 3

Effetti Tra Soggetti

	Somma dei Quadrati	gdl	Media Quadratica	F	p	η^2	η^2_p
Residuo	3.19	9	0.355				

Nota. Somma dei quadrati Tipo 3

Ipotesi

Test di Sfericità

	W di Mauchly	p	Greenhouse-Geisser ε	Huynh-Feldt ε
Training[aaa]	1.000	NaN	1.000	1.000
Blocco	0.854	.532	0.873	1.000
Training ✻ Blocco	0.227	.003	0.564	0.589

[a] Le misure ripetute hanno solo due livelli. L'assunzione di sfericità è sempre soddisfatta quando le misure ripetute hanno solo due livelli.

Test Post Hoc

Confronti Post Hoc - Training

Confronto											
Training		Training	Differenza Media	SE	gdl	t	p	p_{tukey}	$p_{scheffe}$	$p_{bonferroni}$	p_{holm}
T1	-	T2	0.572	0.0611	9.00	9.36	<.001	<.001	<.001	<.001	<.001

Confronti Post Hoc - Blocco

Confronto											
Blocco		Blocco	Differenza Media	SE	gdl	t	p	p_{tukey}	$p_{scheffe}$	$p_{bonferroni}$	p_{holm}
B1	-	B2	0.3474	0.0986	9.00	3.525	.006	.016	.020	.019	.013
	-	B3	0.3974	0.0869	9.00	4.575	.001	.003	.004	.004	.004
B2	-	B3	0.0500	0.0696	9.00	0.719	.491	.759	.778	1.000	.491

Confronti Post Hoc - Training ✻ Blocco

Confronto					Differenza Media	SE	gdl	t	p	p_{tukey}	$p_{scheffe}$	$p_{bonferroni}$	p_{holm}
Training	Blocco		Training	Blocco									
T1	B1	-	T1	B2	0.6279	0.1898	9.00	3.308	.009	.071	.145	.137	.073
		-	T1	B3	0.7727	0.1819	9.00	4.248	.002	.019	.046	.032	.021
		-	T2	B1	1.0089	0.1689	9.00	5.972	<.001	.002	.006	.003	.003
		-	T2	B2	1.0759	0.1402	9.00	7.674	<.001	<.001	<.001	<.001	<.001
		-	T2	B3	1.0310	0.1496	9.00	6.891	<.001	<.001	.002	.001	<.001
	B2	-	T1	B3	0.1449	0.0702	9.00	2.064	.069	.380	.547	1.000	.276
		-	T2	B1	0.3810	0.1116	9.00	3.413	.008	.061	.128	.116	.069
		-	T2	B2	0.4480	0.0742	9.00	6.038	<.001	.002	.005	.003	.003
		-	T2	B3	0.4032	0.1295	9.00	3.112	.012	.094	.183	.187	.087
	B3	-	T2	B1	0.2362	0.0795	9.00	2.970	.016	.115	.216	.235	.094
		-	T2	B2	0.3031	0.0653	9.00	4.642	.001	.011	.028	.018	.013
		-	T2	B3	0.2583	0.0878	9.00	2.941	.016	.119	.224	.247	.094
T2	B1	-	T2	B2	0.0670	0.0691	9.00	0.969	.358	.917	.960	1.000	1.000
		-	T2	B3	0.0221	0.0681	9.00	0.325	.753	.999	1.000	1.000	1.000
	B2	-	T2	B3	-0.0448	0.0814	9.00	-0.551	.595	.992	.997	1.000	1.000